UNIV. OF
TORONTO
LIBRARY

2354

NATURE AND DEVELOPMENT OF PLANTS

CURTIS

I

NATURE AND DEVELOPMENT OF PLANTS

UNIVERSITY OF TORONTO LIBRARY

BY

CARLTON C. CURTIS, A.M., PH.D.

Instructor in Botany in Columbia University

SEEN BY PRESERVATION SERVICES
DATE DEC 1 2 1991

NEW YORK
HENRY HOLT AND COMPANY
1907

105-960
1/11/10

Copyrighted 1907, by

HENRY HOLT AND COMPANY

PRESS OF
THE NEW ERA PRINTING COMPANY
LANCASTER PA.

PREFACE

In the present volume an attempt has been made to acquaint the reader with the more essential aspects of the subject. Especially have we had in mind a purpose to make familiar our common plants—which knowledge we believe to be fundamental in any botanical work. In a word the work is designed to give the reader a view point. It is not put forth as a text-book but it is hoped that the discussion will give the student such a comprehension of the subject that he will come to the lecture room in a proper attitude and that he will approach his laboratory work with a desire for investigation. If the book shall quicken the reasoning faculty, and create a desire for further examination of the subject, its object will have been fulfilled.

Acknowledgments are made to Professors L. M. Underwood, H. M. Richards, T. E. Hazen and to Doctors G. G. Scott and Chester A. Darling, for valuable assistance in the preparation of various portions of the text. The author desires to express his especial indebtedness to his good friend, Mr. H. O. Hanson, for the preparation of many of the illustrations, to Mr. W. A. Andrews for microscopical studies from which drawings have been made and to Miss Mary L. Williams for the execution of many of the drawings.

CARLTON C. CURTIS.

COLUMBIA UNIVERSITY,
August, 1907.

CONTENTS

PAGE

INTRODUCTION 1

PART I

NATURE OF PLANTS

I. THE LEAF 7
II. THE ROOT 42
III. THE STEM 62
IV. THE FLOWER, FRUIT AND SEEDLING 101

PART II

THE DEVELOPMENT OF PLANTS

V. CLASSIFICATION OF PLANTS 130
VI. THALLOPHYTA 134
VII. BRYOPHYTA 242
VIII. PTERIDOPHYTA 282
IX. SPERMATOPHYTA 320
X. ANGIOSPERMAE (SPERMATOPHYTA CONCLUDED) 345

NATURE AND DEVELOPMENT OF PLANTS

INTRODUCTION

1. The Nature of the Plant.—It is a familiar fact that most plants have stems, leaves and roots and that these organs are variously modified. Why have such organs been developed? Why do they assume such varied forms and arrangements? How does it come about that the stem reaches up in the air and that the branches and leaves are arranged in a very definite order while the root penetrates the soil and grows toward moisture and other soil material? Is a plant, like an animal, conscious of its surroundings, seeking suitable foods and avoiding unfavorable conditions? It is evident to the casual observer that the plant is sensitive to its surroundings; some plant organs bend toward a light that we recognize with difficulty or react to a touch that we cannot appreciate. Has the plant a nervous system and faculties like ourselves that enable it to adjust itself to its surroundings? To answer these questions it will be necessary to



FIG. 1. Diagram of a cell showing its three dimensions; mesh-work of granular cytoplasm which incloses colorless cell sap, the vacuoles, and a denser dark body, the nucleus.

consider the nature of the living substance of the plant. The plant body is composed of cells that are like boxes of living matter having length, breadth, and thickness (Fig. 1). While the cells assume various forms and perform different work or functions, they all possess cell walls which contain during the

life of the cell a substance termed protoplasm. This substance resembles somewhat the white of an egg, being viscid and rather tenacious. The protoplasm is not of uniform consistency. The larger portion of it is finely granular and is called the cytoplasm while a denser rounded part is termed the nucleus. Other bodies, plastids, also denser than the cytoplasm are of common occurrence. Such plastids as contain a green pigment, chlorophyll, are called choroplastids or chloroplasts, and these produce the green color

FIG. 2. Structure of the cell: *A*, cell of ditch-moss, *Philotria; n*, nucleus, *v*, vacuole, *ch*, chloroplast. *B*, cell of carrot; *ch*, chromoplasts, *n*, nucleus. *C*, chromoplasts from rose hip. *D*, cells of *Begonia; s*, starch grains, *c*, crystals of lime. *E*, leucoplast, *l*, of potato forming a starch grain.

of the vegetation (Fig. 2, *A*). Still other plastids, the chromoplasts, contain red or yellow pigments that give the color to many fruits and flowers, as the tomato, rose hip, squash, nasturtium, etc. (Fig. 2, *B, C*). Colorless plastids, leucoplasts, often occur in cells hidden from the light. One group of these leucoplasts forms the starch that appears in underground storage organs as

in the potato (Fig. 2, *E*). Usually spaces or vacuoles, that appear as cavities also occur in the granular cytoplasm (Fig. 2, *A*, *v*). In reality they are filled with watery solutions of various substances, the so-called cell sap. Many other structures (Fig. 2, *D*) appear in the protoplasm to which attention will be called later. It is well to remember that these visible structures may not represent the real composition of the protoplasm. It has been supposed that a variety of units beyond the range of visibility, grow, multiply, and build up the various structures which we recognize as constituting the protoplasm. The cell walls make up the body of the plant and give stability to its various organs but the living part of the cell is the protoplasm. This substance reaches through delicate pores in the cell wall to the protoplasm of adjacent cells so that all the living substance of the entire plant body is in contact and forms one united mass.

2. The Nature of the Living Substance of the Plant.—The protoplasm possesses most remarkable powers. It can absorb various fluids and gases, decompose them into simple elements, reunite them into foods or discharge from the cell such substances as are not required. Furthermore the protoplasm effects those changes termed growth by transforming the foods into the substances that compose the cell walls and other parts of the cell, as the protoplasm itself. How are these changes brought about? Every substance is composed of elements. Water consists of two elements, hydrogen and oxygen. The elements that compose any substances are held together with great energy, owing to their mutual attraction for one another. When two different substances are brought together, it may happen that the attraction of certain elements of one substance is greater for one or more of the elements of the other substance than for its own elements. The result is, that the elements will be torn away from their respective substances and united into new combinations. We say that a decomposition and a re-combination has been effected, or, that a chemical reaction has occurred. We see an illustration of these chemical changes when iron-ore is heated with charcoal. The ore is composed of the elements iron and oxygen and the charcoal consists of carbon. In the presence of

heat the carbon has a stronger attraction for the oxygen than the iron has. Consequently the oxygen is drawn away from the iron and unites with the carbon, forming a new combination of carbon and oxygen, and leaving the iron free. This decomposition and re-combination is only a part of the chemical change that takes place, but it will serve to illustrate the nature of many chemical reactions. The energy that was required to hold the oxygen and iron together begins to be set free as soon as the decomposition starts and it contributes to the rise of temperature during the reaction. The living substance of the plant is composed of a great variety of chemical substances that are very readily decomposed by gases and other substances which it absorbs. Light, heat, gravitation and moisture are even more powerful in producing these changes. These substances and forces are often referred to as stimuli (sing. stimulus) because they start the chemical changes referred to above. So the various substances and forces in nature stimulate or cause chemical changes in the living substance of the plant and the energy thus set free is used by the protoplasm in the performance of its work. The cells of a seed have no power of their own to grow or perform any duty. It is not until a certain amount of heat, moisture, gases or other stimulating forces have acted upon the protoplasm or substances contained in it and so aroused chemical changes that it is furnished with the requisite energy to begin growth. Furthermore, these forces exert a very definite influence upon the protoplasm and cause it to accomplish very definite results. This is due to the fact that not all portions of the protoplasm of the plant body are equally influenced by, or we may say equally sensitive to heat, light, and other forces and consequently the various parts of the plants do not respond alike. The force of gravity, for example, acts upon some stems and the living substance is influenced by this stimulus so that a growth is aroused that bends the stem into an upright position. The protoplasm in the cells of the root, however, is so constituted that the stimulus of gravity causes a growth that bends the root down into the soil. Light stimulates the mustard plant in such a way

that the stem grows toward the light and the root away from it So the living substance of the plant while forming one united whole varies in its sensitiveness in various parts of the plant. In this respect the various parts may be compared to the receiving stations of a wireless telegraphy system. The receiving stations may have instruments sensitive to certain intensities of electrical currents and other currents do not affect them. So the various organs of the plant are sensitive, each in its own peculiar way, to certain stimuli. As a result of this adjustment the various parts of the plant, being attuned or sensitive to certain forces, are led or induced to grow so that they come into the most helpful and beneficial relation to these forces. Thus the leaves, branches, and roots are stimulated to grow and develop so that each part becomes properly related to light, moisture, gravity, etc. The broad blades of the majority of leaves are exposed to the direct rays of light. This arrangement is of the greatest benefit because they are using the light in the manufacture of foods. This position is assumed, however, because they are sensitive to light, gravitation and other forces which direct and cause the arrangement. This is the most noteworthy feature about the sensitiveness of the protoplasm. Forces stimulate to growth in such a way that the results are helpful and the greatest good comes to the plant. In other words the reactions are purposive. These growths are so elaborate and beneficial that they often appear as the result of reason and will. The bending of the root into the soil brings it into contact with water and other foods; the tendril coils about a branch and binds the plant firmly to the support. The plant, however, does not direct these adjustments. These reactions do not involve consciousness. These movements and all others are absolutely directed and controlled by the various forces which act upon the sensitive protoplasm. A definite reaction follows a stimulation which the plant has no power to alter or control. The root will bend down into a dish of mercury with the same energy and directness as that with which it penetrates the soil, and the tendril will clasp your finger or a branch of its own body as firmly as a serviceable support.

We will now proceed to study the plant from this standpoint, giving attention to the fitness or adaptation of the leaf, root and stem, to the conditions under which the plant lives, and to the work which it performs.

PART I

THE NATURE OF PLANTS

CHAPTER I

THE LEAF

3. The Work of the Leaf.—Each plant has a character or a personality of its own. What is it that gives this individuality to the plant? When the work that it performs is understood it will be seen that this character is very largely due to the adaptation of the leaves to the performance of certain duties. The



FIG. 3. Forms of leaves: *A*, leaf of white birch with netted veins—*p*, petiole; *b*, blade. *B*, leaf of Solomon's seal with parallel venation and blade clasping stem without petiole.

leaves bring the plant into harmony with its surroundings and give to it a subtle individuality owing to the perfection of their arrangements, structures and forms for the work in hand. The

more important work performed by the leaves is the construction of foods, the giving off of water or transpiration, and breathing or respiration. The magnitude of this work far exceeds the energy expended in all the industries of the world. The leaves, however, accomplish this work so quietly and economically that most people are scarcely conscious of it.

In order to understand why the leaf has a petiole and green blade (Fig. 3) or why it assumes various forms and arrangements it will be necessary to examine the structure of the leaf and see the character of the apparatus that is used in the performance of its work. The blades of the majority of leaves are flat. A



FIG. 4. Blade of lilac leaf cut across showing the more compact arrangement of cells upon the upper side of the leaf.

section through such a blade, cut as in Fig. 4 so that we can look into the end of the blade, shows that the leaf is composed of a complicated arrangement of cells. Fig. 5 is a greatly enlarged view of Fig. 4 taken at *X*.

4. The Epidermis.—It is now seen that a layer of compact cells (Fig. 5, *e*) surrounds the leaf on all sides. In fact such a layer covers all parts of the plant body. This layer of cells, the epidermis, is provided with minute openings or stomata (sing. stoma), Fig. 5, *s*. The stomata are especially abundant in the epidermis on the under surface of the leaf and often quite lacking from the upper surface. A better idea of these openings and of the epidermal cells may be gained by stripping off the epidermis from



FIG. 5. Greatly enlarged view of region *X* in Fig. 4: *e*, epidermis; *s*, stoma; *p*, palisade mesophyll; *ch*, chloroplast; *sp*, spongy mesophyll; *i*, intercellular spaces; *v*, small vein cut across; *l*, end of vein seen from the side, consisting of elongated and banded cells (tracheids); *col*, collecting cells that transfer material to and from the palisade cells.—H. O. Hanson.



FIG. 6. Surface view of epidermis of lilac leaf: *g*, guard cell; *s*, stoma; *ch*, chloroplasts; *n*, nucleus.—H. O. Hanson.

such leaves as the live-for-ever, blue flag or hyacinth. The cells of the epidermis are very compactly put together and may be stripped off by means of a penknife as a thin white skin. Examined under a microscope, such preparations show the stomata as minute openings guarded by two rather lens-shaped cells, termed the guard cells (Fig. 6). Chloroplasts usually occur only in the guard cells, the other epidermal cells having a watery, colorless cell content. In addition to these features, Fig. 7 shows that the outer portion of the walls of the epidermal cells is modified so as to form a delicate, skin-like layer over the outer sur-



FIG. 7.



FIG. 8.

FIG. 7. Section across a stoma shown in Fig. 6: *s*, stoma; *g*, guard cell; *c*, cuticle; *a*, air chamber.—H. O. Hanson.

FIG. 8. Hairs from the epidermis of squash leaf.

face of the epidermis. This part of the cell wall is called the cuticle. It contains a substance, cutin, that renders the epidermis nearly impervious to gases and fluids, so that the leaf is entirely surrounded by a water- and gas-proof coat. In many leaves hairs of various forms grow out from the epidermal cells (Fig. 8).

5. The Mesophyll.—The cells within the epidermis have, for the most part, a greenish color due to numerous green bodies, chloroplasts, contained in the cells (Fig. 5, *ch*). These green cells of the leaf, termed the mesophyll, consist of one or more layers of rather elongated and compact cells on the upper side of the leaf (the palisade mesophyll, Fig. 5, *p*), and below they are joined to rather irregular and loosely arranged cells (the

spongy mesophyll, Fig. 5, *sp*). Numerous air spaces extend in all directions through this spongy mass of cells reaching up to and between the palisade cells and down to the stomata (Fig. 5, *i*). Bundles of small cells also appear here and there between the palisade and spongy mesophyll (Fig. 5, *v*, *l*). These structures are the so-called veins that appear as fine lines in the majority of leaves (Fig. 3). These veins or vascular bundles branch again and again and so extend to all parts of the leaf. In Fig. 5, *v* one of the veins has been cut square across while at *l* a very small one has been cut in half lengthwise. The walls of the mesophyll cells are smooth, thin, and elastic and composed of a substance called cellulose. Thin places or pores are often formed in the walls which appear as openings owing to the extreme thinness of the wall at such places. The term parenchyma is applied to all cells possessing the character of walls noted in the mesophyll cells. In the vascular bundle, however, a portion of the cells have thicker walls which are more rigid, hard and woody. Such cells are often referred to as prosenchyma.

What is the significance of the peculiar and varied arrangements of the cells in the leaf? We naturally anticipate that these different structures are adapted to performance of special duties. Attention will first be directed to the work performed by the green portion of the leaf, the mesophyll. Similar cells containing chloroplasts are of common occurrence in various parts of the plant, being often found in stems and unripe fruits. In the leaf such a tissue is termed mesophyll but a more general term, Chlorenchyma, includes all chlorophyll-bearing tissue wherever found.

6. The First Function of the Leaf, Photosynthesis.—Chlorenchyma is the most important tissue in the plant since it has for its special function or work the construction of carbohydrates, such as sugars and starches, which are among the most important foods of the plant. The construction of these carbohydrates by the chlorenchyma is called photo-synthesis. This work easily ranks as the leading wonder of the world because its processes are so complicated and of such magnitude and importance in the economy of this world of ours. It will be seen that this tissue not only keeps the air in a wholesome condition for breathing

and so makes life possible upon the earth but that all animal life is absolutely dependent, either directly or indirectly, for food upon the products built up by the chlorenchyma. Let us see how these foods are formed. Various substances are absorbed by the roots, and the vascular bundles, which extend throughout the plant body, transfer them to the chlorenchyma. This absorbed material is in large part water, but various mineral substances are dissolved in it. Various gases, as carbon dioxide, also find their way through the stoma to the chlorenchyma. All of these absorbed substances are composed of elements. Thus water consists of two elements, hydrogen and oxygen, in the proportion of two parts of hydrogen to one part oxygen. This composition is indicated by the abbreviation, H_2O. So the gas, carbon dioxide, contains one part of carbon and two parts of oxygen, indicated thus CO_2. Chloroplasts have the power under suitable conditions of decomposing or tearing apart these compounds. They are able with a suitable light and temperature to separate CO_2 into the elements carbon and oxygen and H_2O into hydrogen and oxygen. Furthermore they can reunite these elements into new compounds that are quite different from the original ones. In this way CO_2 and H_2O are decomposed and the elements reunited into complex compounds or foods consisting of carbon, hydrogen and oxygen. Such complex foods as cane sugar, consisting of twelve parts of carbon, twenty-two of hydrogen, and eleven of oxygen are formed in this way. This very common food is expressed by the formula $C_{12}H_{22}O_{11}$. So also other foods are formed, as grape sugar $C_6H_{12}O_6$ and starch usually indicated as $C_6H_{10}O_5$. The cellulose walls of the parenchyma cells have the same composition as starch. It will be seen that all these foods have carbon united to hydrogen and oxygen in the same proportion as it exists in water. Cane sugar has carbon united to eleven times H_2O, and starch five times H_2O. It was supposed that these substances were a combination of carbon and H_2O. For this reason they were called carbohydrates. This name is still applied to these foods although it is known that the base is not H_2O but the hydroxyl OH. The actual changes that are effected in the formation of these various carbohydrates is

not known. It has been suggested that CO_2 and H_2O are at first decomposed into simpler forms and finally re-combined into a compound called formaldehyde (HCOH, written also CH_2O). This change is represented by the equation, $CO_2 + H_2O = CH_2O + O_2$; the two parts of oxygen passing from the cells and escaping from the stomata as a gas. In the presence of certain alkalies and acids, formaldehyde increases the number of elements which compose it, thus six times CH_2O would give $C_6H_{12}O_6$ or grape sugar. This is but a theory based upon the fact that CO_2 and H_2O do split up into compounds which may unite to form CH_2O. Formaldehyde can be detected in plant hairs and it is possible to produce sugar from this substance by treating it with alkalies and acids. All that can be definitely stated about the changes going on in the leaf, during the formation of the carbohydrates relates solely to the beginning and end of the process. Water and carbon-dioxide enter the chlorenchyma cells. A series of changes follows, the nature of which can only be conjectured. Finally, as the end result, sugar appears in the cells and oxygen is set free, escaping as a gas.

These complex changes are brought about in the leaf in so subtle a way that we are not conscious of them or of the great amount of energy that is required to effect them. It requires the energy expressed by a temperature of 1300° C. to decompose CO_2 into its elements. Where does the plant obtain the energy to bring about the decompositions and recomposition? This work is accomplished by the energy of the sunlight acting upon the chloroplasts. The chloroplasts of the seed plants are minute, rather lens-shaped grains, and increase in number by the division of the plastid into two equal parts (Fig. 9). These bodies are denser portions of the cytoplasm and like it are nearly colorless. In the presence of light a green, oily substance, chlorophyll, is formed in the plastids, thus producing their green color. Chlorophyll is rarely formed except in the presence of light. Plants grown in cellars or in the dark are of a pale color owing to the absence of chlorophyll in the plastids. Similarly green plants lose their chlorophyll when placed in the dark. This is the principle employed in the blanching of celery. The stalks are

surrounded by earth or the light is excluded from them by some other means; chlorophyll disappears from the plastids and fails to develop in the new cells that are formed in the dark. Plants in which the chlorophyll is not developed are said to be etiolated. The plastids are in the cells, however, for if an etiolated seedling is removed from the dark to the light the chlorophyll will begin to appear in a few hours. The greening of potatoes when removed from the ground and left in a strong light is a familiar example of this. It is evident that the formation of chlorophyll is closely connected with the action of light. Another important point regarding the influence of light is the fact that such a food as starch is not usually found in the leaf in the absence of light. This may be easily verified by cutting the initials of your name in rather large letters from a piece of black paper or smooth tin-foil and fastening the letters in the early morning with mucilage to the upper surface of a well-sunned leaf of a starch forming plant. Before the sunlight fades remove the leaf from the plant and place it in alcohol. This will dissolve the chlorophyll and the leaf will become quite white in a few hours. Now place the leaf in a tincture of iodine which turns starch to a blue or blue-black color. Wherever the light was excluded from the leaf by the letters there is only a pale yellow color but all portions of the sunned leaf show an abundance of starch as revealed by the blue or blue-black coloration. No starch can be detected in leaves taken from plants that have been growing in the dark for twenty-four hours. Plastids that do not contain chlorophyll, examples of which are found in the cells forming the white bands or blotches in many variegated plants, do not form starch in the light. It is evident from these facts that the chlorophyll and light co-operate in some way in the formation of starch. It is known that chlorophyll absorbs a certain portion of the sunlight. If a beam of sunlight is caused to pass through a prism it is separated into seven colors, *i. e.,* red, orange, yellow, green, blue, indigo and violet. When, however, a beam first passes through an alcoholic solution of chlorophyll or through a green leaf a portion of certain colors and all of other colors do not appear. The chlorophyll has taken

up or absorbed certain of the light rays especially portions of the red, blue and violet. It is reasonable to suppose that the energy in these rays is utilized in part by the plastids in effecting the changes noted above in the formation of foods. The sun gives to the earth great quantities of energy in the form of light. A portion of this light is taken up by the chlorophyll, and the plastid which is the living substance containing the chlorophyll, uses the energy of the light to bring about the decomposition of CO_2 and H_2O and the formation of sugars, starches, and other carbohydrates. Because of the importance of the sunlight in the formation of carbohydrates this process is called photosynthesis. This term means the putting together by means of light.

7. The Magnitude of the Work of Photosynthesis.—Carbon is an important element in the composition of plant foods and also in the walls of the cells. It forms one half of the dry weight of the plant. All the carbon appearing in the plant is derived solely from the CO_2 in the air, of which it forms a very small part, only three or four parts in ten thousand. Furthermore the carbon comprises only 3/11 by weight of the CO_2, 8/11 being oxygen. A square meter of leaf surface, however, can withdraw all the CO_2 from 1000 liters of air in one hour. This would furnish sufficient carbon for the construction of one gram of starch. In this way such large quantities of CO_2 are drawn into the chlorenchyma cells as to make possible each year the harvest and the renewal of vegetation of the earth. In the United States alone in this way there was in 1906 built up over 2700 millions of bushels of corn, 770 million bushels of wheat, 850 million bushels of oats, etc., representing a value for all crops of seven million dollars. These figures represent only a small portion of the work performed, since the grain is but a part of the plant, and furthermore the larger portion of the land is covered with forests and other forms of vegetation. The fact must not be overlooked that this process of photosynthesis is of vital importance to our welfare in another way. Owing to the large volumes of CO_2 that are constantly formed by fires and the respiration of animals such an excess of this gas would accumulate in the atmosphere that all animal life would soon cease were it not for its absorption in

photosynthesis. So nearly balanced are the rates of formation and absorption, however, that the percentage existing in the air does not materially vary.

8. The Construction of Proteids.—A second group of foods formed by plants are called proteids. These differ from the carbohydrates in that they contain nitrogen in addition to carbon, hydrogen and oxygen. They are more complex compounds than the carbohydrates and may contain sulphur and phosphorus in addition to the four elements mentioned above. Less is known about their formation than of the carbohydrates. It is probable that they are largely formed in the leaves and by a process similar to that of photosynthesis. The sulphates, phosphates, and nitrates absorbed from the soil are decomposed and the elements of nitrogen, sulphur or phosphorus are united to simple carbon compounds and complex proteids are the result. Light does not co-operate directly in this construction although it may do so indirectly. These foods are formed in much smaller quantities than the carbohydrates but they are of the greatest importance in the nourishment of the plant. This is especially true as regards the living substance, protoplasm, which resembles somewhat in composition some of the more complex proteids.

9. The Distribution of the Foods.—Let us now consider what becomes of these foods. A small part is consumed on the spot by the manufacturing cells themselves, a larger portion is transported through the vascular bundles to all the living and growing cells of the plant body, but as the plant approaches the completion of its annual growth a larger and larger part of the food is transferred to special parts of the plant, such as buds, roots, seeds. Here it is stored as a reserve food to meet the needs of the plant at such times as it is not able to manufacture food. This transfer of foods is slow and consequently the rapidly constructed sugars gradually accumulate in the chlorenchyma during the day. This would result in the saturation of the cells with sugar and so stop the work of photosynthesis were it not for the fact that the chloroplasts quickly change the sugar to insoluble starch, thus leaving the cells free to receive more sugar. If chloroplasts of well-sunned, starch-forming leaves are examined

in the afternoon, they will be found to contain minute glistening bodies, the starch grains (Fig. 9, *s*). The chloroplasts have the power to absorb sugar and secrete starch. The construction of sugars and their transformation into starch is effected with surprising rapidity. If some plants of pond scum are placed in the dark for twenty-four hours so that all sugars and starches may be consumed, starch will re-appear in the cells in these plants in from three to five minutes after their return to the light. Consider the absorptions, decompositions, recompositions, and trans-



FIG. 9. Greatly enlarged chloroplasts from leaf of moss: *a*, plastid with three starch grains, *s; b*, plastid elongating preparatory to division; *c*, *d*, later stages in division.

formations that have been effected in this short time. Is it surprising that the exact nature of the changes effected in the manufacture of foods is not known? It is evident as a result of the rapid formation of sugar that starch must accumulate in the plastids during the day. At night sugar is no longer formed but the transfer of food continues as in the day. The insoluble starch is now dissolved by means of a ferment or enzyme into sugar so that by morning the cells are quite emptied of the sugar and ready for another day's work. This accumulation and transfer of foods can easily be demonstrated by cutting off in the afternoon a leaf from clover, bean or other starch-forming plant and placing them in alcohol. When blanched and tested with iodine the blue or blue-black color shows that they have accumulated a large amount of starch. Test in the same way leaves taken from these plants early in the morning. No starch reaction will be seen because the cells have been freed from their starch during the night. The formation of starch is often used

as a test for photosynthesis, but it must be borne in mind that photosynthesis results in the formation of various substances, and that some plants indeed do not form starch at all.

10. The Storage of Foods.—It has been stated that the foods not required for the nourishment and growth of the plant are stored in the region of buds, the growing portions of stems, in bulbs, in roots, in seeds, and in fruits. These are the so-called storage foods and they remain in these regions for considerable periods or until the conditions are favorable for a renewal of growth. They generally assume much more definite forms than the products constructed in the chlorenchyma which were designed for immediate use or translocation to other parts. If we examine a potato the cells will seem to be filled with very regularly constructed starch grains (Fig. 10, *A*). In the



FIG. 10. Storage food: *A*, starch grains in cell of potato—*n*, nucleus. *B*, cells of bean, the smaller particles being proteid grains.

pea seed and in wheat and other grains proteid granules are associated with the starch grains (Fig. 10, *B*). These starch grains are formed by rather colorless plastids called leucoplasts (Fig. 3, *E*) that closely resemble the chloroplasts. The starch grain first appears in the leucoplast as a minute point. Its stratified structure is due to the successive deposits of starch by the leucoplasts. If the grain is built up equally on all sides by the leucoplasts it will possess regular and concentric strata as in the bean (Fig. 11, *A*). If the bulk of the leucoplast lies on one side of the grain, this side of the grain will receive more material and consequently the starch grain will become more or less one-

sided as in the potato (Fig. 11, *B*). Frequently two or more grains originate in one leucoplast and compound grains result (Fig. 11, *C*). Proteids and other foods are likewise transported in solution to the storage organs where they may or may not be deposited in solid form. This is usually effected directly by the protoplasm of the cell without the aid of any plastid. Animals have learned to use the foods stored in these organs just as does the plant. It is interesting, however, to note that we cannot use them as economically as the plant does. In the

FIG. 11.



FIG. 12.

FIG. 11. Starch grains: *A*, from bean. *B*, from potato. *C*, compound grain from potato.

FIG. 12. Section of the outer portion of a grain of wheat: *p*, cells containing proteid grains. The larger cells are filled with starch.—After Strasburger.

wheat, for example, the proteid material is largely confined to the outer portion of the grain (Fig. 12). This layer is of a darker color and in the "best flour" it is excluded in the milling process, leaving the flour very white. So the most nutritious part of the wheat does not appear in white bread, but it should be remembered that white bread is better than the much exploited "whole wheat" products because the proteid material is digestible to only a slight degree. These reserve foods are made available to the plant in the same way as to the animal. They are first put into solutions by ferments or enzymes. Saliva in the animal and diastase in the plant are common examples of enzymes which change starch into sugar. So there are enzymes which transform each reserve food and render them capable of

transport and incorporation into the substance of the plant body. The distribution of the foods has been compared to the currency of a country; the storage foods are the bank reserve while the sugars and other solutions are the circulating currency.

11. Significance of the Leaf Structure.—We now begin to undersand the significance of the structure and form of the leaf. The broad blade is a device to catch the energy of the sunlight. The stomata afford access of CO_2 and escape of oxygen. The intercellular spaces in the spongy mesophyll increase the area for the absorption of CO_2 and excretion of oxygen to such an extent that the internal surface of the leaf exceeds the external many times. It will be noticed that the cells of the spongy mesophyll vary in character. This is because of the different functions that they perform. The collecting cells (Fig. 5, *col*) are closely applied to the palisade cells and collect the sugars and other foods from them while the elongated conducting cells transport this food to the cells of the vascular bundles.

12. The Second Function of the Leaf.—Respiration or breathing is a second function of leaves. While work of this kind is performed by all living cells it may be considered at this point because in the leaves the nature of the work is very well illustrated. The work of respiration is usually associated with the absorption of oxygen and the giving off of CO_2. Animals and plants are constantly inhaling oxygen and exhaling CO_2. This interchange of gases is the reverse of that occurring in photosynthesis. Furthermore respiration goes on all the time when conditions are suitable for life while photosynthesis only is possible in the light. Let us consider the significance of this interchange of gases. Why do animals and plants breathe? The construction of the foods to which attention has been called may well be termed the storing up of the energy of the sunlight. When the chlorophyll absorbs certain rays of light it does not destroy any of the energy in those rays. The energy is changed to another form and it still exists in the sugars and other compounds that are built up through its power. The energy is locked up, so to speak, in the food compounds. If these compounds are decomposed into their simple elements again the

energy will be set free unimpaired. Oxygen is the principal agent in the decomposition of foods. It has a great attraction for nearly all the elements that compose the foods. It pulls them away from each other and we say the foods are decomposed. The changes occurring in the foods as the result of the action of oxygen vary in different plants according to the nature of the plant foods. In the majority of cases the decomposition of the complex food compounds continues until such simple compounds as CO_2 and H_2O are formed which we have seen are utilized in photosynthesis in the manufacture of foods. These products escape from the stoma as gas and water vapor. Accordingly respiration is usually indicated by the absorption of oxygen and the giving off of CO_2. Some plants, however, have sufficient oxygen in their tissues to enable them to breathe for considerable periods without absorbing oxygen from the air and on the other hand in some plants CO_2 is not always exhaled because the decomposition of foods is not carried so far as to result in the formation of so simple a product. We eat in order to gain possession of the energy locked up in the foods. We breathe in order that the oxygen may decompose these compounds and set free this energy which gives us power to work and move and keep our bodies warm. The plant lives in the same way. It only differs from the animal in that it has the added power to build up the complex food compounds from crude material. By the decomposition of these foods it gains energy to grow and carry on its other vital functions. The work of respiration is carried on more economically by green plants than by animals since in the animal the CO_2 escapes in the breath as a waste product, while the plant uses this CO_2 during the day time for the construction of foods. Consequently the escape of CO_2 cannot be detected in the light unless an examination of very rapidly growing organs be made. If a jar be nearly filled with opening buds of dandelions or rapidly growing shoots and then closed air tight, sufficient CO_2 will be respired in a few hours to extinguish a light that is lowered into the jar. In such instances as these very rapid respiration is necessary to furnish the required energy for growth and the volumes of CO_2 expired

exceeds many times the volume of CO_2 utilized in photosynthesis. A handful of germinating peas or beans placed in a closed jar for a few hours better illustrates the giving off of CO_2 because here there is no green tissue to absorb any of the CO_2. Plants are often considered unhealthful in sleeping rooms at night because of their exhalation of CO_2. It is well to remember that the amount of CO_2 expired by a plant is small and that a gas jet would furnish more CO_2 to the air than a window full of plants.

We are now in a position to understand the importance of photosynthesis and respiration. Photosynthesis keeps the air pure for breathing, decomposes the simple inorganic compounds and recombines them into foods which represent a certain amount of stored-up energy. Respiration breaks down the foods into products which form the tissues of the plant and sets free the energy necessary for the accomplishment of its growth and movements. In the animal and plant the oxygen decomposes the foods rather slowly and the energy set free is principally manifested as heat and as the power that enables them to perform their various activities. If the decomposition is sufficiently rapid light as well as heat appears. This is the nature of the reaction when plant tissues, such as wood, are burned. The oxygen so rapidly decomposes the products which compose the wood that heat and light are produced. In our fires we are dealing with the heat and sunlight that have been slowly stored up by the plant.

13. The Third Function of the Leaf, Transpiration.—This function refers to the giving off of water by the plant. While other parts of the plant assist in transpiration, the leaf is the principal organ upon which this very considerable work devolves. Water is given off from the plant as a vapor and for this reason transpiration is a more familiar phenomenon than photosynthesis and respiration, where we are dealing with an interchange of invisible gases. We see the vapor from plants growing in a window precipitated on the cool window panes in the form of drops. On hot summer days the leaves of plants droop. This is because they have transpired so much water that their cells are no longer distended by the water, consequently the cells shrink

and the leaves contract or wilt. When plants are covered by a bell jar or placed in a tight glass jar the water given off soon saturates the air and collects in drops on the sides of the jar. The amount of water transpired by a plant is surprisingly large and it is probably safe to state that usually it amounts daily during the hot summer months to more than the plant's weight. An oak with seven hundred thousand leaves was estimated by Ward to transpire from June to October 244,695 pounds of water. A birch with 200,000 leaves transpired 700 to 900 gallons on hot summer days. From careful measurements of the amount of water given off by grass plants it has been calculated that six and one-half tons per acre may be transpired daily during the summer. The question naturally arises, why is the transpiration of such large volumes necessary? We have noticed that the veins carry water to the green cells of the leaf. If water were the only food that the chlorenchyma required, the amount of transpiration would doubtless be greatly lessened. This water, however, contains other foods that are very necessary for the plant and which exist in very dilute solutions, probably not more than one part in ten thousand. In order therefore to secure the necessary foods from the soil the plant is obliged to absorb large volumes of water. Only a small portion of the water is required and the surplus is forced out of the cells and finds its way in the form of vapor along the intercellular spaces in the spongy mesophyll through the stomata to the air. The transpiration of these large volumes of water must be of importance also in keeping down the temperature of the plant during the burning summer heat. Transpiration is often compared to the evaporation of water from a dish. While it is controlled to a limited extent in the same manner as evaporation, it should be borne in mind that the giving off of water is intimately associated with the vital activities of the cells and that the loss of water is to a degree under the control of the plant. The cuticle, which is practically impervious to water and gases, extends as a thin coat over all parts of the plant body, and materially assists in controlling the amount of transpiration. As a result of this water-proof coat the vapor can only escape from the leaf through the stomata. These minute

openings are one of the most interesting features of the leaf. When there is an adequate supply of water the elliptical guard cells are drawn apart and give free exit to vapors and gases. However in cases of drought when a continued loss of water would prove harmful to the plant, or when an interchange of gases is no longer required then the guard cells close and so offer such a barrier against further loss of water that only the severest conditions, such as prolonged heat and drought, can overcome. The stomata in the majority of cases range in size from .0002 sq. mm. to .0008 sq. mm. and in numbers from four to seven thousand to a sq. cm. so that these microscopic openings furnish adequate space for the interchange of gases and the transpiration of water.

14. Noteworthy Features of Leaves.—Now that some idea has been gained of the nature and extent of the work performed by the leaves we are prepared to comprehend the meaning of the large extent of the leaf surface and of the arrangement, structure and modification of leaves. The broad blades are devices for gathering or absorbing as much of the sunlight as possible and they increase the surface of the plant many hundreds of times. Contrast the extent of surface of a tree in full foliage with the bare branches of the winter. A mature maple develops annually from one to two thousand yards of leaf surface.

One of the most noteworthy features about the leaves is their arrangement or "hang" on the branches. In some plants they are arranged in two rows, in other instances three, four, five, eight or more rows. By fastening a thread to a leaf and winding it about the stem so as to touch the petiole of each succeeding leaf the arrangement of the leaves becomes more obvious. Some plants have their leaves opposite in two rows or ranks, others opposite in four ranks, each succeeding set of leaves being at right angles to the lower set. This latter arrangement is called decussate (Fig. 13). More commonly the leaves are spirally arranged and the thread passed once, twice, thrice, five, eight, etc. times around the stem before a leaf is reached that is exactly over the one from which we started (Fig. 14). The number of leaves passed before reaching one that stands over the first leaf indicates the number of rows or ranks of leaves on the stem. This varia-

tion in the arrangement of the leaves is simply a device to bring the leaves into the light and prevent the shading of one leaf by another. In opposite two ranked leaves each succeeding leaf is over the one below it but the cutting off of the light is pre-

FIG. 13.

FIG. 14.

FIG. 13. Branch of *Forsythia,* leaves decussate, in four rows.
FIG. 14. Branch of poplar with leaves spirally arranged in five rows.

vented by lengthening the internodes and thus separating the leaves. The decussate arrangement of the leaves is a more economical arrangement because more leaves can be developed on a given length of stem without danger of shading. The spiral distribution of leaves is a still better device since each succeeding leaf is placed a little above and to one side of the next lower leaf. By this means the maximum of leaf surface can be exposed to the light in a given length of stem. No one can look at the leaves of the maples, dogwoods, ailanthus, creeping and climbing vines, etc., and not be impressed with the fact that the leaves are arranged so as to catch the most favorable light without shading or interfering with one another. Furthermore the arrangement

is such as to occupy practically all the space about the stem that receives light. This becomes very noticeable if we look directly down upon a shoot or better if we stand under a tree and look up into the branches. It will be seen that very little direct sunlight finds its way through the branch, so nicely are the leaves adjusted to each other. Some plants like the hickories, catalpa, etc., occupy all the available space with a few large leaves at the tips of the branches. In other cases, as in the willows, some lilies, etc., the same result is accomplished by many small leaves that can be arranged along the stem for considerable distances without shading. The nicety of leaf arrangements is especially noticeable in many horizontal and creeping stems. In such cases the leaves can only be exposed on the sides of the stem and consequently the stem may become twisted or the petioles variously elongated and curved in order to bring the leaves into proper relation to the light (Fig. 15). Notice the change in leaf ar-

Fig. 15. Horizontal branch of *Forsythia* with leaves in two rows owing to the alternate twisting of each internode, assisted by the curving of the petioles. Compare Fig. 13.

rangement in horizontal and erect stems of maple. In an example like the maple if the leaves could not change their position they would all be standing edgewise to the light on horizontal branches and therefore receive little of it. They not only place their blades at right angles to the light but owing to the greater elongation of each succeeding petiole from the apex toward the bottom of the stem, all the leaves are arranged one beyond another so as to overlap very little. This same device is noticed in many plants that produce their leaves close to the ground in rosettes as mullein, wood betony, plantain, etc. (Fig. 16). Compare the erect and horizontal branches of a variety of plants noting

by what devices the leaves are brought into the light. Especially instructive are the leaf arrangements of plants growing in windows or creeping over trellises, etc., where complicated twisting

FIG. 16. Leaf rosette of evening primrose. Each leaf blade has a longer petiole than the one above it, so that the leaves are not shaded by one another.

and elongation of stems and petioles are necessary to adjust the leaves to the one-sided illumination. The angular leaves of some begonias furnish excellent illustrations of this. Perhaps so many plants bear angular leaves because they can be better adjusted

FIG. 17. Leaf of Aralia. Note the angular shape of the leaflets and the smaller ones filling the space between the larger ones.

and fitted together without loss of space than would be the case in rounded leaves, and smaller leaves can also be more advantageously introduced between the larger ones (Fig. 17). The

lobing and branching of leaves has become a characteristic of many plants because such variations permit the illumination of a larger leaf surface. This is particularly noticeable in our oaks where the outer leaves are often deeply lobed, thus permitting considerable light to pass through to the underlying leaves. Observe also that these lower leaves are larger and less lobed, thus catching as much as possible of this rather feeble light (Fig. 18).





FIG. 18. Leaves of red oak: *A*, sunned leaf. *B*, shaded leaf.
FIG. 19. Compound leaves: *A*, red ash. *B*, horse chestnut.

In many plants the lobing extends quite to the middle of the leaf and the lobes are often attached to the midvein or midrib by a petiole. In this latter case the leaf is said to be compound (Fig. 19). All such modifications permit the development of numerous leaves upon the branches without the danger of shading. If there is still any doubt as to the perfection of this light catching arrangement of the leaves, try to substitute the somewhat similar leaves of two different trees as the birch and elm or the water beech (*Carpinus*) and the beech, noting how the leaves of one fit when placed upon the branch of the other.

15. The Cause of Leaf Arrangements.—How is this perfection in the arrangement of leaves accomplished? The leaf or

any other organ does not remain in a fixed position during its growth and development but owing to the fact that first one side and then an adjoining side of the organ is growing faster than any other part it comes about that the organ is bent from side to side by the more rapidly growing cells and the apex of the organ is often caused to travel through a rather irregular circle. These growing organs are sensitive to light, gravity, moisture and other stimuli. As a result of these movements the organ is brought into relation with various intensities of light, heat, etc. In certain positions the stimuli are not favorable and they cause it to grow away from this position while in other positions the stimuli act in the most favorable way and the organ is stimulated to its best growth. So during the development of the organ it is brought into varied relations with the forces which affect it and as a result of this experience the leaf or other organ is directed by the stimuli and at maturity finally comes to rest in a fixed position that is the most advantageous. The leaf ordinarily so

FIG. 20.



FIG. 21.

FIG. 20. Shoot of wild lettuce, *Lactuca,* with leaves turned edgewise to the light.

FIG. 21. Position of young leaves of hickory: *A,* scales of bud curving back showing the tightly folded leaves. *B,* later stage, leaves unfolding but still erect.

reacts to these stimuli that its blade is exposed at right angles to the rays of light, but in the iris, many grasses and rushes the blades are nearly erect. Quite a large number of plants bear their leaves edgewise. The giant trees of Australia (*Eucalyptus*) and the so-called compass plants are familiar examples (Fig. 20). Possibly these blades are driven into this position because they are more sensitive and the direct sunlight upon the broad surface of the blade would be injurious. This certainly appears to be the case in young leaves when emerging from the bud. Note the character of the foldings and the erect positions of such leaves (Fig. 21). The leaves of the horse chestnut assume at least three different positions during their growth. The folding of such leaves and their erect positions expose less surface. They consequently lose less heat and moisture, and the delicate growing cells are better protected against the intense sunlight. You must notice as the cells mature and become better protected that their irritability changes and they respond in a different way to light and gravity, etc. They begin to unfold and turn away from their erect positions and at maturity assume a fixed position to light, etc., that is quite different from the original position. The leaves of many plants do not have a fixed position but show more or less motility during their entire life. Interesting examples are seen in the sensitive plants and in members of the bean and oxalis family where marked changes result in the position of the leaves from alterations in the intensity of light, temperature, or moisture. Many of these plants fold their leaves when the temperature falls at sundown. Many flowers close in the same way. These changes are popularly termed sleep movements. They are caused not only by the changes in the intensity of the light but also by changes in temperature and moisture. In the case of these leaves the cells in the swollen organ (the pulvinus) at the base of each leaf are kept full of water so long as the temperature and light exert a suitable stimulus upon the plant, but when the conditions are unfavorable certain of the cells on one side of the pulvinus lose water and contract while the cells on the opposite side of the pulvinus remain rigid; consequently, the leaves droop and fold in various

ways. This is due in part to the fact that the contracted cells of the pulvinus are no longer capable of assisting in the support of the leaf and also to the fact that the rigid cells continue to exert a pressure which bends the pulvinus towards the side where the contracted cells are situated (Fig. 22). This folding of the

FIG. 22. Leaf position of the sensitive plant, *Mimosa: A,* in light. *B,* in darkness. The same movements occur as a result of unfavorable temperatures and humidity.

leaves materially reduces the area exposed to the atmosphere. Consequently there will be less loss of heat by radiation when the temperature falls. This appears to be the reason in the case of many plants for the folding of the leaves at night. Transpiration will be reduced in the same way. Many of our sensitive clovers (*Meibomia, Lespedeza*), sensitive peas (*Cassia*), sensitive plant (*Mimosa*), are able to check the excessive loss of water on hot dry days by the folding of their leaves. As soon as the loss of water becomes detrimental some of the leaves begin to fold and the reduction of leaf surface continues until the loss of water by transpiration is met by root absorption. Very commonly leaves that do not have this power of adjustment are able to accomplish the same results by a rolling of the leaf. This is particularly noticeable among the grasses, which, on hot, dry days, roll up their leaves so tightly as to change the appearance of the plants. The interesting feature of all this is that

these reactions and adjustments are purposive but not intelligent. The movements are not called forth by consciousness but by stimuli to which the irritable living substance is attuned or sensitive.

16. The Significance of Certain External Leaf Structures.— Many features connected with the structure of the leaf furnish admirable illustrations of the fitness of the leaf for the performance of its work. The stimulus of light and moisture have a marked influence upon the external structure and form of leaves. The epidermis is strikingly modified by such forces. In shade plants the epidermis consists of a rather delicate layer of cells with very thin cuticle. This gives sufficient protection to such plants, but in leaves exposed to intense sunlight and a hot and dry air, the epidermal cells become greatly thickened and often of two or more rows, while the cuticle may often form the larger part of the outer cell wall or even extend in between the cells (Fig. 23, *c*). Cell walls that are filled with cutin in this manner are said to be cutinized. Such features are particularly noticeable in desert plants, in long-lived leaves of many evergreens, as laurels and rhododendrons, in the needles of cone-bearing trees, and in all plants that are exposed to the hot, dry air of summer or the drying winds of winter. Such leaves may be further protected and strengthened by thick walled elongated cells, stereome (Fig. 23, *st*). Coatings of wax, mucilage and



FIG. 23. Cross-section of the outer cells of a leaf of pine, showing the firm character of the outer cells of the tough leaf: *s*, stoma; *e*, epidermis; *c*, cuticle; *st*, stereome; *m*, mesophyll cells.

lime are also frequently developed upon the cuticle to further re-inforce the impermeability of the epidermis.

The stomata are also developed as to meet the conditions under which the plant grows. In dorsiventral leaves they are more numerous on the under side of the leaves because they are less liable to be filled with water by rains and with dust which would prevent the interchange of gases. The plugging of the stomata by dust is one of the causes of the sickliness of plants in homes. The arrangement of the stomata on the under side of the leaf is also of especial advantage because the direct light does not fall upon them and cause an excessive loss of water. The stomata of floating leaves, however, are upon the upper surface and their stoppage with water is prevented by waxy coatings, as can be easily demonstrated by dipping a leaf of a water lily or spatter dock in the water. When the leaf is removed the water runs off of the waxed surface without wetting it. Some leaves are more or less erect, as the cattails, rushes and grasses, and these have the stomata developed more or less evenly on both surfaces. Doubtless the intense light of midday is not beneficial to these leaves and the blade of the leaf is consequently placed parallel with the sun's rays. This arrangement of the leaf permits a direct illumination of the leaf only in the morning and afternoon when the intensity of the light is feebler. Leaves develop fewer stomata in a dry atmosphere and intense light than in moist and shady places. Light and moisture also influence the position of the stomata in another interesting way. Plants living in the



FIG. 24. Cross-section of a leaf of the inch plant, *Tradescantia,* showing the delicate character of the cells and the raised stoma of this shade-loving plant.

shade or in the presence of an abundant soil-moisture develop the stomata on a level with the leaf surface (Fig. 24), because there is no necessity of conserving the water supply. For the same reason some aquatics have lost altogether the power of closing their stomata. On the other hand plants that are exposed to arid conditions or drying winds develop the stomata well below the surface of the leaf, as in the cactus and in the needles of conifers (Fig. 23, *s*), or in furrows, as in certain grasses, or at the bottom of minute pores, as in the oleander. These depressions remove the stomata from the dry winds and prevent the direct contact of the moist air in the leaf with the dry atmosphere. The pores in the oleander contain hairs which would check transpiration just as a plug of cotton in the neck of a flask would lessen the evaporation of the fluid in the flask. The chief purpose of this arrangement, however, is to prevent the stopping of the stomata with water. This plant grows naturally along the banks of streams where it is subject almost nightly to heavy dews. When the stomata become filled with dew it requires several hours of sunshine to drive the moisture from these capillary openings. Consequently during this time there could be no interchange of gases and the vital functions of the leaves would be largely stopped. Many devices appear that prevent water entering the stomata. Attention has been called to the development of stomata on the under surface of leaves and to waxy coatings as protective features of this nature. Every one is familiar with the waxy appearance or bloom of many fruits, as the plum; or of leaves, as the cabbage; and of stems, as the raspberry. Water is unable to spread over the wax and enter the stomata because a thin layer of air clings to it. The silvery appearance of many grass leaves and of the jewel-weed when immersed in water is due to this thin film of air which reflects the light. Accordingly the water does not really touch the leaf and you notice that it is quite dry when it is removed from the water save for a few drops which readily run off. Protection against wetting is also obtained by the development of coatings of hairs. This is noticeable in many Alpine and polar plants where they are constantly exposed to heavy fogs, dews, and rains.

Plant hairs do not always serve this purpose. Desert plants and those living in dry localities are frequently characterized by hairy coatings. Such plants are frequently termed xerophytes in contradistinction to aquatics or hydrophytes and to plants living in moderately moist soils, which are called mesophytes. The hairy coatings of xerophytes give them their familiar silvery or hoary appearance as seen in the sage-brush, dusty miller, and mullein. The hairs are usually empty tubes or cells which reflect a large portion of the intense light and heat and consequently materially lessen the loss of water, just as straw sprinkled upon the ground keeps the earth below moist and cool. In many instances plant hairs develop mucilage which protects as with a varnish growing regions as buds or young shoots and leaves of birches, alders, poplars, and peach; or oils and poisonous substances are formed which repel by disagreeable odors or tastes the attacks of animals. The nettles are shunned because of the burning sting produced by the plant hairs. The upper portion of these hairs (Fig. 25) is practically a delicate glass tube since



FIG. 25. Stinging hair of nettle: *A*, portion of epidermis bearing hair. *B*, tip of hair enlarged, showing easily detachable knob.—I. D. Cardiff.

the walls are filled with silica. The basal portion of the hair is composed of soft, yielding cellulose so that the fluid collecting in the hair distends this portion of it. The knob at the end is fastened to the hair by so thin a ring of wall, as seen in Fig. 25, *B*, that the least touch breaks it off. In this way the hair is transformed into a veritable hypodermic needle which easily punc-

tures the skin while the distended basal portion contracts and forces out some of the fluid. The burning sensation or sting that immediately follows the puncture of the skin is caused by the injection of formic acid while a variety of other poisons produce the subsequent irritations. These poisons are so powerful in some of the East Indian nettles as to produce serious results, even tetanus or lock-jaw.

17. Significance of Certain Internal Leaf Structures.—The changes or modifications produced in the chlorenchyma by various stimuli are quite as striking as in the case of the epidermis. The palisade cells are an excellent illustration of this point. These cells are developed as a result of the stimulating influence of light (Fig. 26). Leaves growing in the deep shade show little indication of palisade structure. The compact arrangement of these cells prevents to a large degree the access of the hot, dry air and consequently loss of water is less than would be the case if the cells were separated by intercellular spaces. Shade



FIG. 26.



FIG. 27.

FIG. 26. Section of a leaf of *Rhododendron*. Note the compact palisade tissue which enables the plant to withstand drying winds.

FIG. 27. Section of a leaf of skunk cabbage, *Spathyema*. Note the poorly developed palisade tissue and the loose arrangement of the cells of this plant which lives in moist, shaded places.

and moisture loving plants are not obliged to conserve the amount of water received and these two forces, feeble light and moisture, produce a larger and looser arrangement of tissues which is favorable to an interchange of gases and transpiration (Fig. 27). Particularly is this noticeable in aquatics as in the water lilies and many rushes, etc., where the tissue is loose and spongy and permits a ready circulation of gases from the leaves to the roots and to all parts of the plant body even when submerged. This loose arrangement of tissues also renders water plants very buoyant and consequently they are less liable to injury from the currents of the water. Aquatic plants have little need of strengthening and conducting tissues because they are supported by the water and the crude foods being absorbed by nearly all parts of the plant body, do not require elaborate bundles for their conduction. The palisade arrangement of tissues is also favorable to photosynthesis. In intense light the movements of the protoplasm usually distribute the chloroplasts along the sides of the palisade cells so that they are edgewise to the light, but in feeble light they are distributed throughout the cell with their broad surfaces to the light. Just why this disposition of the plastids is made is not known but that it is of vital importance to the formation and distribution of substances in the cell is not to be questioned. We now understand why the shade leaves are thin and broad and soft while leaves of desert plants and those exposed to severe drying winds of summer or winter are thick, compact, firmer and often leathery and hairy.

One of the most interesting adaptive features of leaves is seen in the leaf fall of our deciduous trees and shrubs. In the tropics the leaves remain on the perennial plants often for long periods, but in temperate climates the severe winters necessitate the annual dropping of the leaves except in a comparatively few evergreens where the thick leathery leaves are able to endure such conditions. In our deciduous plants, when the conditions are no longer favorable for the performance of leaf work, the cells at the base of the petiole begin to divide and a delicate layer of cells, the separating layer, is formed across the petiole (Fig. 28). Various conditions induce this growth for there is

more or less leaf fall at all seasons of the year. Doubtless lack of nutriment has much to do with it. If a leafy branch of horse chestnut is placed between moist paper in a few days the separating layer will have formed and the leaf will have dropped from the branch. The delicate cells of the separating layer may become changed into cork cells or the outer cells of the layer may



FIG. 28. Leaf fall: *A*, branch of horse-chestnut showing scar formed by the fall of the leaf. The dots on the scar show the position of the vascular bundles that are finally broken by the weight of the leaf. At the left the base of the petiole is shown. *B*, diagram of a section through a twig of hickory—*s*, separating layer at base of petiole; *v*, vascular bundles. *C*, enlarged view of the separating layer—*c*, cork cells that heal the wound caused by the fall of the leaf. The granular cells are the outer region of the separating layer, and they are beginning to break down, as seen in the upper part of the figure, at *x*, thus causing the fall of the leaf; *st*, cells of the stem containing starch.

break down or they may become rounded off so that water collects in the intercellular spaces. Thus the leaf is attached to the stem chiefly by the veins since the delicate separating layer

offers little support. If now the water in the intercellular spaces of the separating layer should freeze, the expansion of the water as it freezes would so rupture the remaining tissue that we would have the familiar sight of the leaves falling in a shower in the morning after a frost—either of their own weight or with the slightest breeze. It will also be noticed that owing to the formation of the cork layer or the drying up of the delicate cells of the separating layer, the scar formed by the fall of the leaf is nicely healed and closed against any loss of fluids or the entrance of any organism (Fig. 28, *C*).

The casting off of the leaves reduces the area of the plant that is exposed to the unfavorable conditions to a minimum. This reduction of the surface is effected in one way or another by a great variety of plants. Much of our spring vegetation is possible by reason of it. The spring beauties, anemones, fawn lilies, jack-in-the-pulpits, etc., practically complete their growth before the larger summer forms appear. During this short period sufficient food is manufactured by the leaves to mature the seeds and fill the storage organs in the underground bulbs and stems. When the larger forms appear later in the season these plants are overshadowed, their leaves wither, and they are reduced to the small bulbs and stems hidden in the earth. In this condition they remain until the next spring when the abundance of stored food enables them to complete their growth before other competing forms appear.

This ability to reduce the surface of the plant body makes possible the existence of much of the plant life of arid and desert regions. During the short rainy seasons the stored foods enable the plant to quickly put forth the leaves which in turn manufacture the foods which are stored in seeds, stem and roots. Then practically all trace of these plants is destroyed by the drought. Only the seeds or a greatly reduced portion of the plant remain alive and this is often further protected by being hidden in the ground. A return of favorable conditions quickly awakens these plants to growth. Mr. Dan Beard relates an interesting experience in the desert regions of Texas that illustrates this feature of plant life. At the close of the dry season

this territory was a bare, sun-cracked plain swept by hot, dry winds. Not a green leaf was to be seen. Heavy rains transformed this level tract of land into an inland sea. In a few days the water disappeared and one morning he was surprised to see green blades appearing everywhere and in a few days the barren earth was covered with an almost tropical profusion of vegetation and bright flowers. Many desert plants to be sure develop permanent aerial parts but here, too, the most striking feature is the extreme reduction of the organs. Compare the areas of the Spanish bayonet, aloes, grease wood and sage with that of our leafy plants. The cactus represents one of the extreme forms of reduction. The leaves have been dispensed with entirely and are not represented save possibly by the spines. Consequently the work of the leaves devolves upon the stems. But even with the reduction of surface and the development of thickened, epidermal cells, cuticle, hairs, wax, etc., which cause the tough, leathery character of the organs, these plants would not be able to withstand the drying heat of the desert were it not for the fact that the cells are rich in mucilaginous substances and various salts which effectually retain the water. While desert plants receive very meager amounts of water, this is so well protected that the plants are usually rich in water. Drinking water is frequently obtained in desert regions from cacti by pounding up the pulpy interior and squeezing out the water. Animals are well aware of this rich storehouse and will eagerly devour the cacti after the spines are burned off. The development of these storage cells accounts for the fleshy character of many alkaline, saline, and desert plants. Curiously enough there are many examples of xerophytic plants living in bogs and marshes, as, for example, the rushes and sedges, the horse tail ferns (*Equisetum*), the lamb kill (*Kalmia*) and leather leaf (*Chamædaphne*), etc. The cause of the association of these plants with aquatic forms is not known. In exposed moors and heaths these reduced forms would have decided advantage because of their protection against drying winds. Our sedges and rushes are exposed to very intense heat and light which may possibly cause so heavy a transpiration that these plants are not

able to meet the loss by root absorption. That these plants may have some difficulty in absorbing sufficient water, owing to a lack of proper gas interchange in the roots, is indicated by the curving up into the air of the roots of the bald cypress (*Taxodium*) to form the so called knees. These outgrowths are composed of loose spongy tissues and probably supply the roots with air. At least they are not formed when the trees grow in sufficiently dry and aërated soil. For the same reason the roots of seedlings grown in too damp soil grow up to the surface of the soil. It has also been suggested that these aquatic xerophytes at one time lived under really arid conditions and owing to causes not known they have become restricted to bogs and marshes without loss of xerophytic characters.

CHAPTER II

THE ROOT

18. Character of Primitive Plants.—The first forms of life upon the earth were doubtless unattached unicellular plants that lived in the water or moist places. All the varied functions of the plant body, as the absorption of gases and material from the soil and the manufacture of organic compounds, etc., were performed alike by each cell. This was easily accomplished owing to the simplicity of the organism and also because the plant was surrounded by water which contained all the substances required in the construction of foods. Perhaps the first change produced in these plants by their surroundings was a modification of a part of a cell so that it served as an organ of attachment, or root, anchoring the plant to the substratum. As the plant continued to change and become more complex, and especially as it became more and more subject to drier conditions, special organs for the absorption of materials became a necessity. This change from an aquatic to a terrestrial life finally left the root alone in touch with the crude materials. Thus the root which at first was only an anchoring organ, came later to function also as the principal absorbing organ of the plant. Next to the leaf the root commands our attention because of its fitness for the accomplishment of this work. Practically all the water required by the plants, and all the elements, save carbon and oxygen, utilized in the construction of its foods are absorbed by the roots.

19. Root Hairs.—We will first be interested to examine the structure and nature of the absorbing apparatus of the root. If seeds of radish, mustard, or other plant are germinated upon moist sand or moist blotting paper it will be seen when the roots have attained a length of several cm. that a portion of each root is covered with delicate hairs (Fig. 29, *A*). There are several interesting features about these root hairs. In the first place they

are extremely delicate tubular outgrowths of the epidermal cells (Fig. 30). Note that they always begin to grow a few mm. back from the root tip and at the opposite end of the zone of root hairs the tubes are withering or dying off, leaving the older portion of the root covered only with the epidermis. If a root is marked into 2 mm. spaces with India ink it will be found after 12 to 15 hours that only the portion below the root hairs is elonga-

FIG. 29.



FIG. 30.

FIG. 29. The absorbing surface of roots: *A*, seedling of mustard showing extent of surface covered by root hairs. *B*, an older branching root, the shaded areas near the tip are due to particles of earth clinging to the root hairs. The remaining portions of the root are free hairs and take little or no part in absorption.

FIG. 30. Origin and character of root hairs: *A*, first appearance of a root hair due to the pushing out or growth of a definite portion of the surface of an epidermal cell. *B*, older hairs that have become irregular through coming in contact with soil particles.

ting. The first space from the tip has grown very little, the second and third much more, and in the other spaces there is a graded retarding of the growth until 8 to 12 mm. from the tip elongation has ceased. It is at the point where elongation has ceased that the root hairs begin to appear, but the total number existing on the root at any one time does not materially vary owing to the fact that they are dying off farther back on the root with the same rapidity that characterises their formation near the tip (Fig. 29, *B*). Before the significance of these peculiar features can be understood it will be necessary to con-

sider the nature of soils and the relation of the substances absorbed by the plant to the soils.

20. The Nature of Soils.—The soil consists of minute mineral particles derived from disintegrating rocks, and organic particles, or humus, formed from decaying plants and animals. These particles are loosely put together and are associated with a varying volume of water and air. From this source are derived the necessary mineral or crude foods, *i. e.,* potassium, calcium, magnesium, iron, nitrogen, phosphorus, sulphur, and also water. These substances are distributed in the soil in a rather remarkable manner. The water, for example, either fills a portion of the space between the earth particles and is then known as capillary water or it forms thin films about the earth particles, termed hygroscopic water. In addition there is to be considered the free water which runs off through the larger spaces in the soil. The capillary water is more or less mobile according to the fineness or coarseness of the soil particles and may move up or in any direction towards drier soils. In this way water is drawn to the dry surface of the soil from a depth of four feet in sand to six feet in clay. The hygroscopic water, however, remains untouched even on the hottest days and may only be driven off by heating the soil to 100° C. The mineral particles not only retain water, but some of them furnish necessary crude foods, while others are equally important because they enter into combination or fix important soluble mineral substances. The nitrogen in ammonia is fixed in this way by the compounds of aluminium in clay soils and soluble phosphoric and potassium compounds are fixed and rendered insoluble by lime and magnesia. Were it not for this fact these important substances would be washed out of the soils by the first rain.

21. Relation of the Root Hairs to the Soil Particles.—We are now prepared to understand the significance of the structures noted in the root and their relation to the physical properties of the soil. In the first place we see that these microscopic root hairs can readily penetrate any cavity and they are so delicate and soft that they become moulded about any soil particle in their line of growth (Fig. 30, *B*). So firmly are they attached

to these particles that it is impossible to detach them without injuring the root hairs (Fig. 29, *B*). This arrangement admirably adapts them to gathering up the water and other crude material in the soil. The tubes are filled with organic acids and other substances which have an affinity for water and such earth substances as the plant requires. In this way the material in the soils are drawn through the delicate walls of the root hairs, a process termed osmosis. The acids in the root hairs also reach through the cell walls and so come in contact with the insoluble substances, dissolving them and thus rendering them also capable of absorption by osmosis. This acid reaction of roots can readily be seen by placing a fresh root upon blue litmus paper which will turn red owing to the acid on the root hairs. The ability of the roots to dissolve mineral matter may be shown by allowing roots to grow over the clean surface of oyster shells or a piece of polished marble buried in the soil. Delicate grooves in the marble show where the roots have dissolved the marble in the course of their growth. It is important to note that not all the space between the soil particles is filled with water because living cells of the root as in the leaf and stem are constantly respiring, absorbing oxygen and giving off CO_2. The circulation of air in the soil is not only very important for supplying oxygen and bringing about a diffusion of CO_2, but these gases also materially assist the root in the decomposition of various minerals and insoluble compounds.

We are now prepared to understand why the root hairs are developed just back of the zone of elongation. Any disturbance of these delicate absorbing organs, such as the extension of that part of the root to which they are attached, would result in tearing and killing them.

22. Familiar Facts About Roots.—The character of the root and its relation to the physical properties of the soil explain many familiar phenomena of plant life. We see why plants must be repotted. The root hairs are constantly being formed back of the growing zone and dying off behind. During their short existence they absorb much of the foods in their immediate neighborhood, and unless opportunity of extending into new territory is

offered them, the root hairs soon perish and with them the plant. Continued absorption is only possible so long as the root continues to grow. For this reason a plant showing signs of starvation is removed from the jar, the tangled mass of roots next to the jar is cut off together with a portion of the earth and then repotted with fresh earth. In this way room is given for the rootlets to grow out into the earth and develop a new absorbing surface and additional materials are placed at the disposal of the root. We see now that transplanting of the plant should be done in September or October or in the early spring in order to permit the rootlets with their absorbing surfaces to become established in the soil before the leaves develop and bring about an excessive loss of water through transpiration. Why are plants often pruned or cut back in transplanting?

You have often noticed that fields of grain and other crops turn yellow and die during a long rainy season. This is due to the fact that the air spaces between the earth particles become filled with water and there is no longer possible an interchange of gases. At least two-fifths of the space in these capillary pores should be filled with air to ensure healthy plants. This is particularly noticeable in clay soils which are referred to as cold, wet, and sour soils. The reason of this is that the earth particles of such soils are very fine and consequently compact and the small capillary spaces between the mineral particles pull up and hold the water with great tenactity. Consequently acids from decaying vegetation and CO_2 from the roots as well as water accumulate in excess and render the soil unwholesome. Clay soils, while richer in plant foods than other soils require an admixture of sand and an adequate drainage to render them sufficiently porous for a proper circulation of air. Sandy soils are made of coarser particles and therefore looser. For this reason the capillary water is not retained for any considerable time and the soil is properly aerated. Injudicious sprinkling of lawns, gardens, or potted plants will result in the formation of a compact surface crust. This acts exactly like a clay soil and draws the water away from the region of the roots to the surface of the soil where it is lost by evaporation. Potted plants are best watered by stand-

ing the jar in water until the soil is wet. Further watering is not required until the jar sounds hollow when tapped—a matter of several days. So gardens and lawns should be thoroughly soaked during the night. The ordinary sprinkling only moistens the surface and leaves the soil in a worse condition for holding the moisture than before, because the hard crust draws up the water to the surface where the dry air readily evaporates it. The top soil of the garden should be kept loose by raking or hoeing, since the loose soil can not withdraw water from the root and it thus acts just as a mulch of straw or leaves or as the grass of the fields in retaining the water in the soil. In this connection the fact must not be lost sight of that a very considerable portion of water in the soil can not be withdrawn by evaporation. It is often surprising to note how plants continue to flourish during a drought when the earth is sun-baked. This is because the films of water about the soil particles cannot be withdrawn by the dryest and hottest air. The bulk of this water, however, is available to the roots, which, branching through the soil, send out root hairs in all directions and so absorb practically not only all the water in their neighborhood, but owing to the capillary action of the soil, water is drawn to them from the adjacent soil at a distance of from four to six feet.

23. Extent of the Root Surface.—It may seem surprising that the roots are able to take up the large volumes of water that are lost daily through transpiration. The spread of the root, however, is much more extensive than we think, usually quite equal to that of the branches, so that the absorbing portion of the root is beneath the drip of the branches or exposed to the rains and dews. Furthermore, when we come to add together the lengths of the numerous rootlets, the extent of the total root system is still more surprising. The total length of the roots of an oat plant has been estimated at 154 feet; of a corn plant, at 1320 feet; and of a squash plant at fifteen miles. These roots in ordinary soil reach a depth of from three to five feet, but in dry regions much greater depths are attained; for example, alfalfa 31 feet and mesquite 60 feet. The most important increase in the extent of the root system is due to the root hairs, 480 having been

estimated on .01 in. of a root of corn 1/17 in. in diameter, and 230 on a square mm. of a pea root. By this means the absorbing surface of roots is increased from fifteen to seventy-five times. It should be stated that these measurements have been obtained under unsatisfactory conditions and therefore only approximately represent the extent of the root surface.

24. The Structure of the Root.—You have noticed that only the apical portion of the root elongates and it may be well to first examine the structure of this region in order to see how growth is brought about and what changes are effected by it. The tip of the root is composed of very delicate cells. Certain of these cells are rapidly dividing by forming new walls through the middle of the cells, thus dividing each cell into two new cells. The division of a cell is effected in a very elaborate manner. The first indication of this growth appears in the nucleus. This organ of the cell enlarges (Fig. 31, *A*) and we now see that it has a very



FIG. 31. Cell division in root of corn: *A*, cell with nucleus enlarging preliminary to division—*l*, linin; *ch*, chromatin. The central dark body is the nucleus. *B*, later stage, the chromatin has increased and appears as a ribbon-like skein. *C*, formation of the chromosomes, *cr*. *D*, formation of the spindle and the arrangement of the chromosomes in the center of the spindle.—I. D. Cardiff.

complex structure, consisting of a protoplasmic mass surrounded by a delicate membrane and traversed by a network of delicate threads (linin) and containing one or more large granules (nucleoli). The most important portions of the nucleus are the minute particles (chromatin) which are associated with the linin threads. It has been supposed that this substance controls the

activity of the cell and determines the kind of cell that shall be formed and the work that it shall perform. As the nucleus enlarges the chromatin increases in amount and often forms a ribbon-like structure (Fig. 31, *B*) which finally contracts and divides, thus forming numerous small masses of chromatin, called chromosomes (Fig. 31, *C*). At this stage in the division delicate colorless strands begin to appear and ultimately grow into a spindle as shown in Fig. 31, *D*. There are two sets of these strands or fibrillæ; an inner and an outer series. The outer fibrillæ, in some way not undersood, arrange the chromosomes in the center or equator of the spindle (Fig. 31, *D*) where each chromosome divides by a longitudinal division into two equal parts. The fibrillae now separate the two halves of each chromosome and pull them to the opposite poles of the spindle so that each pole receives a half of every chromosome (Fig. 32, *E*).



FIG. 32. Later stages in the division of the cell: *E*, the fibrillae pulling the separated halves of the chromosomes to the opposite poles of the spindle. The inner series of fibrillae are now seen at the equator between the two groups of chromosomes. *F*, the chromosomes arranged at the poles and the inner fibrillae increasing in size and number. *G*, the fibrillae have increased in number until they nearly reach the opposite walls of the mother cell. Their thickening at the equator is the first indication of the wall separating the two new cells. *H*, position of the new wall clearly indicated.—I. D. Cardiff.

The chromosomes now become rearranged at the poles and two new nuclei are gradually formed like the original nucleus (Fig. 32, *F*, *G*). In the meantime the inner fibrillae of the spindle have shortened and become thicker at the equator (Fig. 32, *G*). This

thickening goes on assisted by the addition of new fibrillae at either side of the spindle until a delicate line (the cell wall) reaches across the old cell (Fig. 32, *H*), and the division of the mother cell into two daughter cells is completed. In this way new cells are being constantly added to the root. It will be noticed, if a longitudinal section through the middle of the root is examined, that various regions of the elongating root may be recognized owing to the difference in the character of their growth. In such a section (Fig. 33) we see that the tip of the root is covered with a mantle of cells, the root cap. This cap protects the delicate cells within like a thimble so that they are not exposed or injured as the root extends through the soil. At the tip of the root, just within the root cap, the cells are actively dividing and adding new cells to the end of the root and some cells are also added to the inner side of the root cap (Fig. 34). Owing to this unique arrangement it does not matter if the outer cells of the cap are injured or destroyed as the root pushes through the soil, because the cap is constantly being renewed from within, and so always furnishes adequate protection to the delicate cells within. The cells that are added to the tip of the root divide several times after their formation, so we find that the tip of the root for a distance of one or two mm. is composed of small cells that are in a process of division but that are enlarging to only a slight degree (Fig. 34). This region is called for this reason the formative region of the root. Back of the formative region for a distance of two to four mm. the cells are dividing to a less degree but are elongating very rapidly and changing in form (Fig. 35). This is the region of rapid elongation. Still further back elongation has ceased, but the walls of the cells are becoming thicker and the cells are changing in character, so that they perform different duties. This is illustrated in the cross section of the root (Fig. 36), where we see that the outer cells have become modified, forming an epidermis



FIG. 33. Diagram of a section taken through the center of a corn root: *r*, root cap; *e*, epidermis; *c*, cortex; *p*, central region.



FIG. 34. Portion of the formative region of the root taken at *a* in Fig. 33; *r*, root cap; *v*, cells which by repeated division add new cells to end of root and to inside of root cap. Note the numerous dividing cells and their uniformity in size.—I. D. Cardiff.



FIG. 35. Portion of the region of elongation of the root taken at *b* in Fig. 33. Note the larger size of the cells, increase in cell sap as indicated by the larger vacuoles and absence of cell division; *e*, epidermis; *c*, cortical region; *p*, central region.—I. D. Cardiff.

with root hairs for absorption. Within the epidermis is a broad zone of cells, the cortex, often used for the storage of manufactured foods, while in the center of the stem are the vascular bundles. The woody portion of the bundle, or xylem, radiates



FIG. 36. Cross-section of root taken above section shown in Fig. 33: *e*, epidermis with root hairs; *c*, cortex bounded on inner side by endodermis, *end*. Within is the central region containing vascular bundles; *x*, xylem; *p*, phloem.—I. D. Cardiff.

outward from the center and the soft portion, or phloem, alternates with it (Fig. 36, *x*, *p*). The materials absorbed from the soil are largely transported up to the stem and leaves through the xylem, and the foods manufactured by the leaves reach the root by means of the phloem cells. The branches of the root originate in a very curious way from the cells just outside the xylem. These cells by repeated divisions form lateral roots which gradually dissolve the tissues in their way and finally grow out to the surface of the root (Fig. 37). By this arrangement they are provided with a root cap and fully prepared to enter the soil on emerging from the old root.

25. The Transport of Water in the Root.—The inner layer of the cortex, the endodermis (Fig. 36, *end*), consists of a layer of cells which, in the older part of the roots, forms a very compact

and more or less cutinized ring of cells. This tissue is supposed to function in preventing an undue loss of water from the conducting strands of xylem to the cortex. It should be stated here that the root hairs not only absorb fluids but also force out the absorbed substances into the adjoining cells of the cortex. All living cells have this power of absorbing and expressing fluids.



FIG. 37. Cross-section of a root of lupine showing the origin of the lateral rootlets. Lettering as in Fig. 36.—H. O. Hanson.

So it comes about that the absorbed fluids are forced by the root hairs into the cortical cells and by them they are passed on to the cells of the xylem when by some force not known they are drawn up the stem into the leaves. The force exerted by these myriad number of cells in the root exerts a pressure that often amounts to considerably more than one atmosphere, *i. e.*, 15 lbs. to the square inch. It is the steady expressing of fluids by these cells that causes the familiar phenomena of the "bleeding," or flow of water from stumps in the spring; likewise the "bleeding" of injured branches, or the flow of sap, for in the stem the living cells are constantly absorbing and giving off water as in the root. In the summer, stumps and stems do not "bleed" as a rule because the water lost by transpiration nearly empties the cells,

whereas in the spring before the leaves appear they become filled with fluid.

26. The Sensitiveness of the Root.—The question may now be asked how does this elaborate root mechanism come into contact with the soil. If the root of a pea or bean seedling is placed horizontally in sawdust, after one or two hours it will begin to curve down toward the earth center. No matter in what position it is placed the result is always the same. We are so familiar with the downward growth of roots into the soil that we never stop to consider how it gains its sense of direction. Gravity is the stimulus that acts upon the irritable protoplasm of the cells and so brings about a growth reaction that sends the root in the right direction. When roots are placed in a horizontal position and slowly revolved on their longitudinal axes, no curvature results, since all parts of the cells are stimulated alike; but when allowed to rest a curvature results because possibly heavier particles in the cells fall to the lower sides of the cells and so produce an irritation through their unusual position. Only the cells in the first millimeter and a half of the root tip, *i. e.,* in the formative region, are sensitive to the stimulus of gravity. If this region of the root is carefully removed with a very sharp razor the root is no longer capable of responding to gravity although it may curve in various directions owing to the irritation produced by cutting. Furthermore, while the tip alone perceives the stimulus of gravity, the curvature occurs two or three mm. back of the tip *i. e.,* in the region of rapid elongation, so that we have the transmission of the impulse somewhat after the manner of our own nervous system. To be sure there are no specially constructed cells for conveying this impulse comparable to the nerves, but it has been shown that chemical changes are set up that extend from the tip to the region of curvature where a more extended growth of the cells on the upper side of the root is induced than on the lower side. This results in bending the root down into the soil. These reactions apply to the main or tap root. The lateral branches, or secondary roots, tend to grow more or less at right angles to the stimulus of gravity, while the short tertiary branches radiate out in all directions. Why the stimulus of

gravity causes a more extended growth on one side of the tap root than on the other or why the lateral roots are so differently affected can not yet be answered, but the nicety of these reactions in enabling the roots to reach in all directions and thoroughly explore the soil in their quest for food, is very apparent. The cells of the root tip are also sensitive to touch or contact. This enables the root to avoid obstacles and grow around rocks and stones. In this case the irritation of the root tip produces a more considerable growth in the zone of curvature on the side where the root is irritated with the result that the tip of the root is bent from the object. In this way roots work their way through the soil and avoid obstacles and grow around rocks, for as soon as the irritation of the obstruction is removed by the curvature, gravity will cause the root to grow down again. This sensitiveness to touch is very well shown by fastening with mucilage a bit of sand paper to the tip of the root when the irritation will cause such an excessive growth in the cells of the curving zone as to sometimes cause the tip to move through a complete loop.

The cells of the root tip also react to light and temperature and moisture and food. Roots avoid light and cold and they are thus brought into soils at the proper temperatures. In arctic regions where often only a few inches of the surface of the soil thaws, the roots will be directed straight down to within a short distance of the frozen earth when they will curve at a sharp angle. Every one is familiar with the fact that roots are caused to curve towards and follow the course of decaying tree trunks, being stimulated by the moisture and foods contained in them. Numerous instances might be cited where roots are attracted to wells, drains, etc., even surmounting considerable obstructions in order to reach the water. This localization of the sensitiveness to all external conditions that affect the root at the very tip is altogether admirable, because it enables it at the very start to come into the most helpful relations to its surroundings. It was a very clever fancy of Darwin to compare this localization to a brain center as found in certain low orders of animals.

27. Other Root Relations.—While 70 per cent. of our plants

develop roots adapted to absorbing substances from the soil there are several interesting modifications of the structures noted above. The roots of aquatics often do not develop root hairs because the roots are so finely divided and delicate as to enable them to absorb their crude materials directly from the water. Many plants develop roots in the air. These are sometimes simply clinging devices, as in the poison ivy; or they may be true absorbing organs, in some cases reaching down to the ground or to cup-shaped leaves that hold water, as in many tropical climbing plants. One of the most interesting of these modifications is found in the orchids that live on the damp tree trunks or on dripping rocks of tropical countries. Here the root is covered by a thick mantle of cells that are capable of holding the water that comes to them either in the form of dew or drainage or rain. Such plants, called epiphytes, are able to flourish in the air without the assistance of soil roots. Quite a large number of plants, parasites, depend entirely, or in part, upon other plants for their foods. The mistletoe is an example of this. The sticky seeds are carried by birds to the branches of oaks and other trees where they germinate, the roots penetrating the branches and withdrawing the necessary foods. The mistletoe withdraws largely crude materials from the branch because it has pale green leaves and is therefore capable of manufacturing some food on its own account. Our common dodder (*Cuscuta*) is entirely dependent upon other plants and illustrates the results that follow from the disuse of any function. It has lost its chloroplasts, the leaves are reduced to inconspicuous scales and the stem resembles a coarse yellow string while the root has disappeared entirely. The seeds produce a small thread-like shoot which soon perishes unless it comes in contact with a plant upon which it can live. These thread-like plants are sensitive to touch and so coil about the branches of any plant that they may chance to hit in their growth. If this plant happens to contain foods suitable to the dodder then a second stimulus is aroused which causes root-like branches to form. These organs penetrate the branches upon which the dodder is growing and absorb foods from it. Many plants live as parasites on the roots of a variety of herbaceous and

woody plants, as the beech drop, orobranches, broom rape, etc.

One of the most remarkable features of roots and one of the most important in the economy of the earth is found in that large family of bean or leguminous plants that are characterized by flowers and fruit like those of the pea. Minute plants called bacteria gain access to the roots of these plants through the root hairs and cause wart-like enlargements, tubercles, on the roots (Fig. 38). These bacteria have the power of combining the

FIG. 38. Tubercles formed on roots of lupine by nitrogen fixing bacteria.

nitrogen of the air with other elements, termed the fixation of nitrogen, and so forming a nitrogen-bearing compound that can be absorbed by the plant. Nitrogen is one of the very essential elements required in the construction of the living substance of the plant. Strangely enough, though the atmosphere contains over 70 per cent. of this gas, the plant has no way of utilizing it directly but only in compounds, such as ammonia and nitrates, etc. Compounds of this kind exist in very meager quantities upon the earth. The only sources of supply of any consequence are the decaying animal and vegetable life and manures; the guano deposits, now practically consumed; and the saltpeter beds of Peru and Chile, which will be exhausted by 1925. Consequently widespread alarm has arisen lest famines ultimately result through lack of these nitrogen compounds, and the situation becomes the

more serious since from one half to two thirds of the nitrogen compounds placed on the soils are lost annually in various ways, especially by the leaching out of these substances by rains and drainage waters, see page 44. Naturally of late years much attention has been directed to devices for uniting the nitrogen of the air with other elements in order to find a substitute for the rapidly disappearing nitrogenous compounds. Working upon the fact that nitrogen compounds are formed in the atmosphere by electrical discharges, several countries are now manufacturing by electrical processes large quantities of nitric acid. A much more serviceable source of nitrogen compounds for fertilizing purposes has been discovered in the production of calcium cyanamide.

Calcium carbide (CaC_2), which forms acetylene gas when placed in water, is produced in the manufacture of coke and when heated red hot unites with nitrogen forming calcium cyanamide ($CaCN_2$) and pure carbon. $CaCN_2$ slowly gives off ammonia and consequently when mixed with the soil it becomes an excellent fertilizer, being quite as valuable (weight for weight of the nitrogen contained) as the rapidly disappearing saltpetre. Calcium cyanamide is now being extensively manufactured on the Continent and it is not to be questioned that the construction of nitrogen compounds will be so extended and the process so improved as to meet the rapidly increasing demands of the cultivators of the soil.

Much study has also been given to the nitrogen-fixing bacteria as a further source of supply for nitrogen compounds. Failures attended the numerous attempts to cultivate these important plants with practical results until our own government through the work of Moore succeeded in making cultures of the bacteria that are now sold to farmers for mixing with their clover or other leguminous seeds. See Bulletin 71, 1905, U. S. Dept. of Agriculture. By mixing these bacteria with leguminous seeds the farmer is now able to raise a crop that contains a large amount of nitrogen derived from the air and by plowing under this crop a valuable nitrogenous fertilizer is added to the soil and the way is thus prepared for another crop, as wheat, which is dependent for nitrogen upon nitrogenous compounds. This alternation of crops

is of prime importance is successful farming. Each kind of plant takes from the soil certain elements and for this reason the alternating or succeeding crop should be of such a kind as to require other elements. As these elements are removed from the soil suitable fertilizers containing nitrogen, phosphorus, lime, potash, etc., must be added to the soil to replace the absorbed foods.

The majority of our trees and many of our shrubs are dependent upon certain fungi at least for such foods as are derived from humus. These fungi consist of delicate, cobwebby threads such as are seen on mouldy bread. These threads spread through the soil and either enter the outer cells of the root or simply form a mantle about the root. The fungi live upon decaying animal and plant life and transfer a portion of this food to the root and doubtless secures in return certain substances from the root. This mutually helpful relationship of two plants is termed commensalism. The majority of our heaths, evergreens, poplars, willows, beeches, and oaks have become dependent upon these fungi and do not thrive in soils where the fungi are not found. Perhaps this explains why it is so difficult to transplant certain shrubs and trees. The fungi are easily injured and do not become established in the new soil soon enough to keep the plant alive. Some plants have become so perfectly adapted to these fungi as to receive all necessary foods from them. Accordingly, their green leaves and roots have largely disappeared since they are no longer of service. Examples of this are seen in the white Indian pipe and pine-sap and in the coral root orchid.

28. Roots as Store-houses for Foods.—In all the cases heretofore considered the root has functioned in one way or another as an absorbing organ. It may also serve in other capacities, one of the more important of which is as a storage organ. Such roots become fleshy and filled with foods and are of great economic importance, furnishing a variety of nutritious vegetables, as the sweet potato, beet, turnip, etc. Many of these valuable plants are biennials. During the first season the plant develops only leaves and stores up food in its fleshy roots which is utilized in the following season in the production of flowers and seeds, after which the plant perishes. It is noteworthy that the biennial habit

has been induced in many of these plants by cultivation. If they are left to themselves in a wild condition they will soon revert to an annual growth, *i. e.*, producing seed and perishing during the season. Perennial plants continue to produce new root and stem growths and live on from year to year.

29. Anchoring and Supporting Roots.—Roots also play a very important role in anchoring and supporting the plant. Excellent examples of the supporting roots are seen in the Indian corn where numerous roots spring from the stem a short distance from the ground and reach out on all sides like guy ropes steadying the plant in the ground. Similar devices appear in our elms, maples, and beeches where enlargements of the roots at the base of the trunk rise up like girders bracing the tree against winds. More marked illustrations appear in many tropical plants, as the stilt roots of the mangrove and screw pines, and the buttressing roots of one of the Indian rubber plants (*Ficus*), etc. In the banyan tree of India the branches have an almost unlimited lateral growth owing to the fact that they are supported by a succession of roots that reach from the branches to the ground. Kerner cites an example of one of these trees with 300 large and 3,000 small prop roots. This tree sheltered a village of 100 native huts and an army of 5,000 men.

30. Binding Action of Roots.—The landscape would be constantly subject to great change either by erosion or through the action of winds were it not for the solidifying and binding action of roots upon the soil. The difficulty of breaking up the prairies and the turf of abandoned fields or meadows indicates the extent and completeness of the ramifications and interweaving of the roots. Miles of sandy reaches are held from shifting through the restraining action of roots and stems of grasses and other plants. The little town at the end of Cape Cod would have been submerged long ago by the shifting sand dunes had not suitable plants been planted to hold in check the loose sands. Roots not only bind the soil together but the older portion of the root usually possesses the power of contraction. This property results in the pulling down and fixing of the stem in the ground. You must have often wondered how stems and bulbs become so

deeply buried in the soil although the seeds are scattered on the surface of the soil. This is also well illustrated in the tips of raspberry bushes which come in contact with the soil through the bending of the stalk. Roots strike out from the tip and when thoroughly established in the earth the older portions contract and bury the tip in the soil. In the same way seedlings of various plants are slowly pulled down into the earth so that they finally become planted at considerable depths. . . . . .

CHAPTER III

THE STEM

31. The Function of the Stem.—The stem has for its chief function the production and display of the leaves and roots and the conduction of the materials which these organs are especially concerned in handling. It serves as a connection between them, carrying up the material absorbed by the roots and distributing the various substances received from the leaf. Like the traffic of a city this material is received at many stations and transported along various channels to many points. At one place some of it is used in the manufacture of food, at another point material is required for the nourishment and construction of the cells. Here a portion is stored, or again, there is the useless or waste material to be carried away. These movements are not to be considered as currents of water flowing now in this, now in that direction. Just as root hairs take up fluids from the soil and force them out into the adjacent cells, so each cell in the path of transport is constantly receiving material and forcing it out again into the surrounding tissue.

32. Structure of the Stem.—In order to comprehend the nature of the work performed by the stem it will be necessary to examine its structure and note the character of the apparatus utilized. If thin sections are made across the stems of seedlings, as of the castor bean, or of shoots just emerging from the buds, the tissues will resemble the arrangement shown in Fig. 39. All very young stems of dicotyledons, *i. e.*, plants whose seedlings have two leaves, as the squash, the bean, etc., show more or less clearly three regions; an epidermis, a cortex, and a central region surrounded by the cortex, an arrangement already noticed in the roots. The epidermis does not materially differ in structure, modification, or function from that already noted in the leaf, see page 8. The cortex is largely composed of paren-

chyma and extends from the epidermis to the endodermis which it includes. The endodermis is not so well marked as in the root



FIG. 39. Cross-section of young stem of castor bean: *e*, epidermis; *c*, cortex; *p*, pith or inner portion of central cylinder; *v*, vascular bundle, arranged in outer part of central cylinder.—H. O. Hanson.

and owing to the rapid growth of the tissues in this region it is often impossible to detect it. The cortex assists the leaves in photosynthesis, the outer portion as a rule being well supplied with chloroplasts, and it also serves as a storehouse for foods. Particularly is this true of the endodermis and adjoining cells which are often temporary receiving stations for the carbohydrates during their transport through the stem. The cells in the outer portion of the cortex frequently become thickened and more or less elongated to give strength to the stem. One of the most common modifications of this kind is shown in Fig. 40, *A*. This tissue, collenchyma, is characterized by the thickening of the cells at the angles or on all sides and by the silvery luster of the walls. The walls, though very elastic and tough, are capable of growth and so especially adapted to the support of the young elongating stems. In older stems, where elongation has ceased, greatly elongated cells, called stereome fibers or scleren-

chyma fibers, are formed. These cells have thick walls and tapering ends which interlock and bind the cells very firmly together (Fig. 40, *B*). These strengthening cells form a compact zone about the stem or they may be arranged in separate



FIG. 40. The stereome or strengthening cells of the cortex: *A*, collenchyma cells in growing stem of *Begonia*, showing above the cells in cross-section and below in longitudinal section. *B*, a group of elongated thick-walled cells, called sclerenchyma fibers or stereome fibers, from the stereome of mature flax stem.

bundles, in this latter case they often appear to the eye as light colored bands extending along the surface of the stems of many herbaceous plants. The central region of the stem is characterized by a mass of parenchyma with several vascular bundles arranged in a circle (Fig. 39). The central mass of parenchyma is called the pith.

33. The Vascular Bundle.—The vascular bundle contains three distinct regions; an inner thick-walled portion (the wood or xylem), an outer thin-walled portion (phloem), and between these two regions a delicate layer of cells, the cambium (Fig. 41). The transport of all substances is largely confined to the vascular bundles, the xylem conducting principally the crude materials while the bulk of the organic substances passes through the phloem. We are now interested to study the character of these cells and note their adaptation to the work in hand. In the xylem occur various large spaces, the vessels or ducts (Fig. 41, *v*), and smaller spaces, wood cells of different kinds. The cells of the phloem are much smaller, thinner walled and less numerous than those of the xylem. Therefore very thin sec-

tions are necessary in order to make clear all the different tissues. Sections from a squash stem may well be studied for this purpose because the various cells of the phloem are comparatively large and easily distinguishable. It should be stated that the bundle of the squash is peculiar in that a phloem region is developed on both sides of the xylem. The phloem, like the xylem, is also characterized by large cells, the sieve tubes, so named because these cells are elongated, tube like structures with the cross walls



FIG. 41. One of the vascular bundles shown in Fig. 39 enlarged: *x*, xylem; *v*, vessels or ducts; *p*, phloem; *s*, sieve tube; *ac*, accompanying cell; *c*, cambium; *st*, stereome.—H. O. Hanson.

perforated like a sieve (Fig. 41, *s*). A small cell, the accompanying cell, is associated with the sieve tube. Usually a varying amount of parenchyma also occurs in the phloem and often thick walled stereome fibers (Fig. 41, *st*). Between the xylem and phloem is a region of very delicate and regularly constructed cells, the cambium (Fig. 41, *c*). The growth of the bundle in

diameter and, in fact, of the entire stem is brought about very largely by the formation of new cells through the division of the cells in the cambium. The cross section of the bundle reveals the arrangement and distribution of the various tissues but it will be necessary to examine a section taken parallel with the length of the stem, *i. e.*, a longitudinal section, in order to arrive at an understanding of the structure and character of the cells themselves. Fig. 42 shows such a section of a bundle. The



FIG. 42. Longitudinal section of the bundle shown in Fig. 41: *x*, xylem; *ph*, phloem; *p*, pith; *v–v'''*, annular, spiral, scalariform and pitted vessels; *c*, cambium; *s*, sieve tubes; *ac*, accompanying cells; *st*, stereome.—H. O. Hanson.

vessels are now seen to be tubular structures with peculiar thickenings of their inner walls that assume the form of rings (annular vessels), spirals (spiral vessels), or the spirals may branch more or less (reticulate vessels), and often to such an extent that they cover the entire surface of the wall with the exception of numerous small spaces, pores, thus forming the pitted vessels (Fig. 42, *v*). These peculiar sculpturings prevent the crushing in and closing of these tubes and the annular and spiral thickenings also permit considerable elongation of such ducts which would not be possible if the thickening were more uniform over the wall. It will be noticed that the later formed

ducts are characterized by reticulate or pitted walls since the thickening largely occurs after the elongation has taken place. The ducts are composed of elongated cells but owing to the absorption of the majority of the cross walls the ducts finally come to resemble hollow tubes that often run for considerable distances through the stem without any cross walls at all. The smaller cells of the xylem assume various shapes. Some of them, the tracheids, have pointed ends and are characterized by markings similar to the ducts. Strengthening fibers, similar to those noted in the cortex, are of common occurrence and also short cells with blunt ends, wood parenchyma. The ducts and tracheids soon lose their cell contents but continue to function in the conduction of water. The wood parenchyma and other living cells retain their vitality for several years and generally function as storage cells. It is not improbable that they may furnish some of the energy required to force the water through the ducts and tracheids. It is significant that living tissue is always associated with these water conducting cells of the stem. Passing now to the phloem we see that the sieve tubes are made up of elongated cells with perforated or sieve like cross walls (Fig. 42, *s*). These minute openings adapt these cells to the transport of albuminous substances which do not readily diffuse through cell walls. The accompanying cells (Fig. 42, *ac*) which are cut off from the sieve tubes probably assist them in this work. At any rate they have abundant cell contents and contain nuclei, whereas, singularly enough, the nuclei of the sieve tubes soon disappear although the rest of the protoplasm of the cells remains active for one or more seasons. Elongated parenchyma cells are associated with the sieve tubes and accompanying cells, serving chiefly for the transport and the temporary storage of the more readily diffusible carbohydrates. Fibrous cells, stereome, are also of common occurrence in the phloem (Fig. 42, *st*). The cambium is composed of very delicate cells, owing to the fact that in this region cell division is taking place and the cells consequently have not yet attained their characteristic form (Fig. 42, *c*). We may best gain an idea of the growth of the cambium if we consider it as a single layer of cells between the xylem and

phloem. Each of these cells divides into two daughter cells (Fig. 43), usually the inner cell (*i. e.*, the cell next to the xylem)

FIG. 43. Diagram showing the mode of division of the cambium cells. The cambium cell is shaded to distinguish it from the cells derived from it. Note in the last division at the right that the inner daughter cell becomes the cambium cell while the outer develops into a phloem cell.

develops into a duct, tracheid, or other element of the xylem, while the outer cell, after increasing to the original size of the cambium cells, divides into two cells as at first. This process is repeated again and again, usually the inner cell growing into one of the cells of the xylem while the outer cell retains the power of further division. Less commonly the reverse method of growth takes place and we have the outer cell enlarging and forming one of the cells of the phloem while the inner cell acts as a cambium cell. In this way new cells are added to the xylem and phloem and the enlargement of the vascular bundle is effected.

34. The Conducting System of the Plant.—These vascular bundles extend from the root up through the stem and branches to the leaf where they divide again and again, reaching all parts of it. In this way the bundles become much reduced in size, the free end of the vein often consisting of a single tracheid to conduct the crude materials to the surrounding cells and elongated parenchyma cells to collect the manufactured foods (Fig. 5, *l*). We now see how admirably the vascular bundles are adapted to the transport of fluids. The xylem carries the substances received from the root hairs to all parts of the plant body quickly and with a minimum expenditure of energy because of the elongated character of its cells and the absorption of many of the cross walls of its cells. So the manufactured foods located at any point in stem or leaf are gathered up and distributed by the phloem, the more soluble and diffusible substances

being handled by the thin walled parenchyma, while non-diffusible, albuminous substances can readily be transported from cell to cell through the perforations of the sieve tubes.

35. The Strengthening Tissues of the Stem.—Attention may be directed at this point to the perfection of the arrangements that give rigidity to the stem. The delicate cells of very young stems or of any other part of the plant are distended by fluids which they absorb. In this way a very considerable force, known as the turgor of the cell, is exerted that may amount to over 200 pounds to the square inch. This pressure at first gives sufficient rigidity to the rapidly elongating stem, but as it increases in size and the strain upon it becomes more considerable, thick walled collenchyma cells are formed that keep pace in their growth with the elongation of the stem. When elongation finally ceases, tough fibers of stereome appear which are able to meet the increasing strain upon the stem due to the formation and development of its various organs. These strengthening tissues, with which must also be included the tissues of the xylem, are arranged with the same mechanical effects as are employed in the construction of buildings, bridges, etc. In bending a beam the strain falls especially on the convex and concave surfaces. The cells will be compressed on the concave side and stretched on the convex side, while the tissues in the center will be subject to the least strain. For this reason girders are strengthened on the surfaces receiving the strain and weakened in the center (Fig. 44). So also given a certain amount of



Fig. 44. Common form of girder.

material this is more effectively distributed in the form of a hollow column than in a solid column since less strain falls upon the center of the column. Therefore in stems we find the strengthening tissue very economically distributed at the periph-

ery either as bands or strands of collenchyma and stereome fibers and these structures are frequently reinforced by additional mechanical tissue in the phloem or just outside of it (Fig. 45). The vascular bundles themselves frequently furnish excellent examples of the girder, masses of stereome being found on the outside of the phloem and on the inside of the xylem (Fig. 45, *D*). The resistance of these tissues to cutting and breaking



FIG. 45. Cross-section of stems showing arrangement of strengthening tissues: *A*, burdock with strands of collenchyma—*st*, at periphery; *v*, vascular bundles. *B*, sweet clover, showing a similar arrangement. *C*, moonseed showing cylinder of stereome fibers outside of vascular bundles. *D*, rush with strands of stereome fibers at periphery of stem and also on inner and outer sides of the bundles; *v*, vascular bundles; *st*, stereome.

affords abundant evidence of their hardness and rigidity. It may be surprising to note that the sustaining strength of the stereome fibers is equal to that of the best wrought iron or

hammered steel while their ductility is ten to fifteen times that of iron. The superior quality of these tissues and the perfection of their arrangement result in structures that can not be approximated in any of our buildings. The height of the tallest chimneys scarcely exceeds the diameter of their bases more than fifteen times but many stems of rushes and grasses exceed the diameter of their bases from 200 to 500 times. It would be impossible with any metal to construct a column of the same length and weight and having the same rigidity, elasticity, and resisting power as these stems. We gain an idea of the toughness and durability of these fibers when we consider that linen, rope, matting, etc., are manufactured from them. In stems that increase greatly in diameter, as our trees, the mechanical tissues are confined to the wood or xylem. These are to be sure nearly solid colums, but even here there is sufficient strengthening tissue in the xylem at the periphery of the stem to support the trunk, as is often attested by the sturdy character of trees that have become hollow.

36. The Secondary Growth of the Stem.—The arrangement of the tissues as outlined above, termed the primary growth, remains practically unchanged in some annual plants but in such forms as live on from year to year or that increase materially in diameter there results extensive alterations in the structure of the stems owing to the formation of new tissues, especially in the region of the cortex and in the vascular bundles. The epidermis furnishes sufficient protection to such stems as do not increase materially in size, such as the majority of our annual plants; but in long lived stems, as shrubs and trees, where growth goes on from year to year the epidermis is not able to keep pace with the growth.

37. Cork Tissue.—To meet this condition a new tissue, the cork, is developed from certain cells called the cork cambium, usually situated near the epidermis (Fig. 46). The cells of the cork cambium divide much after the manner noted in the cambium of the vascular bundle, but there is this difference, the outer cell of the two daughter cells becomes a cork cell while the inner cell remains capable of further division. Only rarely does the

outer daughter cell act as the cambium cell while the inner cell develops into one of the cells of the cortex. As soon as the cork cells have reached their growth, a substance called suber



FIG. 46. Cross-section of the outer part of the stem of geranium. The cork cambium, *c*, originating in the cells immediately below the epidermis.

begins to be deposited in their walls. This completely changes the properties of the cellulose walls and renders them impervious to fluids and gases. So the cork cells have the same physical properties as the cuticle of the epidermis and owing to the continued activity of the cork cambium they are able to keep pace with the growth of the stem. It is evident that the cork cells must die as soon as they become impervious to fluids, and it must also follow that all cells lying outside of these cork cells will die since no fluids can reach them from the vascular bundles. These dead cork cells give the characteristic aspect to the outer bark of trees and we would naturally come to think of the coarse, dark bark as composed of rather thick cells. As a rule, however, the cork cells are quite delicate and compactly put together (Fig. 47) and the dark color is more usually due to the dried remains



FIG. 47.



FIG. 48.

FIG. 47. Cross-section of cells of bottle cork showing the delicate character of cork cells.

FIG. 48. Scale bark of pitch pine. The crescent-like lines in the bark show the successive positions of the cork cambium.

of the cell contents. The furrows and seams that occur in the bark of most trees are caused by the continued activity of the cork cambium and the cambium of the vascular bundles which adds each season new cells to the stem and so pushes out the cork cells. In this way it comes about that the cork cells are pushed further and further from the center of the stem and since they are not capable of dividing they are pulled apart, forming the characteristic furrows in the bark. It will often be noticed that the bark of some trees, see the cone-bearing trees, sycamore, etc., cleaves off in shell-like scales. This is due to the formation of a new cambium that joins on to the old cambium in the form of crescents (Fig. 48). Consequently as this bark is pushed out it breaks along these successive crescent shaped cambiums and finally cleaves off in scales. In some trees the cork cells break off about as fast as they are formed, so that a comparatively thin layer of cork cells remains attached to the trunk. Such trees have a smooth bark, as in the beech. In other cases the cork cells are formed in great abundance, and owing to the adhesion of the cells, thick layers of cork are formed, as in the oaks, and giant trees of the Pacific.

It is evident that this mantle of impervious cork cells must prevent the access of the atmosphere to the stem. We have seen



FIG. 49. Cross-section of the outer part of a stem showing the early development of a lenticel. Note the irregular character and loose arrangement of the cells below the stoma and the cork cambium, *c*, extending out on either side of the lenticel.

that all living cells respire. In some plants the air spaces extending from the leaves to all regions of the stem are sufficient to bring about an adequate interchange of gases but in the

majority of plants special devices are required to bring the living cells of the stem into more direct communication with the



FIG. 51.

FIG. 52.

FIG. 50. Later development of a lenticel: *c*, cork cambium; *i*, intercellular space.

FIG. 51. Surface view of lenticels: *A*, lenticels on branch of horse-chestnut appearing as minute brownish swellings. *B*, old lenticels on white birch appearing as dark lens-shaped streaks.

air. This work is effectively accomplished in herbaceous stems by the stomata, but in stems characterized by the formation of cork it is noticed that the cells just below the stomata begin to



FIG. 52. Cross-section of a stem of castor bean showing the formation of the cambium between two vascular bundles: *x*, xylem; *ph*, phloem; *c*, cambium of the bundle. The faint lines, *ic*, are the first divisions of the parenchyma cells between the bundles that result in the formation of the cambium zone.

divide and form a rather loose mass of cells that lift up and finally rupture the tissues about the stomata, thus forming a small, lens-shaped outgrowth on the surface of the young stem,

called a lenticel (Figs. 49; 51, *A*). Soon this growth becomes localized in a layer of cells situated on the inside of the lenticel (Fig. 50). This layer of cells is a part of the cork cambium, but strangely enough only loose cells are added to the lenticel during each year's growth and consequently a passage way is kept open to the living cells within the stem. The lenticels appear as minute points upon the surface of young stems but upon old trunks they often become greatly elongated, forming the characteristic bands on the bark of the birch and cherry, etc. (Fig. 51). In bottle cork, derived from the cork oak of the Mediterranean, the lenticels appear as minute lines often erroneously referred to as worm holes.

38. The Cambium Zone.—Let us now consider the changes that are effected in the vascular bundles through the activity of the cambium. The increase in the diameter of the stems of many annual plants and especially of our trees and shrubs is largely brought about by the formation of new cells derived



FIG. 53. Cross-section of a stem of castor bean in which the formation of the cambium zone, *cam*, as a ring of regular cells, has been completed; *p*, pith; *v*, vascular bundles; *st*, stereome; *c*, cortex. Compare Fig. 39. —H. O. Hanson.

from the cambium. While the vascular bundles are very small it will be noticed that the parenchyma cells separating the vascular bundles begin to divide so as to form a line of cells connecting the cambium of each bundle (Fig. 52). In some cases these divisions are at first somewhat irregular but soon the growth results in the formation of cells with parallel walls and in this way a ring of regular cells, the cambium zone, is formed which continues to divide as already noted in the cambium of the bundles (Fig. 53). Consequently new cells are now added not alone to the vascular bundles but also along the entire extent



FIG. 54. Cross-section of stem of castor bean three weeks older than one shown in Fig. 53. Note the changes that have occurred in the stem and especially the numerous cells that have been added to the xylem. *ck,* cork; *ph,* phloem; *cam,* cambium zone.—H. O. Hanson.

of the cambium zone. This growth results in the formation of a layer of xylem on the inside of the cambium zone and a layer of phloem on the outside, and so brings about the principal enlargement of the stem (Fig. 54). This growth is repeated

each spring in such plants as continue to enlarge from year to year, as in the shrubs and trees. The great majority of all the cells that are to be found in a year's growth are cut off from the cambium zone early in the spring, the first cells, and in fact the majority of all cells formed, being xylem cells and later a few cells are added to the phloem. Consequently the xylem increases faster than the phloem and forms the bulk of the tissues of the stem. The amount of the xylem added to the stem each year is generally indicated by the bands or annual rings (Fig. 55). This is due to the fact that the first cells formed in the spring have thinner walls and often contain a great many ducts, whereas the later formed cells are for the most part small and provided with thicker walls. So there is a sharp contrast between the small thick walled cells of the summer wood and the thinner walls and larger cells of the spring wood (Fig. 56). This appears to the eye as a band, the annual ring (Fig. 55). The



Fig. 55.



Fig. 56.

Fig. 55. Diagram of a cross-section of a stem of black oak four years old: *p*, pith; 1, 2, 3, 4, annual rings of xylem; *c*, cambium zone; *ph*, phloem; *cr*, cortex; *ck*, cork; *m*, medullary rays.

Fig. 56. Magnified view of a portion of one of the bands of black oak in Fig. 55, showing the thick-walled summer wood succeeded by the thinner-walled cells and vessels. This association of cells causes the banded appearance of the annual rings of growth. *m*, medullary ray; *v*, vessels in the spring wood.

difference in the structure of the cells of the spring and summer wood is doubtless due to the large volumes of water that are transported to the rapidly growing shoots, flowers, and leaves



FIG. 57. Diagram of a three-year-old stem of basswood cut so as to show the structure in cross-section, *C*, in radial section, *R*, and in tangential section, *T*; *c*, cortex; *ck*, cork; *mc*, medullary ray in cross-section; *mr*, ray in radial section; *mt*, ray in tangential section; *l*, lenticels; *cam*, cambium zone; *p*, pith.—H. O. Hanson.

each spring. Later in the season a lessened volume of water produces firmer and denser tissues. The age of a tree can generally be ascertained by counting the annual rings. However, two

rings may be formed in one season owing to the checking of the growth by fire, severe drouth, depredations of insects and the subsequent recovery and renewal of growth. These annual rings also reveal the life history of the tree, broad bands indicating favorable seasons and narrow bands telling of fires, drouth, transplanting, and other factors that limit the annual growth.

39. The Medullary Ray.—Another characteristic of most woody stems as seen in cross section is the series of delicate lines, the medullary rays, that radiate from the center of the stem (Fig. 57, *mc*). Some extend from the pith through the phloem and others originate in the various annual zones of the xylem and extend partially through the phloem. When magnified these medullary rays are seen to consist of rather thin walled, oblong cells and varying in width from one to a few cells (Figs. 56, *m*; 57, *mc*). In a longitudinal section cut parallel to the rays, called a radial section, they appear as bars of oblong



FIG. 58. Relation of the medullary rays to the xylem cells in pitch pine: *A*, radial section showing the elongated cells or tracheids, *t*, marked by circular, thin places on bordered pores, *p*; *m*, medullary ray of two cells accompanied by tracheids, *mt*. *B*, tangential section showing medullary rays one cell broad and three to nine high—*p*, bordered pores.

cells running at right angles to the elongated cells of the xylem (Fig. 57, *mr*), while in longitudinal sections cut at right angles to the rays, called a tangential section, we see that the cells of

the rays are arranged in a lens shaped group (Fig. 57, *mt*). The medullary rays are of great importance in the distribution and often also in the storage of foods. Their structure and arrangement in the stem are admirably adapted for this work. Their thin walls can readily withdraw from the ascending current in the xylem as much water as is needed for the phloem and cortex. Tracheids are often associated with the medullary rays to further increase their absorbing power (Fig. 58). Particularly is this true in the spring wood where the ray cells also often become greatly enlarged in order to absorb the large volume of water required at this time of year. In the same way the carbohydrates and albuminous substances transported through the phloem are withdrawn through the medullary cells and conveyed to the active cambium and the growing cells of xylem.

The value of oak and other woods for interior decoration is materially increased by the medullary rays. The trees are sawed into timber in such a way as to expose the medullary to the best advantage, a process called quartering (Fig. 59). Consequently



FIG. 59. Diagram showing a common method of sawing in producing quartered oak. The log is first squared to remove the sap wood and then a number of boards, depending upon the size and grain of the tree, are removed at *a* and *b*. These are the best boards obtained from the log, being large and approximately radial and therefore showing the medullary rays to good advantage. The remaining portions of the log are cut radially, as shown at *c*. One of the ends of quartered lumber is oblique and must be squared, as shown in the diagram.

the rays appear as flecks and bands upon the surface of the board and vary greatly in character owing to the angle of cutting and to the irregularities of the trunk. The medullary rays can always be detected in the grain of wood because they run across or at right angles to the xylem cells (Fig. 57, *mr*). As a rule

the wood cells of the stem remain alive for only a few years. This fact is usually apparent in any log or stump where it will be seen that the outer annular rings are of a lighter color, the sap wood, and the older parts are of a darker color, the heart wood. The sap wood is active in the transport of water and contains many living cells. . For this reason it is not so valuable for lumber. The heart wood is composed of dead cells and while the most serviceable for building purposes, it is of little use to the tree as we often see a vigorous tree whose heart wood has been largely destroyed by decay.

40. Other Types of Stems.—The cone bearing trees, such as the pines and spruces, have essentially the same arrangement of tissues and mode of growth as noted above in the dicotyledons. The xylem, however, consists entirely of tracheids (Fig. 58)



FIG. 60. Cross-section of a single vascular bundle of corn stem: *ph*, phloem; *x*, small cells of the xylem; *v*, vessels of xylem; *st*, stereome that forms a sheath about the bundle; *p*, parenchyma of the stem.

with the exception of a few small spiral vessels that are formed as the first cells of the vascular bundles.

The monocotyledons, plants distinguished usually by their parallel veined leaves and single seed leaf, like the palms, lilies, grasses, etc., are characterized by stems that do not increase materially in diameter. Growth is largely confined to the top of the stem and consequently it can only elongate, forming a very regular, columnar trunk. The reason of this is apparent when cross sections of such stems are examined (Fig. 45, *D*). The vascular bundles are more or less scattered throughout the stem and it is especially noteworthy that there is no cambium separating the xylem and phloem. It will be seen by examining Fig. 60, that the tissues of the bundles are of a similar character and have the same arrangement as in the dicotyledons (Fig. 41) save for the absence of the cambium which prevents the addition of new cells to the xylem and phloem. Consequently there can be no considerable increase in the diameter of the stem and no necessity therefore for the protective layer of cork cells.



FIG. 61. Longitudinal section of the tip of a growing stem: *e*, epidermis extending over surface of the entire tip; *a*, formative region; *b*, upper portion of the zone of elongation; *c*, cortex; *x*, cells of the central region that by further growth form the vascular bundles; *l*, first appearance of the leaves.

41. Apical Growth of the Stem.—All stems are characterized by a more or less extended apical growth. The changes produced in stems by the growth of the cambium only result in increasing their diameters but the elongation of all stems is

effected by the formation of new cells at the apex of the stem. Fig. 61 shows the general features to be noticed in the tip of a stem as seen in longitudinal section. New cells are being formed at the apex by division just as in the case of the root. Below this formative region is the zone of elongation in which the cells gradually change in character so that the cortical and



FIG. 62. Diagram of the tip of a stem as seen in longitudinal section: *a*, formative region corresponding to the part shown in Fig. 61; *l*, *b*, leaves and branches in various stages of development; *v*, vascular bundles; *c*, cortex; *p*, pith.

central regions are apparent and in certain elongated cells we note the first indication of the vascular bundles (Fig. 61, *x*). The leaves also originate in this region through the active division of the superficial cells of the stem. The branches of the stem develop somewhat later in the axils of the leaves, and more

deeply located cells in the cortex co-operate in their formation. This relationship of the leaves and branches to the stem is shown in Fig. 62, which is a diagram of an elongating stem, showing the relation of the apical region of the stem (*a* in Fig. 61), to the lower and older portion. The cells in these young leaves and branches by rapid division and growth soon form the characteristic tissues already noted in the leaves and stems; while a cor-



FIG. 63. Section through the trunk of basswood showing relation of branches to main stem. In the upper portion of the figure are three branches that were killed after a few years' growth by shading and that have been overgrown by the annual rings of the wood. The branch shown at *a* has remained alive and increased in size after the manner of the main stem.

responding growth in the cortex and adjacent regions results in the formation of vascular bundles that connect the vascular bundles of the stem with those of the leaves and branches. When a leaf falls off, the ends of these vascular bundles can be easily seen in the leaf scar (Fig. 28, *A*) but owing to their minuteness it is not an easy matter to trace them through the stem to the point where they join on to the bundles of the stem. In the branch, however, owing to its size and woody character, the union with the stem is very manifest. Fig. 63 shows several very small branches that continued to keep pace for one or more years with the growth of the stem, but eventually they were killed by the overhanging branches and in time became overgrown with the annual layers of xylem. If the branch is favor-

ably situated so that it continues to live then a cone-like structure is developed (Fig. 63, *a*) since the branch increases in girth each year in the same manner as the stem. These branches running through the xylem are the cause of knots that appear in lumber. If the branch is living the tissues of stem and branch are closely bound together and we have a solid knot, but it not infrequently happens that branches after living several years die and begin to decay before they are overgrown. Such a limb will produce in the lumber a knot that is often black and shaky or it may fall out, forming a knot hole. The lower branches of trees growing in a forest are quickly pruned off by the shade of the upper branches. This causes straight, clean trunks to develop and the lumber is free of knots.

It follows from what has been said above that in pruning trees the limbs should be cut off close to the trunk and the trunks should be painted to promote the healing of the wound and to exclude organisms that promote decay. When a wound is made that extends across the cambium as in the cutting off of a limb



FIG. 64. Section of a portion of a stem showing the healing of a wound caused by the cutting off of a lateral branch: *A*, formation of the callus, *cl*, owing to the renewed activity of the living cells exposed by the wound; *c*, cortex; *cm*, cambium; *x*, region of the xylem. *B*, a similar stem three years later—*c*, cortex; *cm*, cambium; *x'*, xylem added since the wound was made; *o*, position of the cambium at the time the branch was cut off; *x*, original xylem of the stem. *C*, trunk from which three branches have been removed, showing the gradual covering of the wounds by new tissue.

or by a cut into the truk that removes the tissues as far as the xylem it will be noticed that the exposed cambium is stimulated to form a mass of large cells, the callus, that gradually extend over the wound if this is not too large. While the callus is still in process of forming an outer protective layer of cork tissue is developed and also a new cambium layer is formed on the inner side of the callus which finally unites the old cambium, thus continuing the cambium of the stem over the surface of the wound (Fig. 64, *A*). In this way wounds are covered or healed and the tissue exposed by the cut is gradually buried deeper by the annual addition of new tissue derived from the cambium (Fig. 64, *B, C*). Grafting is made possible owing to the formation of callus that unites the tissues of the scion and stock (Fig. 65). While grafting is performed in a variety of ways the process consists essentially in all cases of bringing the cambium and cortex of the cutting or scion in contact with the corresponding



FIG. 65. A common method of grafting: *A*, insertion of two scions into cleft of stock at cambium region. *B*, wound protected with wax to prevent drying out of tissues.—After L. H. Bailey.

region of the stock. The junction of the scion with the stock is sealed with grafting wax to prevent decay and the drying out of the exposed cells while the callus is joining together the corresponding tissues in the two parts. When the wound is healed only a slight ring or swelling indicates the point where the scion was inserted—so complete is the union of the vascular and cortical regions. Strange to say, although the organic union is complete, each part, with very rare exceptions, continues a separate and distinct existence. For example, a scion from a plum

grafted upon a branch of a peach tree will develop a normal shoot bearing plums and the peach tree will also continue its growth unaffected. Ordinarily only closely related forms may be grafted, as tomatoes and potatoes, apples and quinces, etc.

42. Comparison of Stem and Root.—It will be noticed that the structure of the growing tip of the stem presents several striking departures from that of the root. In the first place a protective structure such as the root cap is not required (Fig. 61). The leaves, and consequently the branches are formed with great regularity from the superficial tissues of the stem, whereas the rootlets are deep seated in origin and irregularly distributed. Furthermore, the elongation of the stem is distributed over several centimeters, while in the root it is confined to a few millimeters. The leaves and branches are distributed over the entire zone of elongation, while lateral roots do not appear until elongation has ceased. This prolonged growth of the tip of stems is of great advantage in bringing about the proper adjustment of leaves and branches. These organs come into sharp competition with each other as well as with other plants, and so long as growth continues there is a possibility for them to become adjusted to the light without interfering with one another. The mode of elongation of stems and roots is essentially alike. Each has a formative region at the apex characterized by active cell formation and slight elongation, back of this is the zone of rapid elongation and following this is the zone in which elongation is gradually ceasing and the tissues are assuming their characteristic forms and functions. It is also important to note that the growth of the cells on one side of the stem or root is faster at a certain time than any of the other cells. This will cause the stem to be bent towards the opposite side. This condition of rapid elongation does not continue on one side for any considerable time but slowly moves from point to point around the stem so that the stem is successively bent from side to side, thus causing the apex to travel in an irregular circle. By these constant movements, nutations, the stems and roots are more fully exposed to the various forces that influence their development and conse-

quently they adjust themselves to the best advantage (see page 29).

As a rule the irritability or sensitiveness of the stem is not so localized as in the case of the root. It is evident that it is an advantage to have the entire growing surface of the stem sensitive to all those stimuli that influence its growth, because by this means it will be so influenced as to bring the leaves and branches into proper relations with the surroundings. In some plants to be sure the sensitiveness to stimuli is quite as localized as in the root; as, for example, the tip of the cotyledons of many grasses which have an appreciation of light almost the equal of our eyes. These cotyledons will turn toward a light too feeble to enable us to read the time of day on a watch. The significant fact to be noted here is that this keenness and localization directs the plant into a suitable place as soon as it emerges from the soil. Grass plants grow in thick colonies and therefore the stems come into sharp competition. Consequently it is a necessity that the young shoot on emerging from the soil should be able to appreciate the feeblest light and so be able to direct its growth through the smallest openings.

The reactions of the stem to various stimuli, such as light, gravity, etc., are quite as purposive as those noted in the root. Lateral light, as in the case of plants growing in windows, acts as a stimulus and causes a more considerable growth of the cells on the darker sides of the stem. This results in bending the stems towards the light and consequently in the exposure of the broad blades of the leaves to the light. So, also, when erect stems are placed in a horizontal position the stimulus of gravity awakens a more active growth on the underside of the stem, thus bending the tip into an erect position for securing light. Not all stems and, indeed, not all parts of the same plant body are similarly influenced by these forces. Many plants tend to arrange their stems at right angles to the action of gravity and light, as in the case of creeping and underground stems. So also the lateral branches are inclined at various angles to the stem. These adjustments are not to be looked upon solely as the direct response to one or more stimuli but as the resultant of numerous interac-

tions aroused in the plant by various stimuli. The entire plant responds as a unit to these forces. Stimulation and the consequent growth in one part has its effects upon all other parts. As a result of this interaction there comes about a correlation in the growth of all the parts that is expressed in the symmetry of the plant body and in the perfect adjustment of roots, stem and branches, and leaves to its needs. Any window garden, forest, or field furnish abundant evidence of the complicated and varied nature of these reactions. An illustration of this is afforded in the removal of the terminal axis of a tree. An adjoining lateral branch that heretofore has been stimulated to a more or less horizontal growth, now is so directed in its development that it ultimately becomes the terminal axis of the tree. So some parts are constantly being overshadowed and choked out while others are directed into favorable positions and perform the work of the destroyed parts. In this way there is brought about that weaving and twisting of stems and branches and the adjustment of leaves that so often suggest a conscious effort to reach favorable positions.

43. Modifications of the Stem.—In the above discussion attention has been called to the more common characteristics of the stem. It is well to remember, however, that variation is the law in the plant world. No two plants are alike. The environment of the plant is constantly stimulating it and causing it to vary. Frequently the variations are so minute as to escape our attention or totally distinct forms may arise. These departures from the parent type may be of no advantage to the plant and it may therefore perish. On the other hand the variations may be of such a character as to enable the plant to more successfully compete with other plants and as a consequence the plant with its helpful variations will survive. So it has come about that these changes going on during the past ages have resulted in many remarkable modifications. A few of the modifications of the stem that are of especial advantage to the plant and therefore spoken of as adaptive variations will be considered in the following paragraphs.

44. The Climbing Type of Stems.—Climbing stems are char-

acterized by being rather small and having greatly elongated internodes. Stems grown in the dark show a similar development. Perhaps this variation that we call the climbing type has been brought about by competition with larger plants. The feeble light has stimulated these stems so that they attain a very extended growth and finally are able to reach the light and display their leaves. Many variations in the structure and sensitiveness of the stem are associated with this elongation, all of which are designed in one way or another to enable the stem to reach the light. One type of these variations is seen in twining stems, as the morning glory, bean, hop, etc. Young twining plants behave quite like the ordinary plant. The stems are erect and actively nutating, the apex traveling through a rather large circle in one to three hours. When a certain height has been reached the stems are stimulated by gravity so that their upper portions grow more or less horizontally (Fig. 66). This position is a decided advantage since the stem is now revolved through a larger circle and has a greater chance of coming into contact with an object about which it can twine. As soon as the stem comes into contact with any support its nutation will cause it to wind or twine about it. Doubtless the contact also acts as a stimulus, causing the stem to bend more energetically. Certainly in many plants the size and roughness of the support are important factors in inducing the twining. At first the coils are merely horizontal but owing to the elongation of the stem these coils are gradually pushed upwards and become steep and very firmly bound around the support. Twisting of the stems and reflexed bristles often assist in anchoring the plant to its support. It is interesting to note that in the majority of cases the stem twines about the sup-

FIG. 66. Twining habit of wild bean. Note the horizontal position of the upper portion of the stem which results in the apex mutating through a wider circle.

port in left hand spirals or clock wise; less frequently right hand spirals are formed, as in the hop, some honeysuckles and knot weeds. Why the growth of the stem results in left or right hand twining is not known but the fact remains that the plants can not be induced to change their method of twining.

45. The Tendril Type of Stem.—One of the most interesting variations of climbing plants is seen in those stems that climb by means of tendrils. In this type almost any part of the plant may become modified so as to act as a tendril and bind the plant to a support; *e. g.,* the petioles of nasturtium and clematis, stipules in smilax, midrib of leaf in many members of the bean family, or modified branches as in the grape vine, etc. The ordinary tendril as we see them in the gourd, passion flower, etc., are highly specialized stems that are very sensitive to touch. These organs are rapidly nutating, frequently completing two or more circles in an hour, and as they approach maturity they often become slightly curved and hooked (Fig. 67, *A*). Their sensitiveness to contact, which is often localized on the concave upper third of the tendril, also becomes so keen in some forms that a thread weighing one-fiftieth of a gram will cause a curvature of the tendril when placed upon it. In fact a much smaller weight if caused to vibrate will act as a stimulus, so that we have here an illustration of sensitiveness to weight that is keener than our own. When the sensitive part of a tendril is stroked by a pencil or comes in contact with any hard object, this acts as a stimulus and causes a very rapid growth upon the convex side of the tendril. As a result of this growth the tip of the tendril curves, forming a complete loop in from a few seconds to a few minutes. The contact with any object acts as a continued stimulus and the tendril soon becomes tightly wound around it. As soon as the tip is firmly bound to a support a new impulse is frequently transmitted to the free portions of the tendril which in a day or so begin to assume the form of coils that are often reversed in the middle (Fig. 67, *B*). By this unique device tendril bearing stems are firmly fastened to their supports but at the same time the coiling of the tendrils acts like a spring permitting considerable disturbance of the

vine without danger of breaking. This same device is copied in making the attachment of the wires to the telephones and other pieces of apparatus. From many standpoints the tendril bearing stem is an admirable one. It is evident that such stems can reach the light more directly and economically than twiners. This doubtless accounts for their greater abundance and common occurrence. It is also important to note that the tendrils are

Fig. 67.

Fig. 68.

Fig. 67. Tendrils of the bur cucumber: *A*, hooked tendrils in receptive state. *B*, apical portion of tendril coiled about a branch and the remaining portion of the tendril forming reversed coils, thus drawing the vine to the supporting branch.—H. O. Hanson.

Fig. 68. Branch of Japanese ivy attached to wall by means of tendrils with adhesive discs.—H. O. Hanson.

especially abundant at the end of the branches where they reach out considerably beyond the young leaves. Owing to the swaying of the long free shoot of the climber in the wind and the nutations of the tendril and shoot these sensitive organs are brought into contact with branch after branch and thus the vine

is led up to the crown of the tree or to the top of the vegetation. The Virginia creeper, Boston or Japanese ivy, etc., have peculiar tendrils that grow away from the light and form adhesive discs, instead of coils, that bind the vine to the support (Fig. 68). The tip of these tendrils becomes greatly modified so that they fit into every groove and crevice of the substratum. Indeed in some vines these discs become so thoroughly cemented to the crevices and irregularities of the support as to resemble bits of melted wax. It is very interesting to watch the results of the stimulation induced by one of these tendrils striking a wall or other support. The formation and cementing of the disc are accomplished in about four days in the case of the Virginia creeper. Some plants like the poison and English ivy and many tropical plants climb by means of roots. These organs, like the disc forming tendril, avoid the light and seek the dark nooks and corners in the bark of trees or rocks over which the plants grow.

46. Prostrate and Creeping Stems.—It is interesting to note that a large number of plants having rather small and weak stems like the climbers have never acquired the habit of climbing. The young shoots of these plants are often erect at first but soon the weak stems bend over and become prostrate. While these procumbent and creeping stems are at a disadvantage in that they are obliged to arrange their leaves in one plane, on the other hand they are not obliged to build up the elaborate structures that are necessary for the support of the erect stems. Consequently plants of this class are of common occurrence in old fields and unfertile soils since they can subsist on a meager supply of food. They are also protected against injury and therefore adapted to wind swept sandy plains, to rocky hills and mountains, and to districts visited by heavy snows. The majority of these plants also have a very decided advantage owing to the fact that certain nodes form roots and buds that develop into new plants. In some cases, stolons, only the terminal bud develops in this way and with the dying off of the old stem a new plant is established, as in the houseleek (Fig. 69). In other cases, runners, almost any node may form buds and roots which

become separate plants by the decay of the old stem, as in the strawberry, gill-over-the-ground, cinque foil, etc. These new plants repeat the mode of growth of the parent plant and

FIG. 69. House leek forming buds at the end of short branches or stolons.

in this way many prostate stems spread out over the soil, establishing new plants in wider and wider circles. If you will observe the number of new plants established and the distances traversed by some of these creeping stems each year you will see the reason for the common occurrence of the large mats and colonies of plants with prostrate stems.

47. The Rhizome Type of Stems.—In a third type, the stems have become so modified that they respond to the various stimuli in quite a different way from the forms noted above. They creep along in the soil and frequently resemble roots more than stems. For this reason they are called rootstocks or rhizomes. They are, however, real stems, as is attested by the numerous leaves and often erect branches that spring from their nodes as seen in the ferns, sweet flag, cat tail, grasses, etc. Plants of this type are well protected against drought, cold, and fires, and like prostrate stems they are adapted to establishing new plants as is apparent in grassy meadows, colonies of golden-rod and daisies or reedy banks of cat tails, sedges, etc. These stems are also well adapted to propagating new plants because they generally serve as storage organs for food and are therefore often of a fleshy character (Fig. 70). For these reasons it is sometimes very difficult to eradicate plants of this type. This is very well illustrated in the quack grass, often a troublesome pest in cultivated land. Plowing and hoeing only serves to break up the rhizome into numerous parts each of which may

develop buds and roots from their nodes and so establish new plants.

48. The Condensed Type of Stems.—In many cases we find that the food is localized in special regions of the rhizome which consequently become enlarged and rather fleshy. Such modified parts of a rhizome are called tubers, *e. g.*, the potato and Jerusalem



FIG. 70.



FIG. 71.

FIG. 70. Rhizome of Solomon's seal with aërial shoot just emerging from the ground. The seal-like scars mark the successive shoots produced during the past three years.

FIG. 71. Formation of tubers: *A*, old potato or tuber with two shoots reaching up into the air and from the base of these shoots, rhizomes have been formed that are developing new tubers. *B*, mature tuber with spirally arranged buds, the so-called "eyes" of the potato.

artichoke (Fig. 71). The potato is formed by the storage of foods in certain parts of the rather small rhizomes that branch out from the stem of the plant. The "eyes" are the buds that develop at each node of the rhizome and each is capable of forming a shoot, although but a few of them so function, as is the case in an ordinary branch of a tree. By passing a thread around a potato so that it touches each successive bud you will see that there is the same arrangement of the buds as appears in the leafy stem.

A great variety of plants develop only very short stems that are more or less buried in the ground. This type of stem modification is of great advantage in many ways. The vital parts of such plants are only slightly exposed and consequently suffer little from grazing animals or other sources of injury, as in

plantains, dandelions, etc. Frequently these short stems are associated with an abundance of foods that are stored in roots or other organs. Such plants can quickly send up and mature their flower stalks or leafy stems and thus avoid unfavorable conditions, as drought, competition with larger vegetation that will appear later, etc. This habit has been made very conspicuous by cultivation in many plants, as the carrot, turnip, radish, and beet. But in nature the very short, almost flat stems and fleshy roots are also of common occurrence, *e. g.*, the wild carrot, wood betony, dandelion, etc. In many cases the short stem itself is the storage organ and consequently it becomes en-



FIG. 72. Shortened types of stems: *A*, corm of jack-in-the-pulpit. At left surface view showing lateral buds, roots and sheathing leaf arising from top of shortened stem. At right sectional view with folded leaf, *l*, in bud at apex of stem. *B*, bulb type of shortened stems. At left bulb of onion showing the ensheathing leaves which are swollen at their bases with food, thus forming the bulb. At right, section of a bulb of hyacinth showing the fleshy leaves attached to the very short stem and in the center of the bulb a flower cluster.

larged and fleshy, as in the Jack-in-the-pulpit, garden crocus, spring beauty, etc. (Fig. 72, *A*). These short erect stems are termed corms and are suggestive of the tuber. In other plants the bases of leaves attached to the short stems function for the storage of food, as in the bulb type of stems of lilies, onions, etc. (Fig. 72, *B*).

49. The Bud Type of Stems.—The final variation of the stem to which attention can be called at this time is termed the bud type. The bud, as in the case of the bulb, consists of a very short stem which bears a number of more or less modified leaves (Fig. 73, *A*). The outer leaves of the bud are usually scale-like and leathery and closely overlapping to protect the delicate parts within. They are often further assisted in this work by mucilaginous coatings and hairs, as in the horse chestnut, poplars, birches, walnuts, etc. Within the scale-like leaves are the ordinary green leaves, branches and flowers in various stages of development which will appear with the opening of the buds in the spring. This type of stem was doubtless induced by the change of climate that followed the formation of the mountain ranges and the change of the earth's orbit about the sun. As soon as the tropical climate on the earth was disturbed by cold currents of air that swept over the earth as a result of these changes, there resulted a stunting or killing of young and imperfectly developed organs. These changes in temperature were factors in producing modifications of the stem. Thus the bud type came about with its closely united scales which protect the delicate parts within against the drying winds which would rob them of moisture at a time when none could be obtained from the soil. Possibly also the buds prevent the injury that would come from a too sudden freezing or thawing of the delicate parts within. The rapid unfolding of the bud in the spring is a constant source of surprise but when we recall that it contains usually an abundant supply of food and practically all the organs that will appear on the stem during the season we can understand how the bud opens and elongates into a shoot bearing leaves and flowers during the first few weeks of spring. The part of the stem bearing the scale leaves does not elongate ma-

FIG. 73. Bud type of stem: *A*, branch of hickory showing a large terminal bud and several small lateral buds that were developed in the axils of the leaves of the past season. The branch is three years old, as shown by the three rings, *r*, *r'*, *r''*, of the bud scars which mark the successive positions of the terminal bud during the past three seasons; *l*, leaf scars. *B*, branch of pear—after Bailey. The ring at 1 shows the position of a bud which produced the next year a pear, as is indicated by the large scar and swollen branch at *a*. A short lateral branch was also developed the same season with its terminal bud at 2. The next year this bud produced a branch that extended to 3, bearing several leaves (note leaf scars) and of course axillary buds—one of which, *b*, we see grew in the following seasons. No fruit was developed this season. The bud formed at 3 behaved the next season very much as the one previously noted at 1, forming a pear at *a'* and a short lateral branch that reached to 4. Note also that the axillary bud, *b*, of the previous season grew a little and its subsequent history can be followed by the annual bud scars. In the following season bud 4 developed a vigorous shoot reaching to 5. No fruit was formed but three buds survived, *b'*, *b''*, *b'''*, thus nearly duplicating the growth of bud 2. Bud 5 develops the next season fruit at *a''* and also two lateral shoots that extend to 6 and 7 respectively. The three buds below 5 had only a feeble growth during this and the succeeding seasons and bud *b'''* evidently perished during its first period of growth. Bud 6

terially since these leaves are of service only during the winter, and soon fall off after the opening of the bud. Consequently the scars formed by the fall of these protective scales form a ring (Fig. 73, *r*) each year about the stem which marks the position of each successive bud. By counting the number of these rings on a twig you can ascertain its age and observe the extent of elongation during each season, just as the age and increase in diameter is revealed by the annual rings of the wood. From the twigs and branches may be read the life history of the plant, the nature of the season, the competition experienced and the fruit formed (Fig. 73, *B*). Buds are usually found singly in the axils of the leaves and as a rule only a few that are favorably situated as regards the light ever develop. The development of the buds brings about the characteristic appearance or habit of the plant. This growth is not controlled alone by the favorable position of the buds but also by the interaction or correlation of all the growing parts. As a result in some trees we find a single terminal bud that is larger and better developed than all others. This bud will produce the longest shoots and consequently such trees will have spire-like stems as in the spruces, larches, etc. In other cases this same type of growth may continue for a time but eventually terminal buds of equal vigor will be developed upon several branches and consequently equal growths or diffuse types of branching will result as in the maples, elms, etc. Buds that are not favorably located generally perish

produces the next year a pear at a''' and a lateral shoot reaching to 8 and bud 7 at the same time only develops a leafy shoot reaching to 9. Buds 8 and 9 both produce fruit the next season (a^{IV}, a^{VI}) and short lateral branches reaching to 10 and 11. These buds during the present season only formed leafy shoots.

Buds are formed in the axils of all the leaves that appear on the shoots of a season's growth, but notice that but few or none at all develop. Some have a thrifty growth, others remain small and perish after one or more years. This is true of the fruit. Some of the pears mature, as is indicated by size of the scars, while others drop off after one or more months. Note also the almost regular alternation in the production of shoots bearing fruit and leaf shoots. Bud 6 is the only exception to this succession. Observe that the branches assume different positions and that the extent of the elongation of the shoot from year to year varies. Can you explain these facts?

after a few years but not infrequently they remain alive and become overgrown by the increase of the stem. In such cases the bud grows slowly and maintains itself near the surface of the wood. These buds growing and often branching in the wood produce those curlings and twistings in the grain that are commonly known as bird's-eye wood. Dormant buds sometimes develop into normal branches, as when a portion of a tree is removed and consequently an additional supply of food stimulates them to growth. Buds may also arise upon any part of the stem, root or leaf. These are the so-called adventitious buds and their formation upon roots often accounts for the colonial habit of many trees, plants and shrubs, as in the poplars, Ailanthus, sumacs, etc. They appear to rise naturally in some cases, but in other instances their formation is due to the stimulus of a wound or some other cause and like the dormant bud they serve to prolong the life of the plant. Richards has shown that when living cells are exposed to the oxygen of the air, owing to a wound, that these cells are stimulated as a result to renewed activity. This doubtless explains the formation of callus and the healing of wounds, as well as the formation of adventitious buds in many cases. Common examples of buds due to wounds are seen in the vigorous shoots that spring up from the stumps of hard wood trees, pollarded willows, etc. Some buds become fleshy owing to the storage of the food and dropping from the plants serve to propagate new individuals. Examples of this are seen in the fleshy buds on the tips of the branches or in the axils of the leaves of the stone-crop and some lilies and in the flower clusters of some onions. Many aquatic plants have the habit of forming similar buds on the approach of winter. These being compact and heavy with food sink to the bottom of the ponds in the fall and renew their growth in the spring.

CHAPTER IV

THE FLOWER, FRUIT AND SEEDLING

50. The Structure of the Flower.—All the energy of the plant is directed towards the perpetuation of its kind. Numerous examples have been noted where this is accomplished by bulbs, rootstocks, buds, etc. Among our common plants, however, new individuals are usually produced from seeds that are formed in the flowers. We will first examine the structure of a flower and see how it is adapted to the performance of this important work. A flower is a highly modified stem of the bud type. This is very noticeable before the flower opens or is in bloom (Fig. 74, *A*). A series of leaves usually of a greenish color envelop the delicate parts within and protect them, as in the case of the bud, against drying winds. These green leaves are known as the calyx and each individual leaf is called a sepal. As the bud opens a number of organs are disclosed, particularly noticeable are a set of more delicate and variously colored leaves the corolla, each leaf of which is called a petal (Fig. 74, *B*). The corolla is of service to the flower in many ways. Like the calyx it may assist in protecting the other organs. Especially is it of importance in guarding them against dews and rains. For this reason the calyx and corolla collectively called the floral envelop or perianth, in many flowers often close at night or on rainy days. It will be seen later in the work that the perianth is of the greatest importance in assisting insects to visit the flowers in such a way that seeds may be produced. Within the perianth are two kinds of peculiar organs, the central flask-like ones are called pistils or carpels and surrounding them are the stamens (Fig. 74, *B*). These two kinds of organs are collectively known as sporophylls since their special work is to produce certain cells called spores. The sporophylls are the important organs of the flower. The calyx and corolla may be lacking but no seed can be formed without the sporophylls.

51. The Structure of the Sporophylls.—Let us examine the structure of these organs and see how they co-operate in the formation of the seed. The stamens consist of a lobed sack or



FIG. 74. Flower of the stonecrop, *Sedum: A,* bud stage of flower—*ca,* calyx showing three sepals. *B,* open flower—*p,* petals of corolla; *s,* stamens; *c,* pistils or carpels. *C,* stamen, consisting of a four-lobed anther supported on delicate stalk or filament. *D,* pistil or carpel—*o,* ovary; *s,* style which terminates in a small knob or stigma. *E,* longitudinal section of ovary showing row of ovules attached to wall of ovary. *F,* cross-section of ovary, ovules in two rows.

anther which is usually supported upon a stalk or filament (Fig. 74, *C*). If a young anther is cut across four cavities, containing minute or dust-like grains, the microspores, will be seen (Fig. 75, *A*). The microspores, also called pollen spores or grains, are minute cells, provided with a cell wall often variously sculptured and contain a nucleus and protoplasm like an ordinary cell (Fig. 79). At maturity the anthers break open in a manner that varies in different plants exposing the microspores to the air (Fig. 75, *B*). The pistil or carpel is quite different in character. At or near the top is a more or less modified part,

the stigma, below which is an elongated stalk, the style, that broadens out into a base or ovary. By longitudinal and transverse sections of a pistil we gain a better idea of its real character (Fig. 74, *D–F*). The ovary is now seen to contain a cavity from the walls of which one or more rather spherical bodies, the ovules, are developed. The structure of the ovule is very complicated. If a thin section is made through the center of one of them it will be seen to contain usually but one large spore, therefore called the megaspore (Fig. 76, *mg*). This spore is



FIG. 75. Structure of the anther: *A*, diagram of the anther cut across, showing the four cavities, sporangia (sing. sporangium), filled with spores. *B*, cross-section of a mature anther. The tissue about the spores has broken down, thus forming two cavities and at the left the breaking open of the wall of the anther is shown.

developed in a mass of tissue, the nucellus, and the whole is surrounded by one or two coats, the integument, which grows up from the base of the ovule but never quite surrounds it, thus leaving a small opening or micropyle (Fig. 76, *mi*). The megaspore, unlike the microspore, is never set free but is retained permanently in the ovule.

It is not known how the sporophylls have originated but there are indications that they may have arisen through the modification of leaves for the purpose of forming spores instead of constructing foods. It is for this reason that these organs are called sporophylls, or spore-forming leaves, from phylon, a leaf. Those forming small spores are sometimes called microsporophylls instead of stamens and those forming large spores megasporophylls instead of pistils or carpels. Very many modifications of the sporophylls will be noted later in the work.

52. The Germination of the Microspore and Megaspore.—The question now naturally arises as to the meaning of these peculiar

structures. The important part of the sporophyll is the spore. A spore is a cell that is capable of germinating or growing under favorable conditions and thus producing some kind of a new organism, or we may say a new plant. You are always to think of a spore as having this power. In case of the spores under consideration very rudimentary plants are produced that consist of only a few cells. These minute plants are very important,



Fig. 76.



Fig. 77.

Fig. 76. Section of the ovule of the Canada lily (greatly magnified): *mi,* micropyle; *i,* integuments; *n,* nucellus, here consisting of a single layer of cells, but it often forms a large mass of cells; *mg,* megaspore with large nucleus in center. This spore is about to divide; *f,* stalk or funiculus which attaches the ovule to the wall of the ovary.

Fig. 77. Germination of the megaspore: *A,* first division of the nucleus of the megaspore. *B,* second stage in the germination, four nuclei being formed. *C,* final stage in the division of the nuclei.

however, because they contain the male and female cells, or gametes, so called because they unite, thus forming a new spore, the gametospore, quite different from the micro- and mega-spores. So let us not lose sight of the fact that these apparently insignificant micro- and mega-spores germinate and produce minute plants that are quite distinct from the plants that bore them. We will first follow the germination of the megaspore

and note the character of the plant that develops from it. As has been stated this spore is never discharged from the ovule within which its entire growth occurs. The first indications of germination are seen in the enlargement of the spore and the division of its nucleus into two nuclei, which separate and move to either end of the spore (Fig. 77, *A*). The food for the nourishment of the growing megaspore is derived from the cells of the nucellus which are dissolved and absorbed as the megaspore increases in size. Each of the two nuclei at the end of the spore divides again, thus forming two nuclei at either end. This process is repeated once more so that we now find four nuclei at the ends of the spore which has become somewhat elongated (Fig. 77, *B, C*). Now a nucleus from each end, the so-called polar nuclei, move toward the center of the spore and fuse, forming one nucleus. This completes the germination of the megaspore and we see that it has grown into a sac-like plant consisting of seven cells, or nuclei imbedded in a mass of protoplasm (Fig. 78). The cells of this minute plant differ in character and perform very different duties. For example, the larger cell toward the micropylar end of the plant is the female cell or egg cell. It will be called the female gamete because it unites with another cell, the male gamete. The two small cells near the female gamete are the synergidae or cells that assist in the nourishment and later development of the female gamete. The two cells fusing in the center of the plant form the endosperm nucleus which later by repeated division produces a mass of nourishing cells that fill the space within the sac. The three cells at the bottom of the sac, known as the antipodal cells, may increase in number and take part in the later growth, but as a rule they are disorganized and used for food. They are often provided with cell walls, the other cells consisting only of a nucleus and a varying amount of cytoplasm. This plant formed by the germination and growth of the megaspore is called the female gametophyte, because it contains the female gamete, and you will see that a corresponding plant appears in the ferns and other groups of plants.

Returning now to the microspore, we find that its germination

also often begins while in its sporangium but in other cases growth begins after it has escaped from the sporangium and has been carried by the wind or some insect visiting the flower for

FIG. 78. The germination of the megaspore complete. The plant thus formed consists of a female gamete or egg cell, ♀, below which are two synergidae and in the center are the two uniting polar nuclei, *p*, while at the opposite end of the sac are three antipodal cells, *a; mi,* micropyle; *i,* integuments; *f,* stalk or funiculus in which a vascular bundle, *v,* has been developed to transport foods from the ovary to the ovule.

food, to the stigma of the pistil. The stigma is admirably adapted to hold the microspore, being provided with minute outgrowths, and frequently sugary solutions are exuded which fasten the spores to the stigma. The real importance of these sugary solutions is to nourish the microspores. It is evident that these minute dust-like cells can not contain sufficient food to bring about any considerable growth. It is noteworthy that

the strength and nature of the sugar solutions varies in different stigmas and that the microspores are often able to grow only in solutions of such strength as are found on the stigmas of their own kind of plant. So it results that the microspores are often unable to grow when carried to the stigmas of a different kind of plant from themselves. It will also be noted later in the work that the flower may be so developed that the microspore cannot be transferred to the stigma of its own flower but only to that of another. This is brought about in various ways, as by the shedding of the microspores before the stigmas are mature, etc. This transfer of the microspores of one flower to the stigmas of another flower is called crossing, the significance of which will appear directly. The germination of the microspore results in the division of its nucleus and the formation of a large and a small cell known as the tube and antheridial cells respectively (Fig. 79, *A, B*). The latter cell divides once, forming two cells, called the male gametes (Fig. 79, *C*). This stage in the germination of the microspore is reached either in its sporangium or after being transferred to the stigma. Nourished by the foods of the stigma the microspore continues its growth, the tube cell ruptures the outer wall of the spore and forms a tube-like growth, This tube cell grows down between the cells of the style into the cavity of the ovary where it usually curves out and enters the ovule by way of the micropyle. It now works its way through the tissues of the nucellus to the female gametophyte, into the cavity of which it enters after dissolving the wall. In the meantime the two male gametes have moved down to the end of the tube cell as it enters the female gametophyte (Fig. 79, *D*). The question will naturally be asked, what directs the peculiar growth of this minute plant? In the first place the tube cell is repelled by the oxygen of the atmosphere, so as soon as it appears, it is directed away from the atmosphere into the stigma. Its course down the style to the ovary is very largely controlled by lines of loose tissue through which it can easily work its way and also by foods which are deposited in the cells just ahead of the tube cell so that it is led along rather straight pathways to the ovary. The curving out of the tube to the micropyle and its subsequent

growth down into the female gametophyte can only be accounted for by some chemical stimulus that is located in the ovule and female gametophyte. A similar attraction brings about a move-



FIG. 79. Germination of the microspore: *A*, mature microspore. *B*, first stage in its germination—*t*, tube cell; *a*, antheridial cell. *C*, end of cell division—*t*, tube cell; *a*, antheridial cell forming two male gametes. *D*, diagram showing the formation of the tube which grows through the style and finally reaches the female gametophyte. The two male gametes, *g*, are shown passing down the tube; *t*, tube nucleus.

ment of the cytoplasm which carries the male gametes to the end of the tube cell. It will be seen that this peculiar plant (Fig. 79, *D*) consisting of a long tubular growth with three naked cells is more rudimentary than the plant derived from the megaspore. Because this plant produces the male cells or male gametes we call it the male gametophyte. So we see that the flower forms two kinds of spores and that these germinate and form two kinds of plants, a male and a female.

53. Fecundation or Fertilization.—As soon as the development of these two plants is completed a very remarkable change takes place which starts an entirely new growth that is quite independent of the development of the male and female gametophyte. The end of the tube cell becomes distended through the accumulation of material and finally ruptures, discharging the male gametes into the female gametophyte. The chemical composition of the female gamete is such that one of the male cells is attracted to it (Fig. 80) and finally the two gametes unite

forming a single cell. This body is a sexually formed spore or gametospore and like the ordinary spore has the power to germinate and produce a plant. The microspores and megaspores,

FIG. 80. The micropylar end of an ovule of Canada lily (sectional view), showing the process of fertilization or fecundation. The tube, *t*, has grown into the female gametophyte and ruptured, discharging the two male gametes. One, ♂, is seen fusing with the female gamete, ♀, and the other one, ♂′, is uniting with the two polar cells, thus forming the endosperm nucleus; *s*, one of the synergids; *i*, integuments.

however, were not formed by the union of sexual cells or gametes. Therefore we may distinguish them as asexual spores or simply as spores. It has frequently been observed that the endosperm nucleus also attracts the other male gamete and causes a similar fusion with it (Fig. 80). The union of the male and female gametes is called fertilization or fecundation. Unless fertilization is effected the growth outlined above practically ends the history of the flower. But if fecundation is effected then a gametospore is formed that is capable of germinating and producing a plant whose growth and development are attained with most remarkable changes of the ovule and often of the surrounding parts.

54. The Germination of the Gametospore and the Formation of Seed.—The first indication of the germination and growth of the gametospore is seen directly after fertilization. It becomes



FIG. 81. Germination of the gametospore of peppergrass, *Lepidium*: *A*, micropylar end of the embryo sac showing that the gametospore, *g* (fertilized female gamete), has developed a cell wall and become attached to the wall of the embryo sac. Compare Fig. 80. *B*, later stage in the germination—*e*, embryo cell; *s*, suspensory cells; *en*, endosperm cells. *C*, the embryo cell has formed two cells. *D*, later stage. The embryo cell by further division has formed a spherical mass of cells, here shown in section. Note the appearance of an outer layer of cells, the epidermis, and a central or stem region; *s*, a few of the suspensory cells. *E*, still later growth. Two growing regions, the cotyledons are appearing on the side of the stem. *F*, the micropylar end of the embryo sac, showing the stage of development where the parts of a small plant can be clearly recognized —*c*, cotyledons; *st*, stem ending in root, *r*, to which is still attached the suspensory cells; *en*, endosperm cells, now provided with walls, are being absorbed as the plant or embryo enlarges. Note that the embryo and embryo sac are slightly bent to the left. This curvature is due to the fact that the ovule in a great many plants becomes curved in its development with the result that the embryo sac assumes a curved or U-shaped form. This form of the ovule results in a complete bending over of the cotyledons against the stem, as shown in the next figure.

surrounded by a cell wall and attached to one side of the female gametophyte, which is henceforth called the embryo sac (Fig. 81, *A*). By a series of divisions the gametospore now forms a number of more or less elongated cells, the so-called suspensor, and a terminal cell or embryo cell (Fig. 81, *B, C*). In the meantime the endosperm cell has divided again and again and the resulting naked nuclei have become arranged around the sides of the embryo sac (Fig. 81, *B, en*). This formation of endosperm cells continues until they more or less fill the embryo sac and they also become surrounded with cell walls. These endosperm cells are filled with food and serve to nourish the germinating gametospore just as the nucellus nourished the female gametophyte. While the suspensory cells usually soon cease to grow, the embryo cell divides very actively, as shown in Fig. 81, *C–E,* and soon there is a clear indication of a minute plant with root, stem and leaves (Fig. 81, *F*). This minute plant developed by the germination of the gametospore, is called the embryo. In the case of the plant under consideration the embryo, when fully formed, consists of a stem with two laterally placed leaves, the cotyledons and a root. The region of the stem above the attachment of the cotyledons is known as the epicotyl and frequently assumes the form of a minute bud, the plumule. The region of the stem below the cotyledons is termed its hypocotyl and the root and root cap appear at its lower end (Fig. 82, *A*). The ovule containing the mature embryo is called a seed. The interesting feature about the seed is the fact that it is a structure in which growth has ceased and that it is capable, owing to a remarkable series of devices that will be noted directly, of remaining in this condition, *i. e.,* dormant often for one or more years.

Great variation characterizes the germination of the gametospore in the different groups of plants, but the development outlined above is fairly characteristic of those plants that form an embryo with two cotyledons. Such plants are called for this reason dicotyledons. Other plants develop an embryo with but one cotyledon, as our grasses, lilies, etc., and for this reason they

are called monocotyledons. Further variation will be seen in cone-bearing trees, ferns, etc.

It must be borne in mind that the formation of the seed is also attended with profound changes in the structure of the ovule, and often in the pistil and surrounding parts. Frequently the endosperm grows so extensively as to absorb and replace the cells of the nucellus and thus comes to occupy all the space within



FIG. 82. Structure of the seed: *A*, section of an ovule of peppergrass in which the growth of the embryo is nearly complete—*f*, stalk or funiculus attaching ovule to wall of ovary; *mi*, micropyle; *i*, integument; *en*, remains of endosperm. The embryo consists of a stem which is differentiated into a hypocotyl, *hy*, that ends below in a root, *r*, and root cap, and above in an epicotyl (the plumule), *pl*, and two cotyledons, *c*, which curve over and lie one upon the other. *v*, vascular bundles which extend up through the stem into the cotyledons where they subdivide, forming a network of veins. *B*, section of seed of water lily, after Conard—*e*, embryo surrounded by a layer of endosperm cells; *mg*, cells of the nucellus; *i*, integument.

the coats of the integument, as in the castor bean, morning glory, onion, etc. (Fig. 85, *B*). The embryo may remain comparatively small as in the cases just cited or it may in turn absorb and replace all the cells of the endosperm and so come to occupy the space within the integument, as in the bean, pea, etc. (Figs.

82, *A*; 83, *D*). Various other arrangements may result; as for example, there may be remains of the nucellus and endosperm with the embryo (Fig. 82, *B*). The integument also undergoes various changes during these growths, often becoming hard or papery or provided with hairs, hooks, spines, and smooth or pitted. Compare the bean, catalpa, cotton, milkweed, etc. Attention must also be directed to the growths that occur outside of the seed that are induced by fertilization and the germination of the gametospore. Very frequently the pistil becomes tough and hard and invests the seed like a second integument, as in wheat, corn, acorn, etc.; again it forms a sac about the seed as in the sedges and buttercups; or it may enlarge, forming the pods of peas and beans or the capsules of lilies. In other cases it becomes fleshy throughout as in the tomato and currant, or only the outer walls are fleshy while the inner part forms the pit or stone as in the peach and plum. Frequently wings and hooks grow out from the pistil as in the maple, *Ailanthus,* etc. Even the end of the stem, the receptacle, which bears the various organs of the flower, may be induced to grow, as in the apple, where it surrounds the pistils (the core) with a fleshy coat, or the receptacle may enlarge, as in the strawberry, forming the fleshy part of the fruit. In other cases outgrowths below the flower develop, forming the burr of the chestnut, clotbur, beechnut, and the cup of the acorn. The entire growth that is associated with the formation of the seed is termed the fruit and we see that it may include a variety of modified organs. The term seed, however, refers only to the modified ovule and its embryo. Later in the work it will be noted that these growths are often devices to bring about a distribution of the seed or to protect it during its dormant period.

55. The Seed and its Growth.—Spores of all kinds germinate and produce new plants. Seeds do not germinate. They simply continue the growth of the embryo or plant which has been temporarily stopped. The falsely called germination of the seed is but the awakening or renewal of the growth of this plant. The seed is essentially a young plant that is supplied with a certain

amount of food and that has temporarily stopped growing. The seed may be looked upon as one of the many adaptations found among plants to ensure their perpetuation. From a seed a single plant is developed that may produce many, often thousands of seeds, *i. e.*, plantlets. We have already noted that these may be widely distributed and so spread and multiply the number of plants. Some seeds are so fine that they float away as a dust, as in the orchids. Other seeds and fruits are provided with hairs and wings that serve to buoy them up or promote their transport by winds, as in the case of pine seeds, milk weed, willow, and the fruits of maple, dandelion, etc. In other cases spines and hooks or mucilage glands are developed that attach them to any passing animal and so bring about a considerable distribution. Note the common devices of this nature, as in the burdock, stick tight, beggar lice, agrimony or the glands on the nightshade (*Circaea*) and twin flower. So also seeds are scattered by the spring of elastic stems and explosive fruits, as in many lilies, witch hazel, violet, touch-me-not, etc. But especially may seeds be looked upon as an adaptation to tide the plant over seasons unfavorable for growth. In this respect they may be compared to buds, which they also resemble in their renewal of growth when conditions are again favorable. There are some noteworthy exceptions to this statement, as in the mangrove, certain oaks, and many grasses. The seeds of the former plant begin their growth while still attached to the tree and a similar growth has also been observed in the acorns of the white oak. Grain often sprouts while in the sheaves and the seeds from green tomatoes produce earlier and larger fruits. However the majority of seeds and fruits, as well as buds, can not renew their growth immediately after their formation. A longer or shorter resting or dormant period is required during which time chemical changes occur that render the foods soluble and therefore available to the young plant or embryo. For example, in many seeds and buds the storage foods are in the form of starch. This food is insoluble and can not be used by the plant until it has been acted upon by an enzyme which changes it into a soluble and readily diffusible sugar. These enzymes are

slowly formed during the resting period and a renewal of growth is not possible until they have begun the transformation of the insoluble foods into soluble forms. The changes thus effected are so considerable that growing tubers, root stocks and seeds often have a sweet taste owing to the considerable amount of sugar present. So the seed may be looked upon as a variation in the growth of the plant that enables it to multiply and distribute its kind and to meet the various problems in connection with its existence, such as drought, cold, and sometimes even fires.

56. Conditions Necessary for Growth.—The external conditions necessary for starting the growth of the embryo are a suitable temperature, moisture, and the oxygen of the atmosphere. Some seeds will germinate at near the freezing point, while others require a temperature between 10° and 17° C. It is interesting to note that the rising temperature as the season advances from early spring brings out a regular succession in the awakening of various kinds of seeds. A suitable amount of moisture must also be supplied to the seed to assist in the solution and diffusion of the foods to the embryo. This is very necessary since the seed contains very little water. In fact the seed owes its vitality and power to resist long drought and severe cold to its dry condition. Seeds soaked in water cannot endure such extremes as in a dry state. Excess of water, however, is unfavorable to the renewal of growth, because it excludes the oxygen of the air. This is the cause of the failure to grow and the decay of the seeds planted in the spring when the season is rainy. The air spaces in the soil become filled with water and as a consequence oxygen can not gain access to the seed and start those chemical changes that not only supply some of the foods to the embryo but also provide it with the energy for growth. An examination of a few seeds will show some of the more important structural features that adapt them to the conditions that they have to meet.

In the case of the bean the integument is tough and would appear to offer a rather effectual barrier to the entrance of water and air. You have noted that a minute opening, the micro-

pyle, exists in the integument. This opening is seen near the scar or hilum that marks the point where the seed was attached by a minute stem to the walls of the ovary (Fig. 83, *B*). This



FIG. 83. Fruit and seed of the bean: *A*, mature fruit or pod: *s*, style; *c*, remains of calyx; *o*, ovary. At left the pod has been opened, showing four seeds attached to the walls of the ovary by funiculus. *B*, two views of a seed—*h*, hilum or scar marking attachment of funiculus to seed; *m*, micropyle. *C*, wrinkling of seed coat caused by the entrance of water through the micropyle and hilum. *D*, embryo of seed after removal of integument—*c*, cotyledons. At left one of the cotyledons has been removed to show the plumule, or epicotyl, *ep*, and the hypocotyl, *hy*. *E*, half of the membranous integument.

opening permits the entrance of the air and water as does the looser structure of the hilum. This can easily be demonstrated by placing beans in water when the spread of the water from the region of the micropyle will be indicated by the swelling and wrinkling of the integument (Fig. 83, *C*). After a time the entire integument becomes quite soft and water is readily drawn through it owing to the fact that the storage foods in the embryo are in close contact with it and draw the water through by osmosis. Carefully removing the integument from a soaked bean we note that the embryo is well developed, consisting of two fleshy cotyledons and a short stem (Fig. 83, *D, E*). The epicotyl is well developed forming a plumule bearing two leaves and the short hypocotyl and root are enclosed in a sheath formed from the integument. Note that the hypocotyl and root are situated near the micropyle. How does this arrangement work

to the advantage of the embryo? Allow the seed to germinate and it will be noted that the absorption of water first causes the embryo to expand and rupture the integument. This is immediately followed by the growth and elongation of the lower part of the hypocotyl which results in the pushing out of the root (Fig. 84, *A*). Thus it happens that the very part of the embryo



FIG. 84. Renewal of growth of the bean seed: *A*, basal region of hypocotyl elongating and pushing the root into the soil. *B*, upper region of the hypocotyl elongating, lifting the cotyledons and epicotyl above the soil. *C*, hypocotyl erect and epicotyl expanding, forming the first normal leaves.

that shows the first signs of growth is in close proximity to the water as it enters the seed and doubtless the sheath also assists in retaining water in this region. We are already familiar with the properties of the root which direct its growth in such a way that no matter how the seed may be placed the root will ultimately grow into the soil and toward water and the soil foods, page 54. It is noteworthy that many seeds are so fashioned that they naturally assume such a position on the ground as to bring the root as it emerges into direct contact with the soil. Such an arrangement is equally advantageous for the absorption of water. How many seeds can you find that will naturally lie upon the soil in such a way as to cause the root on emerg-

ing to be directed away from the earth? Thus at the very start the growth of the seed is of such a character that the young plant is anchored to the soil by the root and brought into proper relation with the water and other materials necessary for the construction of foods. As soon as this work is well under way the upper end of the hypocotyl begins to elongate, pushing up into the air cotyledons and plumule (Fig. 84, *B*). Here we see that the upper end of the hypocotyl reacts to gravity like a stem and consequently grows away from the earth. This growth is attended with a curving and constant nutation of the hypocotyl which enables it to work its way through the ground to better advantage than would be the case if it were pushed passively up. Darwin compares this work of the seedling in emerging from the soil to the struggles of a man who is prostrate and attempting to arise with a heavy load on his back. The ability of the hypocotyl to overcome obstacles in its way and crowd through small openings is due to the power generated in the enlarging cells. In this way a force is often created that may exceed 80 pounds to the square inch, over five times the atmospheric pressure. Many seeds only elongate the epicotyl, leaving the cotyledons in the ground, as in the pea, acorns, walnuts, etc. This behavior is doubtless associated with the storage of food. In some seeds the cotyledons serve only as storage organs and in such cases the position of advantage is in the soil where they are better protected and in contact with moisture which will assist in the solution and diffusion of the foods. In many cases the cotyledons perform a dual function, first as storage organs and later expanding and developing into the first green leaves of the plant, as in the squash and many of our common plants. In any of these cases it is evident that the young plant requires a certain amount of food to enable it to get established in the soil. It is not until considerable growth has taken place that it develops its first green leaves and so begins to be self supporting. In the case of the bean and in many other seeds and fruits this need is amply met by the abundance of food that is stored in the fleshy cotyledons (Fig. 84, *C*). These organs remain attached to the plant even after it is well established, and it

is only after the food has been withdrawn that the shriveled cotyledons drop off. As the young plant becomes established the plumule unfolds like a bud and develops the stem and leaves and finally the flowers and fruit with which you are familiar.

The castor bean shows quite a variation from the structure of the bean. The hilum and micropyle are more or less concealed by a fleshy outgrowth, caruncle (Fig. 85, *A*). Remov-



FIG. 85. Seed and seedling of the castor bean: *A*, the seed with fleshy outgrowth, the caruncle, *c*, at base. *B*, section of seed. At right showing the two cotyledons, *c*, and short hypocotyl in center of the endosperm, *en*. At left the seed is split parallel to the surface of the delicate white cotyledon. *C*, early stage in the growth of the seedling. The hard integument is being thrown off but the cotyledons are still entirely buried in the endosperm. *D*, later stage. The cotyledons, *c*, still absorbing food from the endosperm, *en*, are beginning to appear and to develop chlorophyll. *E*, the cotyledons and epicotyl are expanding, the endosperm being entirely absorbed.

ing the hard integument we find within a white, oily mass of cells which are attached to the horn-like coat only at the hilum. Cutting across this mass of cells it will be seen that two thin leaves are imbedded in the center. Taking two other seeds and cutting them longitudinally, one at right angles and the other

parallel to the leaves we see that the embryo consists of a very small stem and two delicate leaves imbedded in an oily tissue, the endosperm (Fig. 85, *B*). In the bean the seed consisted of an embryo and integument. In the castor bean, the endosperm forms the bulk of the seed, since it has been only partially consumed by the embryo. It would naturally be inferred that the hard coat would prevent the access of water and gases. Observe, however, that the caruncle readily absorbs water and that the embryo and endosperm are in direct contact with this region of the seed. Many devices will be observed among seeds serving to convey water to the growing parts. Note the mucilaginous coats on flax, quince, many mustard seeds, etc., or the spongy rinds of the fruits of the walnut, cocoanut, etc. It is well worth your time to submerge in water colored a deep red with eosin widely different seeds and examine them from time to time, noting the period required for the penetration of water and the lines along which the water enters and is distributed. It will be found that many seeds and fruits exclude water and oxygen very effectively for one or more years with the result that the renewal of growth is delayed until the decay of the parts permits their entrance. The breaking of the integument would appear as a serious obstacle to the growth of the embryo. However, the power generated by the cells owing to their expansion through the absorption of water is quite sufficient to rupture the seed coat. The early growth of the embryo of the castor bean is similar to that of the bean. The elongation of the lower end of the hypocotyl pushes the root into the soil, while the upper end of the hypocotyl, arching and twisting as it elongates, works the cotyledons and endosperm through the soil into the air (Fig. 85, *C*). In this seed the food is without the cotyledons. Consequently, although the cotyledons gradually enlarge and become green, they remain in contact with the endosperm until all its food is absorbed and but a papery skin remains (Fig. 85, *D*).

A grain of corn will illustrate another modification of the seed. In this case the pistil remains in such close contact with the seed that it appears to form an additional integument. Such

a fruit is called a grain and may be compared to a bean pod with a single seed. We can still see upon the grain (Fig. 86, *A*) at *s* the position of the style while at *p* appears the remains of the stem that fastened the sporophyll to the cob. The long threads, often called the silk, that projects from the ears of corn are greatly elongated stigmas for catching the microspores. While the outer parts of the grain of corn are very hard if it is treated with the eosin solution referred to above it will be seen that at first the water enters by way of the little stalk just



FIG. 86. Grain and seedling of corn: *A,* two views of a grain—*s,* point of attachment of style; *p,* stalk attaching the grain to cob; *emb,* location of embryo. *B,* longitudinal section of a grain. The embryo consists of a plumule, *pl,* root, *r,* and scutellum, *sc. en,* endosperm. *C,* cross-section of a grain showing endosperm, *en,* surrounding the embryo, *emb. D,* diagram of the outer cells of the scutellum, showing the elongated cells that bring about a solution and absorption of the foods stored in the endosperm. *E–G,* stages in the growth of the seedling.

as did the food that was supplied to the growing grain. In this way water is supplied directly to the stem of the embryo which lies near the stalk as shown in Fig. 86, *B*. If two seeds are soaked until soft and then sectioned so as to cut one across and the other longitudinally through the embryo it will be seen

that the endosperm forms the larger part of the grain and that the embryo is relatively small, consisting of a root, plumule, of several closely wound leaves, and a shield like organ, the scutellum (Fig. 86, *B, C*). The entire embryo may easily be removed from the endosperm with a knife if the seed is thoroughly soaked. Unlike the two preceding examples the corn is a monocotyledon and many regard the scutellum as a modified cotyledon. This organ forms the enzymes which put into soluble form the food in the endosperm and it also functions as an absorbing organ of the embryo (Fig. 86, *D*). Similar devices for absorbing the storage foods and transferring them to the embryo are seen in the wheat, onion, date, and cocoanut. The early growth of the embryo goes on as in the preceding examples. The root first emerges and anchors the grain in the soil. The plumule of closely rolled leaves forms a sharp bodkin that reaches up through the soil with ease. This modification of the plumule would appear to be of a decided advantage and you will notice that this method of emerging from the soil is followed by a great many plants with underground stems; see the large leaves of the skunk cabbage, may apple, Solomon's seal, etc. The rest of the grain remains in the ground as in the case of the pea.

57. The Two Phases in the Life of a Plant.—As has been stated the plumule unfolds in all these cases like a bud and gradually develops into the mature plant. This may be effected in one year or several years may be required to complete the work. At some time in this growth flowers will appear bearing sporophylls and spores and thus we arrive again at the starting point in the life history of a plant. There are two distinct phases or generations in the life of a plant. The formation of the megaspores and microspores marks the beginning of the sexual generation or gametophyte which ends when the male and female gametes have been developed. The formation of the gametospore marks the beginning of the asexual generation or sporophyte which ends in the sporophylls when those cells, called the spore mother cells, appear which produce the micro- and megaspores. These two generations are also sharply distinguished from each other by the number of chromosomes, page 49, which

their cells contain. The cells of the gametophyte generation contain only one-half the number of chromosomes found in the cells of the sporophyte generation. If the nuclei of male and female gametes contain twelve chromosomes (the number varies in different plants) the gametospore which marks the beginnings of the asexual generation will contain twenty-four chromosomes. This number will be found in all the cells of the embryo and continue to characterize the subsequent development of the sporophyte. When, however, the spore mother cells appear in the anthers and ovules it is to be noted that these cells in dividing to form the micro- and mega-spores reduce the number of chromosomes by one-half so that the nuclei of these spores and all cells derived from them contain but twelve chromosomes. For this reason the division of the spore mother cell is referred to as the reducing division. The significance of this difference in the number of chromosomes is not known but it furnishes a basis for the sharp separation of the two generations. The evidence at present indicates that the chromosomes of the male and female gametes retain their identity until the spore mother cells are formed. When these cells divide there is a fusion of the male and female chromatic substance, attended with a reduction division of the number of the chromosomes. If this be the case it will be necessary to alter our conception of fertilization for apparently the real fusion of the nuclear substance of the gametes is effected in the formation of the spores rather than in the union of the gametes and the formation of the gametospore.

A sexual and an asexual phase or generations characterizes the life history of nearly all the higher plants and the succession of these two phases is called the alternation of generation. In the case of the bean we have as the most important features of the sporophyte or asexual generation, (1) the gametospore, (2) the seed, and (3) the flowering bean plant. The microscopic plants that comprise the gametophyte or sexual generation are characterized by (1) the micro- and mega-spores which develop respectively into (2) a tubular growth of three cells and (3) a sac-like growth of seven cells, each of which contains

one or more (4) sexual cells or gametes. These two phases are called respectively the sporophyte or asexual generation, and the gametophyte or sexual generation, since in the first phase mother cells are developed which form spores, *i. e.,* this is the spore forming or asexual generation and in the second phase gametes or sexual cells are produced.

58. Significance of Fertilization.—Let us now stop to consider the meaning of the complicated process that we have termed fertilization. No satisfying explanation has been offered. Some see in these changes only a process of nutrition and chemical stimulation. The gametes are lacking in certain materials that are essential to their further growth. According to this view the male gamete is the complement of the female and by the union of the two all the substances are supplied that are necessary for growth.

It has also been suggested that the growth culminating in fertilization is attended with the removal of impurities from the sexual cells. Every cell has its growth, maturity, and senescence when it loses its power for further growth and division. The reduction division and the subsequent formation of the gametes results in the removal of substances that prevent their continued activity and as a consequence they are restored to a youthful condition which appears in the growth of the gametospore.

Much attention has been directed in recent years to the process of fertilization as a method of controlling the character of of offspring. You have already noted, page 49, that granular bodies or chromosomes are constant features of every nucleus. It has been supposed that these complex bodies direct the vital activities of the cell and determine the type of plant that shall be developed. In other words the chromosomes contain the hereditary substances that cause each plant to develop in a certain definite way or to follow a certain pattern in its growth. The chromosomes derived from the parent plants cause the offspring to resemble them in growth. The micro- and mega-spores are formed in the sporophylls, and these spores must therefore contain the same kind of chromosomes or hereditary substances as the plants bearing the sporophylls, *i. e.,* the parent plant, since

they have been derived directly from the parent plant by cell division. The gametes derived from the germinating spores must also contain the same kind of hereditary substances as the parent plants. In fertilization we have the fusion of the gametes. This means that the hereditary substances in the gametospore will be of a dual character since it has been derived from the male and female gametes (Fig. 80). Consequently there is now united in one body, the gametospore, all the possibilities of growth that each gamete inherited from the parent plant. The transmission of hereditary characters may also be stated as follows. Every feature of a plant or animal which we recognize in distinguishing them is due to hereditary substances derived from the parents. These substances, possibly lodged in the chromosomes, cause definite traits (called unit characters) to appear in the offspring. For example, the form of the leaf, the color of the flower, the fragrance of the fruit, etc., are unit characters and each is due to a definite hereditary substance derived from the parents. These unit characters are transmitted by the parent to the offspring. They may fail to appear, *i. e.,* lie dormant, but when active they always cause the same trait or character to appear in the succeeding generation. Offspring derived from parents that differ in possessing one or more unit characters not common to both parents are called hybrids. Hybrids differ from either parent because they have all or only a portion of the unit characters of both parents. Parents of the same kind, *i. e.,* of the same species, have identically similar unit characters and the offspring of such plants and animals must resemble the parents because they only have unit characters common to both parents. For this reason a species is often defined as a kind of plant or animal that is capable of producing offspring like itself.

At present it is impossible to state the nature and extent of the variations that may result in the offspring from the union of the hereditary substances of the parents. The theory has been advanced that the purpose of fertilization is to lessen the pronounced parental traits by blending such characters in the offspring. In a word fertilization has a tendency to diminish the accentuated traits of the parents and to lessen the amount of

variation in the offspring so that they conform to a more constant and uniform type. On the other hand many consider that fertilization is designed to bring about variation in the offspring. So numerous, however, are the traits or unit characters of each parent that it is often impossible at present for the holders of this view to interpret the nature of the variations that sometimes appear in the offspring. In some cases there appears to be a complete mingling of the parental characters. When, for example, the parents vary in color or some other character, the offspring may show a blending of these two colors as when two pigments are mixed. More commonly certain characters of one parent appear to be stronger in controlling the development of the offspring and suppress certain characters of the other parent. As a result the offspring resemble one parent more than the other. Why this occurs can not be explained. We can only say that the hereditary substance of one parent is perhaps more potent and has more influence in the development of the offspring. These stronger features that appear in the offspring are called the dominant characters. It usually happens in the second and succeeding generations that some of the offspring will show certain characters that were suppressed by the dominant characters in the first generation. These weaker traits are called recessive characters because they reappear in the later generations. The number of offspring with dominant characters often stand in definite ratio to those with recessive characters. For example taking a single unit character of two plants, as the black and yellow color of their seeds, Shull has shown when two such plants are crossed that only black seeds may be produced in the first generation. This color is therefore called a dominant character because it suppresses the yellow. If these seeds are grown and the resulting flowers are crossed among themselves it will be found that the seeds produced in the second generation vary in color. Approximately one-fourth are pure black, two fourths a mixture of black and yellow, with the black dominant, *i. e.*, masking the yellow, and one fourth pure yellow. The pure yellow and the pure black seeds continue to breed respectively yellow and black seeds in all subsequent generations and it

is evident that they represent pure stock in which the black and yellow unit characters have been entirely separated. The seeds that were a mixture of black and yellow produce when crossed among themselves in the third generation the same ratio of dominants, recessive and mixtures as appeared in the second generations, namely, one-fourth pure black, two-fourths black and yellow with the black dominant, and one fourth pure yellow. So we see that each generation after the first is characterized by the appearance of a rather definite proportion of offspring with pure dominant or pure recessive characters, and with mixed characters. This ratio of variation is known as Mendel's Law. Plant and animal breeders rely upon these separations in producing new forms and by careful breeding they are able at will to add to or remove from the offspring one or more of the parental unit characters. In some cases this is possible with such certainty that horticulturists have advertised new forms, describing their characters and qualities, before they have been bred. How applicable Mendel's law is to all cases of crossing has not been determined. In the case cited above where only two unit characters were involved there can be but three combinations of them, *i. e.*, pure black, pure white and a mixture of black and white. It is manifest where many unit characters appear that there will result a re-combination of these characters in the offspring that is complicated and difficult to interpret. When however the number of unit characters that appear in any plant or animal are definitely known is is probable that the variation of the offspring can be definitely forecast.

It must be borne in mind that the variations referred to above do not result in the formation of new characters but in the recombination of characters already existing in the parents. As to the actual origin of new characters there is no generally accepted explanation. On the other hand there is a widely prevalent view that insignificant departures may appear in any form owing to some circumstance of environment and that these departures may gradually become intensified so that after a long lapse of time they are sufficiently pronounced as to warrant the recognition of the forms possessing these characters as distinct

species. This view is purely theoretical and is not supported by any demonstrable evidence.

As opposed to this theory deVries has called attention to the fact that entirely new forms may appear at any time in the offspring of the parents. These new forms are called mutants because of their sudden appearance. During his cultivation of Oenotheras, evening primroses, in which only plants of the same kind were crossed, he found after fifteen years of cultivation in which time over 80,000 plants had been under observation that fifteen new forms had suddenly appeared at one time or another. These mutants are distinguished by new characters not possessed by the parents and furthermore they transmit these characters to their offspring. This theory of mutation holds that the unit characters which distinguish the individuals of a kind or species may be influenced by external forces with the result that one or more of them may be blotted out or changed in its nature. The offspring therefore will be characterized by the absence of certain parental characters or will appear with new ones not possessed by the parents. To what extent this sudden appearance of new characters has been a factor in establishing new forms of plants and animals remains to be seen. Illustrations in addition to those previously noted, will repeatedly occur later in the work, showing that plant life is profoundly influenced by the stimuli of its environment. Gager has recently shown that marked variations may be induced in the offspring of *Oenothera* by subjecting the parents to the emanations of radium and MacDougal reports like results from injecting solutions into the ovaries. It seems altogether probable that the hereditary substances may be similarly affected by the varied stimuli which the plant is constantly receiving in nature.

Later in the work it will be seen that nearly all plants have devices of various kinds that assist in the transfer of the microspores of one plant to the stigmas of other plants. In this way the sexual generations are brought together that have been derived from plants that may differ more or less from one another. All these differences in so far as they are due to the hereditary substances must be transmitted to the gametospore, consequently

it will develop into a plant that will show various associations of the parental characters. It is possible that many kinds of plants which we call species have arisen in this way. Mutations may also have produced species and also lead to more pronounced changes in the offspring which we recognize as genera.

This brief consideration will give us an idea of the nature of variation and it also helps us to understand the significance of the elaborate processes leading up to fertilization. The variations and departures of the offspring from the parental type are of great importance. In this way new characters arise that enable the offspring to live under conditions that the parents can not endure, consequently the offspring can occupy territory that would not support the parents and as a result they do not come into competition with them. Variations have gone on from one cause or another during the past ages and thus plants have become adapted to practically all the different conditions of soil and climate on the face of the earth. It must also be borne in mind that these variations may adapt a wide variety of plants to one and the same condition. Consequently in every locality we find plants living together and fiercely competing for their existence. Here again it is the variation of each kind of plant that gives it a slight advantage and enables it to compete successfully with its associates or supplant them. It is instructive to watch the changes in vegetation that occur in the cultivated field that is allowed to remain fallow for several years. Each succeeding year will mark the crowding out of certain forms by plants that possess variations that better adapt them to the locality. These changes in the vegetation will go on until the forms are introduced whose special variations adapt them to the conditions of the field and to the competition of the other plants. This vegetation of the field is now fixed owing to the survival of the fit. In this way the distribution and association of the vegetation of the earth has been brought about.

PART II

DEVELOPMENT OF PLANTS

CHAPTER V

CLASSIFICATION OF PLANTS

59. The Method of Classifying Plants.—We have now arrived at an understanding of the nature of the plant and the work which it performs. In this study the most complex forms have been largely considered. There are, however, a great variety of very simple forms ranging from single-celled plants that often form the green coatings on trees up to the complex types. How have these different forms come about and what relation do they sustain to one another? We will also be interested to compare the life histories of these different forms and their modes of life. In approaching this subject it is first necessary to gain a knowledge of the system employed in grouping or classifying plants. Obviously many plants are closely related as the red, white, scarlet oaks. Such a group of closely related plants is called a genus (plural, genera) and the different kinds of individuals which compose the group are known as species. The red oak is one of the species of the oak genus. In scientific work the Latin or Greek names are used to designate the genera and species. In the case of the oak the genus is known as *Quercus,* an old Latin name for oak, and the red oak species as *rubra,* meaning red. Both the name of the species and the genus are employed in naming a plant and consequently the scientific name of the red oak is *Quercus rubra.* So in other cases we can say white oak or *Quercus alba,* scarlet oak or *Quercus coccinea,* etc. The oak has several characters in common with the chestnut and beech nut genera, as for instance the fruit in each case is asso-

ciated with outgrowths that appear as the burr in the chestnut and beech and as a cup in the oak. On account of these and other common characteristics these genera are supposed to be related and are therefore grouped together in one family. A family is composed of allied or related genera. The name of the family is derived from some characteristic genus in the group. In this case it happens to be the beech or *Fagus*. To distinguish a family from a genus the termination *aceae* is added to the base of the generic word, in this example making the family name Fagaceae. So we have the three allied genera, oak, beech, and chestnut forming the beech family or Fagaceae. In the same way it will be found that families are related and joined together into a still larger group known as the order. The flowers of the birch, alder, water beech, and hazel genera are very similar and consequently form a family known as the birch family or Betulaceae, Betula being the name of the birch genus. It will also be noticed that there is a similarity between the flowers of the Betulaceae and Fagaceae. This relationship is expressed by placing them in the same order known as the beech or Fagales, the termination *ales* being employed to distinguish the order just as *aceae* characterized the family. So also orders are related and grouped into classes. For example, the Fagales and many other orders are characterized by having seeds with two cotyledons. Several other orders have seeds with but one cotyledon. This relationship is expressed by grouping the orders into two classes, Dicotyledones and Monocotyledones. In both of these classes the seeds are inclosed within the pistil, but in the cone-bearing trees the seeds are exposed on a flat scale-like organ. All seeds producing plants belong to one or the other of these two groups. So we have two subdivisions, the Angiospermae, meaning seeds inclosed, and the Gymnospermae, meaning naked seeds. If now we stop to consider still larger groups than subdivisions, it will be noted that many plants do not produce seeds, as in the case of the ferns and mosses, etc. So a final grouping of plants into divisions or subkingdoms may be made. All seed-bearing plants form the division Spermatophyta, meaning seed plants; the ferns comprise the division Pteridophyta, meaning fern plants,

etc. The termination of *phyta,* from phyton a plant, distinguishes the division from the other groups. To repeat, the division is composed of subdivisions which in turn are made up of classes. Frequently the division contains only classes. Classes consist of related orders. The orders are divided into families which include allied genera and these latter groups comprise closely related individuals or species. The vegetation of the earth is separable into four widely differing divisions: I., The Thallophyta, including among others the fungi and algae; II., The Bryophyta or moss plants; III., Pteridophyta or fern plants; IV., The Spermatophyta or seed plants. The character and relationship of these groups will be considered in the following pages.

CHAPTER VI

DIVISION I. THALLOPHYTA

60. Classification of the Thallophyta.—The Thallophyta comprise a multitude of plants that include the most primitive and simple forms of vegetation upon the earth. They range in size from single microscopic cells to forms that are comparable in bulk to some of our shrubs and trees. They are all of simple structure and do not possess roots, stems and leaves in the sense of the seed plants. Such a type of plant body is called a thallus and it assumes a variety of forms. The absorption and manufacture of foods is carried on, as a rule, by any and all of the cells and this is equally true of the reproductive process. It would be impossible, however, to give in a short space a general idea of the nature and character of this division which includes many groups of widely different forms. For the purpose of study the Thallophyta may be divided into five subdivisions:—(1) Myxomycetes or Slime Moulds; (2) Schizophyta or Bacteria and Blue green Algae; (3) Diatomeae or Diatoms; (4) Euphyceae or Algae; (5) Eumycetes or Fungi.

Subdivision 1. Myxomycetes or Slime Moulds

61. The Life History of a Slime Mould.—In one stage of their life the slime moulds have a motility and mode of feeding suggestive of some of the lower animals, while on the other hand the final stage of their existence is more suggestive of the fungi. For this reason it has often been suggested that they are intermediate between plant and animal life. These plants are widely distributed over the earth and may frequently be found on decaying logs and rotting twigs and leaves in forests. A common form, resembling a miniature, brownish puffball, is often seen on stumps and fallen logs (Fig. 87). Other kinds are illustrated in Fig. 88. They range in size from scarcely a pin head to nearly

a foot in diameter and from spherical to cylindrical and cake-like masses. Not infrequently they are of great beauty owing to their coloration and lace-like structures. These small sacs or sporangia (sing. sporangium) as they are commonly called, represent but one stage in the life of the slime moulds. If we begin

Fig. 87.



Fig. 88.

Fig. 87. The sporangial stage of two common slime moulds: *A, Lycogala. B, Arcyria. C,* sporangia rupturing, hair-like structures (the capillitium) and spores protruding. *D,* sporangia emptied.

Fig. 88. Open types of sporangia: *A, Stemonitis,* at left a single sporangium enlarged, showing net-like structure formed by capillitium radiating from the central stalk of the sporangium. *B, Cribraria.*

with an examination of the dust that floats away from these sacs every time they are tapped, the life history will be found to be about as follows: Under the microscope the particles of dust are seen to be minute cells or spores (Fig. 89, *A*) each provided with a cell wall and containing a nucleus and protoplasm. A spore is a cell that is capable of germinating under favorable conditions and producing a plant. In this case it is advisable when germinating the spores to place them in a nutrient solu-

tion made by boiling in water bits of wood upon which the sporangia were growing. Germination will begin in about one day. As the spore wall ruptures the contents escapes as a naked bit of protoplasm (Fig. 89, *B*) which possesses a singular power of motion. This motility is due either to the slow creeping or streaming motion of the soft plastic body or more commonly the body may possess a delicate thread or cilium (plu. cilia) (Fig. 89, *C*). These ciliated bodies are called zoöspores. The cilium is a highly sensitive organ which by rapid and rhythmical contraction and expansion beats the water and thus drives the body along. While in this condition the zoöspores are rapidly multiplied owing to the repeated splitting of the little bodies into two similar parts. Finally this greatly increased number of individuals begins to come together in small groups (Fig. 89, *D*) which in turn merge and so form large slimy masses termed plasmodia (sing. plasmodium) which are quite destitute of all walls and consist of a great number of nuclei surrounded by a watery cytoplasm (Fig. 89, *E*). You will often find this plasmodial stage of the slime mould on the under side of rotting limbs or bits of wood in the forest as a sticky mass resembling in consistency the white of an egg. The mucilaginous character of the plasmodium accounts for the name slime mould popularly applied to these plants, also for the term myxomycetes, from myxos, slime, and myces, mould or fungus.

The plasmodium possesses a sensitiveness or capability of responding to external stimuli to a degree that is remarkable when we consider the extreme simplicity of these plants. For example it avoids too strong a light and slowly moves toward moisture and food. This accounts for your failure ordinarily to notice it since it seeks the darkness and food in decaying logs or the under surface of sticks and leaves. The motion is brought about by little arm-like branches that flow from the jelly-like plasmodium and finally the entire plasmodium will slowly follow along these lines with a complicated streaming movement (Fig. 89, *E*). If bits of wood containing some of the plasmodium are placed in a damp chamber in the dark after a time the plasmodium will be seen creeping up the sides of the

moist dish in a complex skein-like mass or if a glass slide, down which water is slowly allowed to trickle, is brought in contact with the plasmodium it may be observed creeping up the slide toward the source of the water supply. The slime moulds do not contain chlorophyll and are therefore dependent upon organic food. Plants that feed in this way are called saprophytes. Since the plasmodium is not surrounded by a cell wall feeding becomes a simple matter. The jelly like mass engulfs the decay-



Fig. 89.



Fig. 90.

Fig. 89. Mobile phase in life of slime mould: *A*, group of spores. *B*, germination of a spore. *C*, two forms of zoöspores greatly enlarged. The right-hand one mobile owing to a streaming movement of its protoplasm and the left-hand one moves about owing to the rapid movements of the cilium, *c*. *D*, association of the zoöspores preliminary to the formation of a plasmodium. *E*, plasmodium. The arrow indicates the direction of the streaming movement.

Fig. 90. Character of the capillitium: *A*, in *Arcyria*. *B*, in *Trichia*, portion of thread enlarged on the left hand.

ing particles or other foods and after digesting the nourishing portions casts out the worthless part. As the plasmodium approaches the final stage of its life, its nature appears to change completely for now it avoids moisture and seeks the light. It creeps to the surface of the wood or leaves in which it has been

growing and forms the characteristic bodies seen in Figs. 87, 88. These sporangia are formed by the outer part of the plasmodium hardening into a wall, while each of the inclosed nuclei, which have greatly increased in number by division, becomes surrounded by a wall, thus forming the spores. Usually a portion of the substance of the young sporangium is transformed into simple or branching threads or tubes, collectively called the capillitium (Fig. 90). In several of the genera the wall of the sporangium is formed of a net work of the capillitium (Fig. 88). These threads are hygroscopic and their constant motion assists in stirring up the spores and exposing them gradually to the wind as soon as the wall of the sporangium ruptures. A somewhat simpler type of slime mould lives as a parasite in the roots of turnip, cabbage and cauliflower producing a destructive disease known as club-root.

There are several groups of low types of plant and animal life that are suggestive of relationship with the slime moulds. The most important among these, the myxobacteriales, have been made known by Taxter. They are minute plants, very suggestive of the next group, the bacteria, but associated in definite structures that resemble the sporangia of the slime moulds and also of certain fungi. On the other hand, certain aquatic forms, as *Protomyxa*, with a life history very similar to that of the slime moulds, intergrade almost perfectly towards simple animal types, as the protozoans, of which the common amoeba is an example.

Thus we see that the life history of these plants is a very simple one. By the formation of sporangia numerous dust-like spores are produced that are capable of germinating and forming zoöspores that increase rapidly by division. The aggregation of the zoöspores results in the formation of the plasmodium which after a time completes the life history by creeping upon suitable dry objects and forming sporangia. The mingling of the zoöspores in the formation of the plasmodium is not a sexual process, since the nuclei do not fuse. As far as known the entire life history is purely an asexual existence. It is possible, however, that the purpose of reproduction may be effected among

simple plants in some other way than by the fusion of a male and female gamete. The structure and life history of these plants is so simple that it is possible that they may have been derived from forms allied to the first forms of life upon the earth. The common occurrence among simple animals and plants of motile stages or zoöspores in their life history certainly suggests that possibly such forms may represent the first appearance of living matter.

Subdivision 2. Schizophyta or Bacteria and Blue Green Algae.

As in the preceding group these forms are extremely simple and possess some characters suggestive of the lower forms of animal life. They have received the name of Schizophyta, meaning splitting plants, owing to their common method of reproduction by division of the cell into two equal parts. There are two important classes of Schizophyta: A, Bacteria; B, The Blue Green Algae.

CLASS A. BACTERIA

62. The Structure and Nature of Bacteria.—Bacteria are commonly known by such vague terms as microbes and germs. They are, however, unicellular plants that are of almost universal distribution though more abundant about dwellings and less common in cold countries, at high altitudes and on the sea. They include the smallest and simplest forms of plants, ranging from scarcely 1/50,000 in. to 1/10,000 in. in diameter (Fig. 91). Such forms would have many times more room in a drop of water than a whale would find in New York harbor. So minute and simple in structure are the bacteria that the real nature of the plant body is somewhat a matter of dispute. The plants are unicellular and surrounded by a delicate thin wall which incloses a colorless and slightly granular protoplasm (Fig. 91, *A*, 1). There is no nucleus comparable to that of the higher plants, although indications of it are seen in a few scattered chromatin grains. The cell contents is very simple and totally lacking in plastids and other differentiations with which you are familiar. The slimy appearance of bacteria noticeable where they grow

together in colonies is due to the mucilaginous excretion from their bodies which is often brightly colored. Bacteria range from globular to rod-like and curved forms (Fig. 91). Many



FIG. 91. Forms of bacteria: *A, Bacillus subtilis,* a form common in hay infusions. 1, motile state; 2, cells with spores; 3, slimy mass of bacteria, the zoöglea condition, that appears on the surface of infusions, cooked vegetables, etc. *B, Spirillum. C,* a coccus form that appears in pus. *D,* mobile and spore stage of lock-jaw bacillus.

are motile by means of cilia, as in the case of the zoöspores of the Myxomycetes, which project singly or in tufts from the ends of the cells or in varying numbers from all sides.

(*a*) *Reproduction of Bacteria.*—This is a simpler process than in the slime moulds and much more rapidly effected when the conditions are favorable for growth. This process consists of forming a partition through the middle of a cell when the two daughter cells thus formed separate at once or they may remain attached, forming ultimately a chain of cells, since these two new cells continue to grow and repeat the dividing process. This method of multiplication goes on with great rapidity, frequently within a half hour, so that many millions of plants may be formed in a day from a single bacterium. This accounts for the rapidity with which solutions, as beef tea, in which bacteria thrive, become turbid and also for the gelatinous masses and wrinkled

scum that appears upon cooked foods and soups, if allowed to stand undisturbed for a day or more (Fig. 91, *A*, 3). Conn estimates that a single bacterium, if allowed to develop under perfect conditions, would produce in three days a mass of bacteria weighing 4,700 tons. When the conditions are unfavorable for growth many species of bacteria have another form of reproduction and form small bodies, called spores, within the cell (Fig. 91, *A*, *D*). These spores are exceedingly resistant to unfavorable conditions but germinate readily, producing bacteria when conditions are again suitable for growth. Some spores will stand boiling for more than one hour. This peculiarity of the spore, while of great advantage to the bacteria, renders them very difficult to exterminate.

(*b*) *Exclusion of Bacteria.*—The various devices for preserving fruits and meats and sterilizing utensils and instruments in surgery is based upon some method of killing these organisms and preventing the entrance of others. This is the purpose of boiling and canning foods and of salting or drying fruits and meats. The boiling, if thorough, effectually sterilizes while the canning excludes all organisms and the material will remain unchanged indefinitely. Smoking or drying, with addition of salt in the case of meat and of sugar in fruits, is effectual since bacteria flourish only in the presence of moisture. This is the reason that seeds and other structures are not speedily destroyed by bacteria. Seeds are naturally dry and protected by coats that tend to exclude the moisture for some time. Sugar is used as a preservative because bacteria cannot endure over 40 or 50 per cent. of it. Salt for the same reason is useful in preserving meat because it is injurious to bacterial life and smoking also introduces volatile products that prevent their growth.

(*c*) *Economic Importance of Bacteria.*—Insignificant and simple as are the forms and structures of bacteria it is safe to say that no plants are so vitally related to our welfare. A small number of species construct necessary food products; others form certain mineral compounds; some remove the remains of animals and plants through putrefaction and decay; and others pro-

duce a form of decay called fermentation. It is also a familiar fact that many diseases are due to these plants.

Attention has already been called to certain bacteria of the first group mentioned above as very important plants in keeping up the fertility of the soil. These bacteria form tubercles in the roots of plants belonging to the bean family where they combine the free nitrogen of the air into compounds that can be absorbed by the plants, page 57. Another very important kind that may be considered in connection with the tubercle-forming bacteria, live in the soil and change the simple nitrogen compounds, as ammonia, that result from the decay of plant and animal life into nitrites. For this reason they are commonly called the nitrifying bacteria. Were it not for these bacteria the ammonia compounds would quickly escape from the soil as a gas and thus the principal source of nitrogen in nature for the plant would be lost.

Numerous forms of bacteria produce deposits of sulphur and iron by oxidizing the lower compounds of these minerals and are thus the cause of the white and yellow slimy masses of sulphur and iron in springs and streams. Some of these bacteria appear to be independent of organic matter for their support, deriving the necessary energy for their growth from the decomposition of inorganic material. It should be added that the nitrifying bacteria effect the same interchange of gases as noted in photosynthesis and that some of the sulphur bacteria possess a reddish pigment that decomposes carbon-dioxide in the same way. This substance may be related to chlorophyll and it has been suggested that it may represent an earlier state in the evolution of chlorophyll.

(*d*) *Bacteria of Decay.*—The great majority of bacteria live as saprophytes upon organic matter. There are two classes of these plants. The first class live upon nitrogenous matter, causing a decomposition or decay known as putrefaction and thus removing the remains of the animals and plants. The immense importance of this in the economy of nature is evident. The accumulation of material not only is prevented but changes of decay, by a long series of chemical reactions, break down the

complex substances into inorganic and simple organic compounds that serve as food for plants. This in brief is the story of the organic world. Plants construct complex compounds from simple ones and die. Bacteria now reduce them to simple substances which become again the food of succeeding generations. Thus bacteria assist in keeping up the fertility of the soil and prevent the burying of the earth under the annual fall of vegetation. The second class, in common with some fungi, live upon sugars and other carbohydrates, causing a decomposition or decay known as fermentation. Vinegar is due to the decomposition of the alcohol in cider and other weak alcoholic solution by bacteria which appear as slimy masses and are popularly known as mother of vinegar. Milk sours and coagulates through the agency of bacteria which reduce the sugar to lactic acid which in turn coagulates the casein. So also butyric acid, necessary in the manufacture of cheese is produced by bacteria. It will be seen from this that the products of decomposition are not necessarily harmful. It is interesting to note that some of the flavors of cheese and of high grade butter are due to the products of decomposition and the excretions from bacteria. It is evident that this might be true in the case of limburger cheese and rancid butter but also remember that species of bacteria are cultivated and used to impart certain flavors to cheese and also to a less extent to butter. It is altogether probable that this will become a common practice in our dairies. Some bacteria bring about decomposition and produce secretions that are exceedingly poisonous. The ptomaines found in fish, cheese, ice cream, etc., are due to this cause.

(*e*) *Bacteria of Disease.*—In contrast to the kinds of bacteria mentioned above there is another group that live as parasites on plants and animals producing disease either by destroying the tissue and sapping the vitality or by the production of poisonous compounds, toxins. Among the more terrible of these infectious bacteria may be mentioned those producing consumption. These affect especially the lungs of animals and cause over 30 per cent. of the deaths of the human race. This is no longer considered so fatal a disease as formerly. A person of ordi-

nary constitution, if suitably nourished, can be cured by living in the open air. This pest could probably be wiped out within ten years, if the habit of spitting could be stopped. The spores and plants are readily carried away in the air from the dried sputum to be inhaled again and so spread the disease. The bacteria causing lockjaw, or tetanus, are especially abundant in the soil in certain localities. Through wounds they are carried into the system. These and some other bacteria are peculiar in that they can not grow in the presence of oxygen. Consequently the dangers of infection are great in deep wounds. An immediate cauterization of the entire surface of the wound will kill the organisms. Other well known bacterial diseases are pneumonia, diphtheria, Asiatic cholera, typhoid, and splenic fevers and grippe. It should be noted that many diseases as smallpox, malaria, hydrophobia, yellow fever and probably scarlet fever, etc., are due to a low order of microscopic animal life. These bacteria are either localized, as in the lungs, throat, intestines, or are generally distributed throughout the system. The symptoms of the disease are largely due to the products of decomposition caused by the growth of the bacteria or by secretions from the bacteria themselves all of which act as poisons and are commonly referred to as toxins. In other cases the toxins are formed by the infected plants or animals owing to the disturbance of its growth and nutrition by the bacteria in its tissues.

These bacteria are constantly finding their way into our bodies by means of the breath but so long as the system is in a healthy condition there is only slight danger of the organism gaining a foothold. Just how the system is able to combat the growth of these plants is now the subject of earnest investigation. Certain diseases, as mumps, measles, whooping-cough, smallpox, etc., usually occur but once and it is probable that substances are formed in the body as a result of the presence of the bacteria or animal organisms that prevent a second growth of the germs and the individual thus becomes immune to further attacks. The formation of such a substance is known to occur in the case of diphtheria. The bacteria of this disease occur in the upper air passages of the throat and secrete a poison or toxine that affects

the entire body. Gradually the system forms a substance called an antitoxine that stops the growth and eventually kills the bacteria. The vigor of the system and the virulence of the bacteria decide whether the disease shall prove fatal before the antitoxines are formed in sufficient quantity to destroy the bacteria. Note that these disease-producing bacteria vary in their power to produce toxine. Sometimes, as in epidemics, they are so virulent as to attack with fatal results the strongest individuals. In other cases the bacteria appear to be degenerate and only produce mild symptoms of the disease. This is the principle of vaccination. The little animals causing smallpox are cultivated under conditions that weaken them and render them less virile. Consequently when they are introduced into the system they produce only mild symptoms of the disease. The history of the study of bacteria constitutes one of the most interesting and fascinating pages in science. Pasteur was the first to demonstrate that fermentations and putrefactions, and later that certain contagious diseases, were due to bacteria and animal organisms. His work is now commemorated by a tablet on the rue Pasteur in Paris with the inscription: Here stood Pasteur's laboratory. 1857 Fermentations; 1860 Spontaneous Generation; 1865 Diseases of Wines and Beers; 1881 Virus and Vaccini; 1888 Silkworm Distempers; 1864–1888 Hydrophobic Remedies. To-day every state and government has its corps of workers and every city its expert bacteriologists who are studying the nature of bacteria and their relation to plants and animals.

Class B. Blue Green Algae or Cyanophyceae

63. The Structure and Nature of the Cyanophyceae.—These plants are very simple organisms that have some features in common with the bacteria. They are unicellular plants, although the cells are more commonly joined into rows, forming a thread or filament (Fig. 92, *A, B*). In some cases these filaments may be regarded as multicellular plants since they branch and because the various cells perform different functions. They form slimy, blue-green, black, yellow or violet masses and, like the bacteria,

appear to be dependent to a degree on organic material. At least they are specially abundant in the presence of decaying organic matter, as in the drainage from stables and watering troughs, or in pools, puddles and damp places where there is an abundance of filth and decay. The enormous increase of these plants often produces a discoloration of the water, as in the Red Sea, and they are often the cause of the foul odors of muddy ponds and streams and reservoirs and of the pollution commonly known as water-bloom or working of ponds. Like the bacteria also they are associated with varying amounts of gelatinous substances derived from their walls or excreted from the cells and in both cases the walls may become colored with various pigments. Doubtless the mucilaginous character of these plants enables them to retain moisture and so adapts them to dryer conditions than would otherwise be possible. The Cyanophyceae differ radically from the bacteria in possessing chlorophyll which is distributed in the outer portion of the protoplasm, and in the possession of a more definite nucleus. A blue pigment phycocyanine, is associated with the chlorophyll and for this reason they are called the blue-green algae or cyanophyceae. Cell division follows the ordinary method noted in the higher plants, at least this appears to be true in those cases that have been accurately studied. These plants accordingly show a decided advance over the preceding forms not only in the differentiation of the cells but especially because they are capable, to a degree at least, of manufacturing food from inorganic substances by reason of their chlorophyll.

In addition to the multiplication of individuals by cell division, as in the case of the bacteria, many of the thread-like forms have another interesting method of increasing their numbers. A few of the cells, termed a hormogonium, become separated from the other cells and frequently possess for a time a slight motility that enables them to move away from the parent filament (Fig. 92, *hr*). In some species one or several rather large, colorless cells, heterocysts, appear in the filament and may serve to separate it into hormogonia (Fig. 92, *h*). In *Oscillatoria* (Fig. 93) the filaments into which these hormogonia grow per-

manently retain their power of motion. No cilia have been observed for a certainty in any of these forms and the cause of the motion has been ascribed to the interchange of fluids attend-



FIG. 92.

FIG. 93.

FIG. 92. Forms of the Cyanophyceae: *A, Gleocapsa.* At the right a cell has divided, but the two daughter cells are held together in a gelatinous mass. Above numerous divisions have occurred, but all the cells are surrounded by a mucilaginous envelope. *B, Nostoc.* Below appear the gelatinous, spherical masses of the plants as they appear floating upon the surface or resting on the bottom of ponds. At the right a plant enlarged—*h,* heterocyst; *s,* thick-walled resting cells or spores. At the left a spore has germinated, producing five cells. *C, Rivularia.* At left gelatinous mass of plants attached to stem of water plant. At right view of the plants enlarged—*hr,* hormogonium; *h,* heterocyst.

FIG. 93. One of the most common forms of the Cyanophyceae, *Oscillatoria.* The different sizes of the cells show that cell division may occur in any of the cells of the filament—*e,* a decaying cell which will ultimately free the cells below it as a hormogonium.

ant upon the absorption and digestion of foods. A simple type of spore production is also to be found in the majority of forms. This consists merely of a slight enlargement and thickening of the walls of the ordinary cells (Fig. 92, *s*). These simple spores are able to tide the plant over unfavorable conditions and germinate, forming new plants when conditions are again suitable.

The Bacteria and Cyanophyceae are evidently very primitive and ancient forms of life. The fact that they frequently occur in hot springs and that they can endure greater extremes of heat than higher plants may indicate that they are related to forms that appeared upon the earth when just such conditions existed and at a time when the environment was not suitable for the development of higher types. It has been stated on page 142 that certain bacteria are independent of organic foods and that others effect a decomposition of CO_2 as in photosynthesis while in the purple and red sulphur bacteria we have pigmented forms that are perhaps in a transition state to chlorophyll bearing plants. It seems not at all improbable that the earliest forms of life could build up organic compounds without chlorophyll and in such forms as these bacteria we have perhaps an illustration of the gradual acquisition of chlorophyll as seen in the higher plants.

Subdivision 3. Diatomaceae or Diatoms.

64. The Nature and Structure of Diatoms.—The diatoms are among the most common and widely distributed plants (Fig. 94). While microscopic they exist in such large numbers as to form the familiar brown coatings on the bottom of ponds and sluggish streams and often render sticks and stones in the water slippery with slimy deposits. They are widely distributed on damp soil and in fresh and salt waters. Note should also be made of the peculiar species that float upon the surface of the ocean and form the larger part of that organic life known as the plankton (Fig. 95). This is especially abundant in northern waters and has been compared to great pasture lands since it furnishes the principal food of surface feeding fishes and other

marine life. The Diatoms are unicellular plants and exhibit a great variety of forms as circular, elliptical, rod or wedge-



FIG. 94. Common forms of diatoms: *A*, fan-shaped colonies of diatoms, *Licmophora*, attached by gelatinous stalks to seaweed. The repeated division of a solitary diatom gradually builds up the fan-like colonies. *B*, *Tabellaria*, forming colonies grouped in zig-zag blocks. *C*, *Melosira*, cylindrical pill-box-like diatoms, arranged in chains. Below end view of the diatom. *D*, a species of *Navicula* arranged in gelatinous branches. At right enlarged view of a branch in which the diatoms glide back and forth.—After Wm. Smith.

shaped, curved or straight. They are equally variable as regards their association. Some are solitary and free swimming, others are attached by stalks; some form bands or ribbons or zigzag

chains, while others are imbedded in a gelatinous mass of a more or less regular form (Fig. 94, *D*). The cells are covered by two valves, one of which overlaps the other like the cover of a box. Therefore a diatom presents two quite distinct ap-



Fig. 95.



Fig. 96.

Fig. 95. Plankton forms of diatoms: *A, Coscinodiscus, B, Planktoniella.* Below seen from side.—After Gran.

Fig. 96. Structure of the diatom, *Pinnularia: A,* valve view. *B,* girdle view. *C,* cross-section, the two dark bands showing the position of the chromoplasts in the diatom—*p,* pore in wall which appears as a line in *A,* running from the ends towards the center of the valve.—After Lauterborn.

pearances—the top or valve view and the side or girdle view (Figs. 96, 97). This difference is further intensified by the sculpturing of fine lines that appear upon the walls. An interesting feature about these valves is the fact that they are com-

pletely infiltrated with silica, a substance resembling glass. If a diatom is burned in a flame on a strip of platinum or placed in acid to remove the organic substance the appearance of the valves remains unchanged.

The glass-like valves are quite transparent and it can be readily observed that the cell contents is much more highly differentiated than in the preceding group. The chlorophyll is deposited in plastids of definite form, although this color is often masked by a brown pigment which causes the characteristic appearance of these plants when associated in masses. The oil drops seen in the cells are the product of photosynthesis—starch not being formed. Some species, however, can live upon decaying organic matter and in consequence contain colorless plastids. In



FIG. 97. Structure of *Cymbella: A*, valve view. *B*, cross-section showing the difference of the two sides of the diatom. *C*, the two girdle views. —After Pfitzer.

the free swimming forms the motion consists of an irregular gliding movement and this is supposed to be due to the expansion and contraction of minute strands of protoplasm that project through the pores of the valves (Fig. 96, *C*, *p*).

(*a*) *Reproduction of the Diatoms.*—It will naturally be asked how can these plants living in glass houses, grow? As the valves become changed to silica naturally any increase in size must cease. Nevertheless the cells reproduce with great rapidity and in a very interesting manner. Through the growth of the living

substance the valves are pushed apart and the cell contents divides, forming two diatoms with but one valve each. A new valve that fits into the old valve is soon developed on the uncovered side of each diatom and two complete diatoms are thus formed (Fig. 98) which may become free at once or remain at-



FIG. 98. Diagram illustrating the divisions of a diatom and the resulting reduction in size. The brackets connect the two diatoms that were formed by each successive divison.—After Pfitzer.

tached and by further division form the odd groupings or colonies shown in Fig. 94. Since the valves are of unequal size, *i. e.*, one fitting into the other, the two diatoms formed by the division must vary in size and it will be seen that the majority of the offspring will become greatly reduced in size as a result of the rapid and repeated divisions. That this reduction may not go on too far, when a certain minimum size has been reached the cell contents, with or without dividing, throws off the valves entirely and grows to the full size of the diatom, when new valves are formed and the diatom is ready to repeat the dividing process. In some cases we find quite a different method of reproduction. Two diatoms become enclosed in a jelly-like mass, into which the cell contents, with or without dividing, is discharged. These naked cells now unite in pairs and the bodies thus formed finally grow into diatoms (Fig. 99). This latter method of reproduction is termed sexual reproduction. The two fusing cells are looked upon as male and female cells or gametes although in these simple plants there is no indication of sex that would enable us to recognize them as such. In the preceding groups, multiplication in numbers or reproduction has been effected by the division of the cells or by the formation of more or less modified cells, termed spores. All such methods of re-

production are called asexual since but one cell is utilized in the process. In sexual reproduction a cell is formed by the union of the contents of two cells. The nuclei and protoplasmic contents of each unite so completely that a cell with but one nucleus and protoplasmic contents results. These sexually produced cells are called gametospores because they are formed by the union of two gametes and are capable of forming a new plant in the same way as do spores. We may distinguish the spores derived from single cells as asexually formed spores or we may call them simply spores, while the spores formed by the fusion



FIG. 99. Sexual reproduction of *Pinnularia: A,* two diatoms enveloped in a mass of jelly. The valves have been thrown off and the contents of each diatom have divided into two sexual cells or gametes. *B,* the fusion of the gametes.—After Karsten.

of the two cells or gametes may be termed sexually formed spores or gametospores. Both kinds of spores are devices to enable the plant to increase in number or to meet some other problem in life, such as conditions unfavorable for growth. The spores are especially to be looked upon as devices to bring about a rapid increase in the number of individuals while the gametospores more usually serve to carry the plant over periods of drought or extremes of temperature which would prove fatal to the plant. For this reason the gametospores are often provided with thick walls and dense cell contents which enable them to

remain in a dormant condition until conditions are favorable for growth. The gametospore is often called a resting spore for this reason.

So rapidly do the diatoms multiply that the larger part of the soil and rocks in many localities is composed of their remains. The constant casting off of the valves through reproduction or death results in vast deposits, known as silicious earth, on the bottom of ponds and in the sea. Beds of silicious earth formed in this way are often seen in districts where ponds and lakes have dried up. Earth of this nature is used as polishing powders and as absorbents in the manufacture of some explosives.

Subdivision 4. Euphyceae or Algae

65. General Features.—This subdivision includes a large number of plants that live chiefly in fresh or salt water. These plants, popularly known as Algae, vary greatly in form and structure and range from microscopic unicellular forms to some of the largest and most highly constructed plants found among the Thallophyta. Their advance over preceding groups appears especially in their well-marked walls and distinct plastids and nuclei (Figs. 100, 104). Chlorophyll is always present in the cells, although in certain groups it is masked by brown or red pigments. These latter pigments are supposed to adapt the plant to varying intensities of light. This is supported by the fact that the marine Algae exhibit a zonal distribution in the water. Forms living near the surface of the water are predominantly green, at depths where they are alternately exposed and covered by the tide, they are more commonly brown, while in the shade of the brown Algae, or, below tidal limits, red forms occur. The Algae appear to have developed along three lines which are indicated by the green, brown and red colors, although the basis for this classification rests upon structural and reproductive characters. These three classes are: A, Green Algae or Chlorophyceae; B, Brown Algae or Phaeophyceae; C, Red Algae or Rhodophyceae.

Class A. Green Algae or Chlorophyceae

66. The More Imporant Orders of the Chlorophyceae.—The Chlorophyceae are one of the most interesting groups of the Algae, because they contain primitive forms that are at least suggestive of low animal types and they also exhibit a gradual modification of the plant body and reproductive processes that help us to understand how variations arose and how complex types have been evolved from very simple forms. Starting with unicellular motile forms, the green algae appear to have diverged along several lines. Among the more important of these may be mentioned the following orders: (*a*) Volvocales, largely unicellular green Algae: (*b*) The Zygnematales or conjugating green Algae; (*c*) Chaetophorales or filamentous green Algae; (*d*) Siphonales or tubular green Algae.

67. Order a. Volvocales or Unicellular Green Algae.—The lower members of this order represent a very primitive type of plant. Their motility, certain features of their life history and delicate cell walls are suggestive of some of the preceding groups, as well as of low forms of animal life. *Sphaerella* (Fig. 100) is a familiar example of this group, often appearing in rock hollows as blood-red stains. If some of this red material is examined in water, the plants appear as spherical cells with dense red protoplasmic contents and thick walls (Fig. 100, *A*). This condition represents the resting state of the plant. If the plants, after being dried, are allowed to stand in water for a few hours, the nucleus will divide, forming usually from 4 to 16 daughter cells (Fig. 100, *B*) which escape by the rupturing of the old mother wall and swim actively about by means of two cilia. These cells, or zoöspores, remain motile for varying periods, during which time they increase in size, but finally the cilia are retracted and the plants remain in a quiescent state for a short period (Fig. 100, *E*). A division of the nucleus now results again in the formation of two or more zoöspores (Fig. 100, *F*) which repeat the life history noted above. This method of reproduction may occur again and again and thus rapidly bring about a great increase in the number of plants. During the

divisions, the red coloring matter is usually rapidly replaced by chlorophyll until but a red speck remains and the zoöspores soon appear as rather ovate green cells surrounded by delicate walls which become widely separated from the chlorophyll owing to the accumulation of water (Fig. 100, *G*). Nearly all zoöspores



Fig. 100. Stages in the life history of *Sphaerella: A,* resting state of the plant. *B,* first division. *C,* second division, the four cells are about to escape from the mother plant. *D,* one of the cells of *C* after escaping. This is a zoöspore of the first generation. *E,* zoöspore at rest. *F,* forming four new zoöspores. *G,* one of these zoöspores of the second generation. Note that the red material, represented by the shaded area in the center of the cell, has become greatly reduced and that the wall is becoming distended and separated from the granular cytoplasm. *H,* third resting stage. *I,* cell dividing. *K,* zoöspore of third generation with greatly distended cell wall and small red area. Delicate strands of cytoplasm connect the cell wall and the central protoplasmic body. *L,* a resting cell dividing into a large number of zoöspores which are consequently smaller.—After Hazen.

that occur among the algae are characterized by a small red body, known as the eye spot, which is located near the ciliated end of the zoöspore. It has been supposed that this body is sensitive to light and gives the zoöspore a sense of direction. While this

is questionable, since the eye spot may be lacking (as in *Sphaerella*), it is certain that these beautiful bodies swim about, ciliated end foremost, with a rotary motion from right to left and adjust themselves to a suitable illumination. It can easily be demonstrated that the zoöspores of *Sphaerella* are keenly sensitive to different intensities of light by placing them in a glass dish by a window when the zoöspores will congregate on the illuminated side unless the light is too intense when a reverse action takes place.

Thus we see that the life of the plants is largely a motile one, each generation being characterized by a short resting stage during which division occurs and a longer motile zoöspore stage. In fact, the zoöspore may be looked upon as the original state of these plants, while the resting condition is a departure due to changes in the environment or the condition of the organism. This common occurrence of motility in the lower types of life indicates that possibly such was the condition of the first life upon the earth.

(*a*) *Conditions Affecting the Life of Sphaerella.*—Changes in the surroundings sometimes produce remarkable variations in the life of the plant. This fact is well illustrated in *Sphaerella.* If the water dries up, a thick wall is formed about the central protoplasmic body which becomes dense and of a deep red color while the delicate distended wall and cilia disappear. This variation adapts the plant to conditions unfavorable to growth, such as drought and severe temperature. The so-called red snow is due to a certain species of these plants that are swept off from the rocks in this resting condition by the winter winds and falling to the earth produces red streaks in the snow. Hazen has been able to show that a most suggestive change in the mode of life of the plants may also be induced by low temperature or by a reduction of the volume of water in which they live. Under such conditions, bodies are formed that resemble zoöspores, but that are not capable of motion owing to the lack of cilia. This helps us to understand how the motionless or stationary forms of plant life came about and how the motile stage became less and less conspicuous. Owing to such causes as these and many

others, plants become stationary. At first, perhaps, simply an aggregate of motionless cells, but finally there resulted chains or filament of cells owing to the repeated division of the plants in one plane. By division in two planes, membranous expansions resulted and the more complex types arose by the division of the cells in three planes and by modification of the cells.

One other feature appears in the life of the *Sphaerella* that indicates clearly how sexual reproduction came about. When the conditions for growth are unfavorable, as for example, through lack of moisture or low temperatures, there result plants that are not so well nourished, or, at least they appear to lack some of the qualifications that characterize the plants growing under favorable conditions. The same condition often arises in these plants after several generations have been formed. Such plants, which may be characterized as weaker, or, at least lacking in certain material, do not give rise to the ordinary zoöspores, but to much smaller though similar bodies that are lacking in cell walls (Fig. 100, *L*). These small zoöspores usually perish or produce small plants unless two of them meet, when a fusion may occur and thus a plant is formed that is capable of repeating the life history of *Sphaerella*. By suitable nourishment, however, these small zoöspores may sometimes be made to develop into the normal plant. It is very evident that these small zoöspores are lacking in some substance that is essential to their growth. This is rarely furnished to them in nature except when two of the bodies fuse. From numerous examples appearing among different groups of Algae, it appears that sexuality arose in this way. Owing to certain conditions of light, temperature, food, etc., zoöspores were formed that were incapable of further growth. But, by the union of two of these zoöspores, a cell was formed possessed of renewed vigor and capabilities of growth. So we can think of the sexual cells or gametes as zoöspores that are lacking in the materials essential to growth and of the sexual process as a union of the two bodies for the purpose of bringing together the missing material and supplying the necessary energy for growth.

A closely allied genus, *Chlamydomonas* (Fig. 101) has a life

history very similar to that of *Sphaerella* but the formation of the gametes reveals a variation that gives us an understanding of how these bodies came to differ and finally became distinguishable as male and female gametes. In some of the species of *Chlamydomonas,* to be sure, the gametes are similar and can not be distinguished as male or female (Fig. 101, *B*). This stage of sexuality is called isogamous, meaning similar gametes. In other species, there is a decided variation in the size of the gametes, certain cells or plants producing large and less active



Fig. 101.



Fig. 102.

Fig. 101. Features in the life history of *Chlamydomonas: A,* character of the mobile plant. *B,* conjugation of isogamous gametes. *C,* a cell dividing to form numerous small male gametes. *D,* a plant forming two large female gametes. *E,* male and female gametes about to conjugate.

Fig. 102. Features in the life history of *Pandorina: A,* a colony of plants. *B,* each plant of the colony dividing to form a new colony. *C,* the plants of a colony escaping as gametes. *D,* the conjugation of two gametes of unequal size. *E,* later stage in the conjugation. *F,* gametospore or resting spore. *G,* large zoöspore formed from the gametospore. *H,* a colony formed by the division of the zoöspore, *G.*

cells that are termed female gametes, while other plants produce numerous minute male gametes that swim actively about (Fig. 101, *C, D*). This stage of sexuality, where the gametes differ

in size, is called heterogamous, meaning unequal gametes. This variation in the size of the gametes which we call the differentiation of sex is due to the amount and nature of the material stored in them, the male gametes being formed in a cell in larger numbers than in the case of the females. These heterogamous gametes do not differ in nature from the isogamous gametes. They are also lacking in some substance that renders them incapable of growth unless fusion of two gametes is effected.

(*b*) *Colonial Unicellular Algae.*—The various stages in the evolution of sex are also well illustrated in several genera of unicellular green algae that live together in colonies. *Pandorina* shows the earlier phases of the differentiation of the gametes. This plant is really a spherical colony of about 16 motile plants, each similar to *Sphaerella* but held together in a mucilaginous mass (Fig. 102, *A*). These beautiful colonies multiply by each plant or cell dividing into 16 daughter cells which finally become an independent colony (Fig. 102, *B*). The gametes are formed in essentially the same way as the colonies, save that the 16 daughter cells finally escape as free swimming bodies (Fig. 102, *C*). The mother plants or cells producing these gametes differ in size, owing, doubtless, to the amount of nutriment which they receive. Consequently, the gametes formed from them must also differ in size. It is also possible that the mother plant may produce a larger or smaller number of gametes and so cause a variation in their size. It is interesting to note that these gametes fuse quite regardless of their size, although there may be a slight tendency for the small and more active gametes to fuse with the larger and slower moving ones. Thus we have an illustration of the first appearance of the differentiation of sex although the sexuality of the gametes is not thoroughly established. The gametospore, resulting from the union of the gametes, develops a thick cell wall, the contents assumes a reddish color and it passes into a resting stage (Fig. 102, *F*). When conditions are favorable, it germinates as shown in Fig. 102, *G–H*. In the beautiful spherical colonies of *Eudorina* and *Volvox* (Fig. 103) the differentiation of the gametes is complete. Certain cells produce but one large female gamete that remains motionless

in the mother cell while other cells form numerous small motile male gametes which are yellowish in color. In *Volvox,* over 200 male gametes may be formed from a single cell (Fig. 103, *D, E*).

This tendency to form gametes of different sizes is manifestly of great advantage. The great number of male cells increases the chances of fusion of the gametes and their smallness promotes their movements through the water. The increase in the size of the female gamete renders her movements more difficult, but the storage of food in this gamete provides for the



FIG. 103. Features in the life history of *Volvox: A,* a colony which may contain as many as 20,000 plants. The three central groups are young colonies which may arise by the repeated division of any of the plants. *B,* side view of one of the plants showing the canal-like connections with the neighboring plants. *C,* surface view of a plant with canals radiating out to the adjoining plants. *D,* a plant enlarging and forming a single motionless female gamete. *E,* a plant forming numerous small male gametes. *F,* male gamete enlarged.

better nourishment of the next generation. Finally the nourishing function becomes so strongly developed that the female ceases to move at all and remains protected in the mother cell. Thus we see that sexuality arose among plants as the result of

definite stimuli as the temperature, light, food and many other conditions that occur in the environment. These stimuli produce enfeebled zoöspores that are incapable ordinarily of growth unless a fusion of two of them is effected. So sexuality arose from the asexual state of the plant as a result of the influence of the environment and differentiation of sex appeared owing to the variation in the amount and kind of material in the gamete producing cells.

(*c*) *Motionless Unicellular Algae.*—In contrast to these motile plants, there is a large group of the Volvocales that have lost their motility and spend a large part of their life as stationary plants. We have already noticed how this condition may have come about, owing to the temperature and moisture conditions that were unfavorable for the motile existence. *Pleurococcus* (Fig. 104, *A*) is one of the most common of the green



Fig. 104. Common forms of stationary unicellular green algae: *A, Pleurococcus* showing stages in the division of the plants, the nucleus and the irregular chloroplast. *B, Scenedesmus.* Below three of the plants are shown dividing. *C, Ankistrodesmus.* At right three plants in process of division.

algae and illustrates many of the features in the life history of the stationary forms. These plants often cause the green coating that appear upon the shaded or moist side of tree trunks, fences and buildings. If a bit of this material is examined under the microscope, it will be seen that the plants are rather spherical cells provided with nucleus, cytoplasm, chloroplasts and well marked walls. This latter feature is characteristic of stationary algae and forms a sharp contrast to the delicate walls of the motile forms. These plants increase in numbers by the formation of a wall through the middle of the cell. The two daughter cells or plants grow to the size of the mother plant and repeat the process in division by forming a wall at right angles to the old wall. It will also be noticed that cell division may occur in three planes. The plants adhere together for a considerable period, or become detached shortly after the division of the mother cell. In this way they multiply with extreme rapidity and so come to form extensive green coatings. It has been noted in allied genera and reported in certain species of *Pleurococcus,* also that changes of temperature and amount of water, etc., will cause the contents of the cell to divide and form zoöspores which are of the same nature and behave in the same way as in *Sphaerella.* Thus we see how changes of the environment may cause these stationary plants to return to the motile condition of simpler types. It is interesting to note in this connection that Livingston was able to change the stationary condition of certain algae to the motile condition at will by diluting the solutions in which they were living. Fig. 104, *B, C* illustrates several other forms of stationary green algae that are common in the drinking water and in ponds and streams. Interesting forms for study may be obtained by tying a woollen sack to a water faucet and allowing the water to run into it for several hours. Invert the sack into a glass, washing off the material that collects on the inside of the sack in the water of the glass. By means of a pipette, a drop of the material that collects at the bottom or sides of the glass or forms on the surface of the water may be transferred to a slide and examined under a microscope.

Some of these forms are free, some attached and others aggregated in a mucilaginous mass.

There are a great many of the stationary genera that show the same habit of uniting into colonies that we noticed among the motile genera. These colonies may be very simple and composed of a few loosely aggregated plants forming flat discs or star-shaped plates (Fig. 105, *A*). In other cases, we find spherical and net-like colonies. The water net, *Hydrodictyon,* is a common illustration of this latter arrangement (Fig. 105, *D, E*).



FIG. 105. Colonial forms of unicellular green algae: *A, Pediastrum,* the plants of the colony being arranged in a flat plate. *B,* a view of the outer cells of the colony showing the formation of a new colony. *C,* one of these new colonies. *D,* a plant of the water net containing a young colony. *E,* enlarged view of one of the meshes of a net showing the geometrical arrangement of the plants.

These plants occur in ponds and sluggish streams, forming sac-like nets three to four inches long, that are constructed in a very regular manner. The reproduction is quite characteristic of many of the Volvocales. Zoöspores are formed in large numbers in any of the plants and after a period of motility, they come to rest and arrange themselves into a new net inside of the

mother plant (Fig. 105, *D*). The walls of the mother plant soon break down and the new colony is set free. Under different conditions, smaller but similar bodies are formed which escape from the mother plant. These bodies are gametes and incapable of growth unless a fusion is effected between two of them. Especial interest attaches to the study of this plant because Klebs was able to change the reproduction character of the plant by subjecting it to certain definite conditions. By exposing the plants to bright light, slight increase of temperature and inorganic culture solutions of definite composition, zoöspore are produced. On the other hand, plants growing in solution of cane sugar in subdued light or darkness formed gametes.

These reactions are of great importance in helping us to understand the nature of plant life. They demonstrate that definite factors may change motile plants to a stationary condition and in *Hydrodictyon* we see that measurable factors, as light, heat and concentration of the solution may bring about the motile condition of the plant and may also cause a variation in the motile condition which we call sexuality.

68. Order b. Zygnematales or Conjugating Green Algae.— This order includes the most common and attractive forms of green algae (Figs. 106, 108). They are of very general occur-



FIG. 106. Two common forms of the Zygnematales: *A, Spirogyra*— *n*, nucleus; *p*, pyrenoid. *B, Zygnema.*

rence in the still waters of ponds and streams where they often form the large floating and frothy green masses popularly known

as pond scum or frog spittle. They can usually be recognized by their slimy nature which is due to the copious exudation of a gelatinous substance from the cell walls. In one group of this order the plants are multi-cellular (Fig. 106), the cells being attached end to end and forming a filament—though it should be stated that each cell is practically independent in its life process and the organism may therefore be regarded as a colony. Such a type of plant would have originated in the case of *Pleurococcus* if the two daughter cells had remained attached and continued to divide in but one plane and parallel to the first division wall. This is exactly the mode of growth in these filamentous plants. Any of the cells may divide into two daughter cells which grow to the original size of the mother cells. In this way the length of the plant is increased. The most characteristic feature about the Zygnematales is the remarkably large and attractive chloroplasts. In *Spirogyra* (Fig. 106, *A*) these appear as bands, from one to several in each cell, spirally wound around the cytoplasm just within the cell wall. In *Zygnema* (Fig. 106, *B*) they assume the form of stars and in other genera they appear as plates, bars and variously modified bodies. Imbedded in the plastids are denser protoplasmic bodies, the pyrenoids, that appear to be connected with the secretion of starch; at least a layer of starch can be detected about the pyrenoids by testing with iodine (Fig. 106, *pr*).

The filaments are fragile and readily become dissociated into smaller portions. In this way, the plants may multiply since the fragments continue to grow after the manner of the original filament. Singularly, there is no indication of a return to the motile condition in the life history of these plants. No zoöspores or motile gametes are formed. Sometimes the walls of a cell become thickened and after a dormant period, as through the winter, may germinate and grow into a new plant. With this rather rare exception, the asexual reproduction of the plant by spores, does not occur. The sexual reproduction is of a peculiar character. This is very well illustrated in *Spirogyra*. When filaments of this plant chance to lie side by side, under certain conditions, tube-like processes from opposite cells begin to grow out towards each

other (Fig. 107, *A*). These tubes finally meet, when the walls at the point of contact are absorbed, forming an open passage between the two cells. While this growth has been going on,



FIG. 107. Sexual reproduction of *Spirogyra: A,* in lower portion of figure the formation of the tubes between the opposite cells of the filaments is shown. Above the contents of two cells has united with two cells of another filament, forming two gametospores. *B,* germination of a gametospore.

the contents of the two cells has contracted somewhat and finally the contents of one cell passes through the tube and fuses with the contents of the other cell. In other genera, there is a considerable modification in the process, as in some species of *Zygnema*, where the fusion is effected in the middle of the tube. This

is only a different form of the sexual process with which you are familiar, the difference being, that but one gamete is formed in each cell and these bodies are motionless or at least not provided with cilia and do not escape from the mother plant. When the fusion of the gametes has been effected the resulting gametospore becomes surrounded by a thick wall and in this resting condition it tides the plant over seasons, such as the winter or drought, unfavorable for the growth of the plant. When conditions return that permit the growth of the plant, the gametospore germinates by rupturing the outer wall and protruding the inner wall of the spore as a delicate tube (Fig. 107, *B*). As this tube continues to elongate the characteristic plastids and cell partitions arise, thus forming a new plant. The loss of motility in the reproductive process in this order may possibly be associated with their exposure to terrestrial conditions. They are often exposed to the soil, owing to the drying up of the water and it has been suggested that they have consequently lost their motility. Their mucilaginous coatings would be of great service in enabling them to meet such conditions by retaining moisture, and indeed, they are often able to flourish in many damp places without the aid of surface water.



FIG. 108. Common forms of Desmids: *A, Closterium. B, Micrasterias. C, Xanthidium. D, Straustrastrum.* At left top view, at right side view. *E, Desmidium* forming a chain of plants. At right end view of chain.

The Desmids.—A second group of the Zygnematales are strictly unicellular plants known as Desmids (Fig. 108). They are the most attractive of unicellular plants and are of common occurrence associated with coarse algae. The desmids are elaborately and variously fashioned but can readily be recognized by the fact that they consist of two similar halves (Fig. 109, *A*). The structure of the cells and the sexual method of reproduction is essentially the same as in the filamentous forms which have probably been derived from these unicellular plants. The asexual method of reproduction, however, is rather peculiar. The region connecting the two halves (Fig. 109, *A, i*), called the isthmus, becomes somewhat elongated and swollen. Soon a constriction appears midway between these two halves which deepens



FIG. 109. Reproduction of *Cosmarium: A,* asexual reproduction, showing the elongation of the isthmus, *i,* and its gradual enlargement and division to form the new lobes of the desmids. *B,* sexual reproduction. At the left the cell contents of two desmids fusing to form a gametospore. On the right the mature gametospore covered with a spiny coat.

until the desmid is cut in half. Each desmid now consists of one of the original halves and one half of the isthmus. This latter part gradually enlarges and forms the second half of the desmid. Owing to the gelatinous character of the cell walls, the desmids

often become aggregated in masses and form chains of cells (Fig. 108, *E*).

69. Order c. Chaetophorales or Filamentous Green Algae.—Here belong a great array of green algae that grow attached to various objects and appear as simple or more usually as branched filaments which frequently end in hair-like tips. They are of common occurrence at the bottom or margins of springs, wells, streams and ponds. *Ulothrix* may be taken as an example of one of the simple forms of the Chaetophorales. Fig. 110 shows that it consists of a simple chain of cells that divide and so bring about the growth of the plant as in *Spirogyra.* However, it is important to note that we have here a higher type of vegetative structure than in any of the algae heretofore studied. This advance appears in the one or more rather colorless cells at the base of the plant which serve to anchor it to the substratum. In other words, there is apparent a distribution of labor or the performance of definite duties by special parts or cells of the plant body. The evolution of higher forms comes about in much this way. So long as each cell of the plant performs the same functions, there can be little variation in the form and structure. But as these cells are differently stimulated by their environment, they tend to vary in character and so become adapted to the performance of different functions. Perhaps in *Ulothrix* the contact of the substratum acted as a stimulus, as Pierce found to be the case in many algae. Thus the plant comes to consist of different kinds of cells, each of which performs a different kind of work. The complexity of the plant is due to the variations of its cells which are thereby adapted to the performance of special duties. In *Ulothrix,* we have one of the simplest modifications of this nature, one or more of the basal cells becoming slightly changed in form and contents and so adapted in anchoring the plant. The other cells of the plant are practically alike, each containing a single nucleus and chloroplast and therefore capable of manufacturing food and forming the reproductive bodies which are of the same simple type as noted in *Sphaerella.*

(*a*) *Reproduction of Ulothrix.*—The contents of any of the green cells may divide into two or more cells which escape as

zoöspores through an opening formed in the cell wall (Fig. 111, *A*). The cells in which the zoöspores are formed are called sporangia. The zoöspores resemble the motile plants noted above, but after a short motile period they return to the stationary condition and grow into new plants (Fig. 111, *C*). In this way,

Fig. 110.



Fig. 111.

Fig. 110. Lower portion of *Ulothrix,* the basal cells are somewhat modified and the lowest one acts as an anchoring organ. Each of the upper cells contains a girdle-like chloroplast.

Fig. 111. Asexual reproduction of *Ulothrix: A,* a few cells of a filament in the upper cells of which the formation and escape of the large zoöspores are shown, while in the lowest cells a large number of small zoöspores appear. *B,* a large zoöspore. *C,* a young plant formed from *B*. *D,* a small zoöspore. *E,* a young plant formed from *D*.—After West.

the number of plants is rapidly multiplied and it should be stated that the asexual method of reproduction may be looked upon as a means of bringing about a rapid increase in the number of individuals of a species while the conditions for growth are favor-

able. When new conditons arise, such as changes of temperature, food, etc., that would interfere with the normal growth of the individuals then a modification of the contents of the cells and its mode of division results. The cells of *Ulothrix* as a consequence of such alterations divide into larger numbers of motile bodies which are consequently smaller than in the case of zoöspores (Fig. 112, *A*). These bodies may behave as gametes and unite, forming gametospores. The cells in which gametes are formed are called gametangia, sing. gametangium. Thus, *Ulothrix* is controlled in the same manner as *Chlamydomonas* and *Hydrodyction.* It is interesting to note, however, that these small bodies frequently behave as zoöspores and develop into small and weak plants, which fact accounts for the common association in this genus of puny and vigorous plants (Fig. 111, *C, E*).

These small bodies of *Ulothrix* represent a curious intermediate condition between a zoöspore and a gamete where the sexual



FIG. 112. Sexual reproduction of *Ulothrix: A,* a few cells of a filament showing the formation and escape of the gametes. *B,* gamete. *C, D,* stages in the union of the gametes. *E,* gametospore. *F,* gametospore germinating and forming a single-celled plant in which several zoöspores are developing.

character is not strongly enough developed to overcome completely the zoöspore character. Thus, we see again in *Ulothrix*

that definite environmental conditions caused motile bodies to be produced that are lacking in some essential substance and consequently do not have the energy for growth, or at least can only develop into feeble plants. This weakness is overcome by the fusion of two gametes which results in the formation of a cell or gametospore with renewed energy for growth.

The gametospore germinates after the period of rest in a radically different manner from previous cases. The contents of the spore, instead of giving rise directly to a new plant, divides, forming several zoöspores (Fig. 112, *E, F*) which after a period of motility become attached to some object and grow into a new plant. The cause of this peculiar behavior of the germinating gametospore is not known. Some changes must have occurred in it that was lacking in the gametospores previously noted, and as a consequence, it is not able to develop directly into a new plant. This variation in the growth of the gametospore marks the beginning of one of the most important departures in the evolution of plant life and it will become more and more significant as the work proceeds. It is evident that the formation of several zoöspores, each of which produces a new plant, is a decided advance over those types in which the gametospore develops directly into but one new plant. Note also that in *Ulothrix* there are two kinds of zoöspores, one formed from the *Ulothrix* plant and the other from the gametospore, the former being frequently called zoögonidia to distinguish them from the latter. This distinction is of great importance, because most plants have two distinct stages or generations in their life history. One is represented by the plant that bears the sexual cells or gametes and therefore called the sexual plant or sexual generation, or, in short, the gametophyte. The real nature of this plant is to produce gametes, although it may for a time produce zoöspores. The other stage is derived from the gametospore and is called the asexual plant, asexual generation or sporophyte, since this plant can only produce zoöspores. Among the green algae this distinction is not very manifest, the *Ulothrix* plant for example being the sexual plant or gametophyte and the gametospore is the asexual plant or sporophyte, in this case being merely a single cell.

It is very manifest, however, that this single cell or plant is decidedly different from the *Ulothrix* plant because it cannot develop into the *Ulothrix* plant, but can only produce zoöspores. So we see that there are two generations in the life history and that they differ radically in their natures and possibilities of growth. In higher types of plant life it will be seen that the gametospore tends to develop into a more and more complex body or plant and it will be one of the most interesting features of the work to watch the evolution of this asexual plant.

The higher members of the Chaetophorales illustrates the same gradual differentiation of the gametes, as has been noted in the motile forms of the Volvocales. The female gametes become distinguishable because of their larger size and shorter period of motility, while the male gametes are small, owing to the large numbers that are found in a cell or because of the small size of the cells in which they are developed. Finally, in several of the genera we find but a single female gamete which remains motionless in the cell. This condition is very well illustrated in one of the genera of the next order.

70. Order d. Siphonales or Tubular Green Algae.—This order includes a large number of odd forms that are filamentous in character and they differ from all other algae in that the filaments contain numerous nuclei, but with rare exceptions no cell partitions. Such plants are called coenocytes. They assume various forms and often resemble a small plant with stem, root and leaf, but in all these cases the plant is essentially a huge cell or tube without partitions and containing numerous nuclei. Most of the Siphonales are marine, but one genus, *Vaucheria,* is well represented in shallow streams, damp places and on the earth of flower pots in greenhouses, where it forms rather coarse green felt-like masses. The plant body consists of long tubular threads, often branching and anchored to the ground by colorless outgrowths (Fig. 113, *A*). The protoplasm forms a thick lining layer on the inner wall of the filament and embedded in it are numerous minute chloroplasts, nuclei and oil drops. A watery cell sap fills the center of the tube. Partitions only appear when reproductive bodies are formed or to close a wound in case of injury

to the filaments. Asexual reproduction is effected by very large zoöspores which are developed in the enlarged extremities of the filaments (Fig. 113, *B*). These tips are cut off by a transverse wall and a single zoöspore escapes through an opening in the tip of the sporangium thus formed. The entire surface of the zoöspore is clothed with cilia arranged in pairs, each pair being associated with a nucleus so that the zoöspore resembles the motile



Fig. 113. Structure and asexual reproduction of *Vaucheria: A,* portion of a plant showing the branching tubular filament and colorless root-like outgrowth, *r. B,* end of a filament enlarging to form a zoöspore. *C,* zoöspore. *D,* germination of a zoöspore.

colonies previously noted (Fig. 113, *C*). After a very short motile period, the zoöspore comes to rest and grows into the characteristic tubular plant (Fig. 113, *D*). When the plants are exposed to too dry conditions, the tips of the filaments often enlarge and finally become detached as motionless spores that germinate when conditions are again favorable.

The male and female gametes are produced in gametangia that are formed from short branches. The males are developed in large numbers in curved branches that become cut off from the filament by a cross wall (Fig. 114, *A*) and the gametes finally escape through an opening that forms at the apex of the branch.

A gametangium that produces clearly differentiated male gametes is called an antheridium (plu. antheridia) and the gametes are frequently called antherozoids or sperms. A single female gamete is found in a similar branch which becomes rather egg-shaped and at maturity opens at the beaked end (Fig. 114, *B*). A single-celled female gametangium is called an oögonium and the female gamete is frequently referred to as the oösphere or egg. The male gametes are discharged a few minutes after the oögonium opens, when they swarm about the open end of the female gametangium and readily enter it. As soon as fusion has been effected the gametospore becomes invested with a thick wall and in this condition can endure a limited drought. In germinating it develops directly into a new plant. This feature of the life history of *Vaucheria* is therefore of a more primitive nature than in the case of *Ulothrix*. In the following studies you will



FIG. 114. Sexual reproduction of *Vaucheria: A,* portion of a filament that has formed two branches which have grown into a male, *an,* and female, *og,* gametangia. *B,* later stage, the gametangia have opened, permitting the escape of the male gametes and the fertilization of female gamete. *C,* gametospore detached from the filament.

repeatedly notice variations of this nature. In the development of plants advances are often confined to one or another feature of their organism which may become highly developed and specialized; at the same time one or more features may be retained that have been subject to little or no variation and which, therefore, remain in a primitive state.

71. Other Members of the Green Algae.—There are several orders of the Chlorophyceae, some of which are marine, that cannot be considered at this time. Two genera, however, deserve attention because they show significant advances in the evolution of plant life. In *Oedogonium,* a member of a small order, the Oedogoniales, we find a still higher form of the sexual reproductive process. These plants are of very common occurrence in ditches, streams and springs where the filaments form for a time greenish masses and coatings upon various objects, but finally become detached and free floating (Fig. 115, *A*). Any of the cells of a filament, save the basal



FIG. 115. Growth and asexual reproduction of *Oedogonium: A,* young plant showing basal cell modified as a hold-fast. *B,* two cells of a filament, in one of which a zoöspore is forming and the other cell has opened, permitting the escape of the zoöspore shown at *C. D,* zoöspore at rest. It becomes attached to an object at its colorless, ciliated end. *E,* later stage in the germination of the zoöspore.—After West.

one, may form a single zoöspore, which are large pear-shaped bodies with a narrow colorless end around the base of which arise a circle of numerous cilia (Fig. 115, *B, C*). These zoöspores develop into new plants (Fig. 115, *D, E*) as in previous cases. The sexual reproduction presents a high degree of specialization. The male gametes are usually formed in pairs in small cells and are similar to but smaller than the zoöspores, while the female gametes are developed singly in the ordinary

cells of a filament which become greatly enlarged and spherical or ovoid in form (Fig. 116, *A, B*). The female gamete is motionless but the colorless region on one side of the gamete certainly suggests the idea that this body is related to the zoöspores and male gametes, but that it has lost its cilia. At maturity, openings of various kinds appear in the oögonium through which the male gametes enter and one fuses with the female (Fig. 116, *A,*



Fig. 116. Sexual reproduction of *Oedogonium: A,* portion of a filament in which a female gamete has been formed—*o,* opening in cell wall for entrance of male gamete. *B,* portion of filament showing formation and escape of male gametes. *C,* gametospore free from mother cell. The germination of this spore results in the rupture of its outer wall and the protrusion of the contents of the spore as four cells which are for a time retained by the delicate inner wall of the spore, as shown in *D. E,* the four cells have become mobile zoöspore and the delicate inner wall of the spore is greatly distended and about to rupture. *F, G,* stages in the germination of the zoöspore.—After Hirn.

o). The thick-walled gametospore that results from this union is set free by the decay of the oögonial walls and germinates after a resting period very much as in the case of *Ulothrix.* The contents of the gametospore is extruded through the ruptured wall as a delicate sac and then divides, forming four zoöspores which produce new plants (Fig. 116, *C–G*). The question will naturally arise as to the force that brings these and other gametes together. The frequently observed aggregation or swarming of

the male gametes about the females renders probable the view that some substances as organic acids, etc., are developed in the gametes that serve to draw them together whenever they come within a certain distance of each other.

The consideration of *Coleochaete,* a plant of the same order as *Ulothrix,* has been deferred to this point because it presents several interesting departures from previous types which indicate that a higher point has been attained in some respects by this plant than by any other of the green algae. The filaments of *Coleochaete* have a pronounced apical growth and are usually associated together in a radiate manner, forming small discs or cushion-like masses on the stems and leaves of water plants (Fig.

Fig. 117.



Fig. 118.

Fig. 117. *Coleochaete,* showing the radiating filaments and hair-like outgrowths of the cells.

Fig. 118. Sexual reproduction of *Coleochaete: A,* end of a filament bearing the male, *an,* and female, *og,* reproductive organs. *B,* the gametospore is being enveloped by the adjacent filaments of the plant. *C,* gametospore germinating. Eight cells have been formed, rupturing the walls of the gametospore.—After Oltmanns.

117). The advance of this type is indicated not only by the localization of growth at definite points, but also by the formation of the zoöspores and gametes in definite regions, *i. e.,* in special cells which are usually located at the end of the filaments. The zoöspores are produced singly from such cells and from smaller pear-shaped cells single male gametes are formed. The female gametes are developed singly in large flask-shaped cells, access to which is afforded by an opening that appears at the end of the long neck of the flask (Fig. 118, *A*). The gametospore develops a cell wall and becomes enveloped by the adjoining cells of the filaments (Fig. 118, *B*). In this condition the winter is passed and in the spring it germinates, forming neither a new plant directly as in *Spirogyra* and *Vaucheria* or zoöpores as in *Ulothrix* and *Oedogonium,* but instead, the gametospore forms a number of cells. This growth ruptures the coat of the spore and finally from each cell a rather irregular zoöspore (Fig. 118, *c*) is derived that develops into a small plant. The plants thus formed multiply solely by zoöspores until finally, after several generations, larger plants are produced that bear gametes, thus completing the life history.

72. Noteworthy Features of the Chlorophyceae.—Two features in the study of green algae should be kept clearly in mind, because they are closely connected with the tendencies that will appear in the development of the mosses. First: The gametospore is essentially a dormant or resting cell that tides the life of the plant over the conditions unfavorable for growth. Second: In passing from lower to higher types, the gametospore tends to vary in its nature and possibilities of growth. At first it forms directly a sexual or gamete-bearing plant as in *Spirogyra* and *Vaucheria.* In higher types, it develops a generation of zoöspores from which the sexual plant is derived as in *Ulothrix* and *Oedogonium.* Finally in *Coleochaete,* we find a further advance in that several non-motile cells are formed by the gametospore. These cells, to be sure, are essentially like the zoöspore because they develop directly into zoöspores, but this condition emphasizes the fact that the nature of the gametospore is steadily departing from that of the gamete-bearing plant.

This development could be summarized by saying that there is a tendency to separate the life history of the plant into two stages; a sexual or gametophyte generation and an asexual sporophyte generation. These two generations follow each other in the life history of all the higher plants. This peculiar relationship is called the alternation of generations and the reason for it will be more apparent when we come to study the mosses.

Class B. The Brown Algae or Phaeophyceae

73. General Features.—With few exceptions the Phaeophyceae are marine plants. They are commonly known as brown algae, owing to the brownish pigments which conceal in a measure the chlorophyll, thus producing their characteristic brown or yellow color. They are of common occurrence along rocky shores and attain enormous dimensions in the northern and southern seas and on the Pacific Coast. As a rule, the plant body is better differentiated than in the green algae and shows a higher type of specialization than yet seen. It would appear probable that the brown algae have been derived from green algae at a point considerably in advance of the unicellular forms. At least, there is no satisfactory evidence of a series of unicellular forms leading up to the rather complex types that are representative of the simplest of the Phaeophyceae. The majority of these lower forms have already reached the filamentous stage and many of the higher genera exhibit a differentiation of the plant body suggestive of the higher plants, as for example, a definite axis with branches and leaf-like outgrowths and root-like organs that anchor the plants to the substratum. Likewise, the tissues of these higher types often reveal many of the features already noted in the epidermis, cortex and central region of terrestrial plants. The life history of the brown algae indicates that they have undergone the same evolution as the Chlorophyceae.

(*a*) *The Simpler Brown Algae.*—This development is very clearly brought out in *Ectocarpus,* a filamentous branching form that is almost universally distributed along the sea shore

(Fig. 119). The zoöspores are developed in certain cells of the upper branches. It is important to note in the brown algae, that the reproductive cells are generally confined to special parts



FIG. 119. Reproduction of *Ectocarpus: A,* branch bearing several sporangia, *sp; g,* zoöspore enlarged. *B,* branch bearing gametangia, *gm. C,* the upper motionless gamete (female) surrounded by several still motile gametes (males). *D,* fusion of the gametes, below the resulting gametospore.

of the plant and not promiscuously developed as in the green algae. This marks a decided advance in the evolution of the

plant. The tissues are becoming more specialized and the work performed by the plant is apportioned to special cells and organs. The zoöspores have essentially the same structure and mode of growing into new plants as noted in *Ulothrix*. The cilia, however, are laterally attached and this is a characteristic feature of the zoöspores of the Phaeophyceae generally (Fig. 119, *A, g*).

The sexual reproduction of *Ectocarpus* exhibits some of the most instructive variations to be found in all the algae. The gametes are formed in gametangia that become divided into great numbers of very small cubical cells, each one of which produces a single gamete (Fig. 119, *B*). Thus the origin of the gametes is more specialized than in the case of the green algae where numerous gametes were formed in a single cell. The gametes themselves, however, show the same range of variations as previously noted. In the simpler types of this genus they are quite alike and on the same plane of differentiation as in *Ulothrix,* since they sometimes develop as zoöspores, directly forming the plant, or they may behave as gemetes and fuse to form a gametospore. The nature and character of these bodies is so imperfectly established that their behavior is entirely determined by external conditions. Low temperatures and bright light tend to develop these bodies as zoöspores, whereas, high temperatures cause them to behave as gametes. (See *Pandorina* and *Hydrodyction.*) Other species of *Ectocarpus* show an advance over this stage. The gametes are alike, but it has been observed that certain ones, the female gametes, have a shorter period of motility and after coming to rest they attract the still motile gametes, the males, and cause one to fuse with them (Fig. 119, *C, D*). This is the simplest distinction that can be pointed to as indicating a difference in the nature of the gametes which we call sex. This variation in the period of motility of the gametes must be due to an essential difference in the material or substance of which they are composed, although there is no external evidence of this. Certain species of this same genus also reveal a variation in the size of the cells from which the gametes are derived. In these species, the sex is clearly indi-

cated by the larger size of the female cell and it is noticeable that there is a decided tendency for the smaller gametes to fuse with the larger and more slowly moving females. It is also worthy of note that these gametes are less able to grow into weak plants than in the case of the similar gametes. Thus, in *Ectocarpus,* we have a most remarkable series of variations that indicate how sexuality has arisen from the asexual condition and also how sex finally became characterized by a shorter motile period in the female gamete. The character of the female finally became more pronounced, owing to its better nourishment and consequent increase in size and slower movements. A more perfect illustration of the evolution of sex is not found in nature.

The differentiation of reproductive parts, as will be frequently noted, does not keep pace necessarily with the evolution of the plant body. It is evident that sexuality has arisen independently at different levels in the plant world and in groups in no wise connected. We have noted the same state of sexuality in some of the motile green algae, *i. e.,* in *Hydrodyction* and *Ulothrix,* and now again in *Ectocarpus* although these plants are widely separated as far as relationship is concerned and exhibit a marked variation in the development of the plant body. It is also frequently to be noted that plants may get along very well indeed with only a sexual or an asexual method of reproduction.



FIG. 120. Three of the larger brown algae: *A, Laminaria. B, Lessonia. C, Macrocystis.*

(*b*) *The Coarser Brown Algae, the Kelps.*—This latter condition is well illustrated in the kelps, plants related to *Ectocarpus*, which include the largest and most highly organized forms of all the algae. Indeed some of these forms are quite comparable in size with our shrubs and trees. The Laminarias (Fig. 120, *A*) of our Atlantic coast have stalked blades ten to twenty feet long. The great bladder kelps of the Pacific, *Nereocystis* and *Macrocystis*, attain great dimensions, the latter genus reaching a length of 500 to 900 feet and *Lessonia* with trunk-like stems and leaf-like segments forms veritable submerged forests in the Antarctic Ocean (Fig. 120, *B, C*). These highly organized plants have apparently lost or failed to develop a sexual method of reproduction and depend entirely upon the formation of zoöspores for their propagation.



Fig. 121. Two common forms of the Fucaceae: *A, Sargassum*, the stem-like axis bearing air sacs, *s*, and leaf-like organs; *g*, reproductive branches. *B, Fucus—s*, air sacs; *g*, reproductive branch. *C*, young plant.

The rock weed or bladder wrack (*Fucus*) and the gulf weed (*Sargassum*) are representatives of a group, Fucaceae, that contain the most specialized of the brown algae (Fig. 121).

The *Sargassum* with its stem and leaf-like organs which may become modified into air sacs and reproductive organs bears the closest external resemblance to the higher plants of any of the algae (Fig. 121, *A*). It forms the major portion of that floating vegetation in the Atlantic known as the Sargasso Sea. The bladder wracks may be found firmly attached to the rocks by disc-like holdfasts in almost all colder, temperate and northern



FIG. 122. Structural features of *Fucus: A,* cross-section of a portion of the central stem-like part of the plant, showing an epidermal, *e,* cortical, *cr,* and central region, *c. B,* section of one of the cavities that appears to the eye as a dot. See Fig. 121, *B, g.* This cavity contains only male gametangia. *C,* section of a cavity from another plant contains only female gametangia.—After Oltmann.

seas (Fig. 121, *B*). The elongation of the flat leathery stems is largely localized in a terminal cell and results in a regular forking of the stem into two equal parts, a method of branching called dichotomy in contradistinction to the axial branching characteristic of the majority of our flowering plants. *Fucus,* like many of the gross brown algae, contains air cavities or

bladders which buoy it up in the water; this feature accounts for its popular name of bladder wrack. A cross section of the stem shows that the tissues of these plants have attained a considerable differentiation (Fig. 122, *A*) as is attested by a rudimentary epidermal, cortical and central region, the latter often containing well marked sieve tubes.

The most characteristic feature of these plants, however, and the one separating them sharply from the Laminarias appears in their method of reproduction. None of the Fucaceae develop zoöspores, although an asexual reproduction may be effected by the detachment of small branches from the plants. Sexual reproduction is a step in advance of any of the preceding types of the brown algae, in that the female gamete becomes still larger and loses entirely the power of motion. The reproductive organs, or gametangia, are developed in specialized branches or enlarged tips of the thallus. In *Fucus* (Fig. 121, *g*), these organs are contained in small pits or cavities that appear as minute points, or as dots when the enlarged tip of a branch is held up to the light. A magnified section taken through such a branch shows the nature of the cavities (Fig. 122, *B, C*). In some species, the male and female gametangia are found in the same cavity, or they may occur separately and on different plants. The male gametes are developed in enormous numbers in numerous little sacs, or antheridia, borne on branching filaments of cells which grow out from the sides of the cavities (Fig. 123). The

FIG. 123. One of the branching filaments from Fig. 122, *B*, greatly enlarged. Some of the antheridia, *an*, have discharged the male gametes, which are still retained in the inner wall of the antheridium, as at *b*. At *a* this wall is ruptured, freeing the gametes.

female gametes are produced in larger sacs or oögonia, which are each supported on a single cell and associated with hair-like chains of cells, paraphyses (Fig. 124, *A*). Usually eight gametes are developed in each oögonium. The gametes when first discharged from the antheridia and oögonia are enclosed in a delicate cell wall which later ruptures and sets them free (Fig. 124, *B*). The female gametes are without cilia and as they float away in the water they appear to attract the male



Fig. 124. Female gametangia: *A*, greatly enlarged view of one of the oögonia shown in Fig. 122, *C*. The oögonium is dividing and forming the female gametes; *p*, paraphyses. *B*, the female gametes discharged but still retained in the inner wall of the oögonium. *C*, greatly enlarged view of a gamete which is surrounded by male gametes, some of which are seen as dark bodies penetrating the cytoplasm of the female gamete. *D*, early stage in the germination of the gametospore. See later stage, Fig. 121, *C*. —After Thuret.

gametes which swarm about them and finally fertilization is effected by one of the male gametes working down to the nucleus of the female and fusing with it (Fig. 124, *C*). In an allied form not only is an attractive substance formed in the female gamete, but probably a repellent one after fertilization

is effected, for Farmer observed that the male cells swarm about the female cells for a time and then suddenly swim away "like a flock of frightened birds." As soon as fertilization has been effected, the gametospores become invested with a cell wall and attached to the rocks. Cell division now proceeds rapidly and soon establishes the characteristic thallus of the plant (Figs. 124, *D*; 121, *C*). *Fucus* is a very prolific plant and the reproduction process can be observed at almost any time. After the plants have been exposed for several hours by the low tide the male and female gametes will often be found forming orange yellow and olive green drops at the mouth of the cavities. In this condition the entire process of fecundation and early stages of germination can be studied by transferring a bit of these two fluids to a drop of sea water on a slide and studying them under a microscope. The differentiation of the gametes of *Fucus* is an interesting one because it represents the stage where the female gamete has become motionless, but is not retained in the mother cell as in *Vaucheria* and *Oedogonium*.

No resting spores are found among the brown algae. This may be connected with the more uniform conditions that obtain in the sea where they are not exposed to the dangers of desiccation as in fresh water forms. It is also noteworthy, that although the reproduction organs are, on the whole, more complex, fertilization is of a more primitive character than among many of the green algae, in that the fusion of the gametes is effected outside of the gametangia.

Class C. Red Algae or Rhodophyceae

74. General Features.—The Red Algae are largely marine plants. Unlike the brown algae, they reach their greatest development and abundance in the warmer waters of the temperate and tropical seas and are usually found attached to various objects below tidal marks. Their red pigments probably adapt them to the feeble illumination of the deep waters in which they generally occur. They range through a great variety of forms, from delicate filaments or flattened ribbon-like bodies to struc-

tures with cylindrical axes and leaf-like branches (Fig. 125). The elegant symmetry of their branching together with the deli-

FIG. 125. Common forms of the Red Algae: *A*, a leaf-like form, *Delesseria*. *B*, a branching thread-like form, *Ceramium*. *C*, a branching form covered with delicate hairs, *Dasya*.

cacy of structure and richness of coloration has always attracted attention and made them the most familiar of all the algae.

FIG. 126. A branching filament of *Callithamnion*, showing the spore mother cells forming tetraspores.

They are popularly though inaccurately known as sea mosses.

Reproduction of the Red Algae.—The reproduction of the

Rhodophyceae, particularly the sexual method, presents so many modifications and specializations that only the general features can be pointed out. Singularly the spores and gametes have lost their motility. Asexual reproduction is extensively brought about by spores that are usually formed in fours from a mother cell (Fig. 126) and therefore called tetraspores. These spore-forming mother cells arise singly on the surface or in the tissues of the plant, or they may be associated in conspicuous groups. The spores escape from the mother cell as naked bodies and finally secrete cell walls and develop into new plants. The sexual method of reproduction is the most complicated among all the algae, and even in the simplest form is far in advance of any method heretofore noted. The female gametangium resembles a flask with a long neck and is usually developed at the end of a branch (Fig. 127, *A*), see *Coleochaete*. The single female gamete is found at the base of the flask. The male gametes are produced singly in small cells also at the ends of branches and they are often closely aggregated in dense clusters (Figs. 127, *an*;



FIG. 127. Sexual reproduction in *Nemalion: A*, tip of branch bearing female gametangium, *c*, and cluster of male gametangia, *an*, from some of which the motionless male gametes are escaping. *B*, first division of the germinating gametospore, *g*. *C*, later stage showing the early formation of the branches from which the spores will be developed. *D*, spores forming at the ends of the numerous branches.

128, *A, B*). After the discharge of the male gamete, they are carried by the currents of the water to the long tube of the female organ. The wall of the tube at the point of contact is now absorbed and the male gamete passes down the tube and fuses with the female. The gametospore does not germinate and produce a new plant similar to the one that bore the gametes. On the other hand, it produces a number of branches (Fig. 127,



FIG. 128. Features in the reproduction of *Polysiphonia: A,* branch of the plant bearing clusters of male gametangia, *an. B,* one of the clusters enlarged, showing the numerous small gametangia. *C,* branch on which the spores derived from the gametospore are enveloped by a sac. From one of these bodies, called cystocarps, the spores are escaping.

B, C), the terminal cells of which develop as spores (Fig. 127, *D*). These spores on germinating produce the characteristic red algae, which bear tetraspores. The mode of sexual reproduction outlined above is complicated in the majority of the red algae, owing to the fusion of the germinating gametospore

with adjacent cells that contain storage foods. In this way, the formation of a larger number of spores is made possible. Frequently a sac-like structure, the cystocarp, is developed about the spores owing to the outgrowths of the adjacent cells (Fig. 128, *C*). The red algae are of considerable economic importance. Irish moss, *Chondrus,* is used in the manufacture of jellies, and agar-agar is obtained from several species of algae. Many tons of various kinds of red algae are annually dried and consumed for food in the East. The swallow's nest, of which you have heard so much as an article of food in the Orient, is constructed of algae.

75. Significant Features in Life History of the Algae.—The germination of the gametospore in the Rhodophyceae is a noteworthy departure in two respects. Unlike previous cases, it is retained on the plant, where it is nourished during its germination and grows practically as a parasite. This relation of the gametospore to the mother plant will become more noticeable among the mosses and lead to pronounced changes in the life history of the plant. In the second place, we notice that the gametospore in the red algae produces a number of cells before the spores are formed. In the green and brown algae it either developed directly into a plant similar to the mother plant, as in *Spirogyra, Vaucheria* and *Fucus,* or zoöspores are first formed which developed into a plant similar to the parent, as in *Ulothrix* and *Oedogonium.* What is the significance of this variation in the germination of the gametospore? Attention has been directed in the study of *Ulothrix*, p. 173, to the fact that there are two phases in the life history of a plant, a spore-bearing phase or generation, and a sexual phase or generation. This is doubtless true of all forms characterized by a sexual reproduction. On page 122 you have noticed that these two phases or generations are also sharply distinguishable by the number of chromosomes appearing in the nuclei of the cells, the sexual generation having only one half the number found in the asexual. This change in the number of chromosomes occurs at two points in the life history, first, when the gametes unite to form the gametospore, at which time the

number of chromosomes is doubled, and, secondly, when the spore mother cell of the asexual generation divides to form the spores, the number of chromosomes is reduced by one half. The two generations are generally not apparent in the Green and Brown Algae, because the asexual generation is greatly reduced and often does not produce spores, as in *Vaucheria, Spirogyra,* etc., where the asexual generation consists of a single cell, the gametospore. This gametospore, however, must have twice the number of chromosomes as the mother plant, since it is formed by the union of the chromatic substance of two gametes and it will doubtless be found that its first divisions in germination are attended with a reduction of the chromosomes to the original number found in the mother plant. This remark applies also to *Ulothrix* and *Oedogonium,* where it is reasonable to suppose that the formation of the four zoöspores by the germination of the gametospore results in the reduction of the chromosomes to the number found in the mother plant. This has been found to be the case by Allen in the division of the germinating gametospore of *Coleochaete,* and Yamanouchi reports that the reduction of the chromosomes in the Red Algae does not occur until the mother cell divides to form the tetraspores (Fig. 126). For example, in *Polysiphonia* the gametospore germinates, producing a small spore-bearing plant, enclosed in the cystocarp, which is parasitic upon the sexual plant. This plant is a part of the asexual generation, as all its cells contain the double number of chromosomes. The spores from the cystocarp form plants externally similar to the sexual plants, but these plants are really different, being characterized by cells with the double number of chromosomes and also by the fact that they bear tetraspores. The formation of the tetraspores, however, is attended with a reduction of the chromosomes by one half, and these spores produce plants characterized by the reduced number of chromosomes and also by the fact that they bear the sexual organs. So in the Red Algae we have very clearly brought before us the two generations in the life of the plant, the sexual generation, characterized by the reduced number of chromosomes and the production of the sexual organs, and the asexual generation, characterized by the double

number of chromosomes which appear in two-spore bearing plants, *i. e.*, the minute plant in the cystocarp and the tetraspore-bearing plant. The distinction between the sexual and asexual generation is emphasized at this point because we see in the algae how it gradually became more and more conspicuous, and in the mosses and succeeding groups it will be noticed that the relation of these two generations is intimately associated with the evolution of the higher types of plant life.

Subdivision 5. Eumycetes or Fungi

76. The Nature of Fungi.—The Fungi are the largest group of the Thallophyta and include such familiar forms as moulds, mildews, toadstools and mushrooms. The absence of chlorophyll is the most striking feature of these plants. They are unable therefore to form sugar, starch and other foods from the elements of the soil and air and must obtain them already manufactured. Consequently, they are either saprophytes living upon decaying organic matter, or parasites preying upon living organisms. The majority of fungi procure their food from decaying plant and animal matter, and in this relation many of them are of the same economic importance as the saprophytic bacteria. Other forms exist as parasites upon living plants and animals. In this relation fungi cause almost incalculable loss annually to the country and frequently crops over large areas are ruined by their depredations. The parasitic forms gain access to the host through a wound, stoma or the delicate tissues of seedlings or young parts of the plant, and in other cases the fungus forms a ferment that dissolves the cell walls and thus opens the way for the entrance of the pest. The cultivation of plants has doubtless weakened their resistance in many cases to the attack of the parasites, and like ourselves the plant may inherit a weaker constitution that is more subject to disease. Every state now employs experts to study the diseases caused by these fungi and to devise means for killing them and to develop more resistant plants.

The fungi are of very simple constructure because they live upon foods already manufactured for them. There is no longer

a necessity for a plant body that is complex, because each part is adapted to the performance of one or another of the many functions that coöperate in the construction and distribution of the organic substances. Consequently, the fungi do not exhibit many of the characteristics of chlorophyll-bearing plants. Their cell walls are thin and inclose a watery, colorless protoplasm in which are usually dispersed many small nuclei (Fig. 129). Whatever



FIG. 129. A few cells from a branching filament of green mould, *Penicillium*, showing the granular character of the cytoplasm and the absence of platids. The colorless areas in the cells, vacuoles, contain principally water.

form the plant body may assume, it will usually be found to consist of filaments of delicate cells or tubular growths, called hyphae (sing. hypha). These fine filaments or tubes are the essential portion of any fungus and as they spread over the substance upon which they feed they form a branching and interwoven mass of threads collectively known as the mycelium. This structure is well illustrated in the hyphae that spread over bread and fruits, forming a cobwebby mass or mycelium that is commonly known as mould or mildew. Even in the larger fungi, as the mushroom, it will be found that the entire body is composed of a mass of these delicate hyphae which spring from a mycelium in the ground. We will expect to find remarkable departures in the fungi from previous types because of their peculiar manner of obtaining food and also because they are largely terrestrial plants and therefore exposed to a much wider series of conditions than in the case of the algae. This will tend to cause variations and so bring about new structures that are in harmony with the conditions under which the plants live. As a rule, these variations have been so extensive and have so completely changed the char-

acter of the plant as to render it impossible to state whence many of the groups have been derived or what relationship they sustain to one another. It may be stated, however, that the lower fungi show unmistakable evidence of relationship with certain of the green algae and that they have become degenerated owing to their parasitic habit and consequent loss of chlorophyll. The fungi may be divided into three classes: A, Phycomycetes or Alga-like Fungi; B, Ascomycetes or Sac Fungi; C, Basidiomycetes or Basidia-bearing Fungi.

Class A. Phycomycetes

77. Alga-like Fungi.—Some of these plants show such a striking resemblance to certain algae, both in the structure of the plant body and in their reproductive processes, that they are called the Phycomycetes from phycos, alga, and myces, mould. Attention will be called to three orders of this group.

78. Order a. Saprolegniales or Water Moulds.—These fungi are aquatic, living upon dead animals and plants, and as parasites they are very destructive to fish. Common examples of this order are seen in the whitish masses that form around decaying insects in the water and in fluffy or mould-like outgrowths that often appear upon the bodies of fish in aquaria. The plant body consists of branching tubular threads without partitions but containing numerous nuclei and thus resembling *Vaucheria* save for the absence of chloroplasts (Fig. 130). The sporangia are also formed by the cutting off of the tip of one of the branches by a transverse wall. The contents of a sporangium, however, generally breaks up into a very large number of biciliate zoöspores (Fig. 130, *C*). In the species that cause so much damage to fish, the spores come to rest upon the fish and form tubular outgrowths that readily penetrate the tissues of the fish, especially where a scale has been rubbed off. Some of the hyphae also extend outward from the body, causing the mould-like blotches on the infected fish, and from the tips of these branches the sporangia mentioned above are formed. Most of the species of *Saprolegnia* also form sexual organs as in *Vaucheria*, from one

to several gametes being formed in an oögonium (Fig. 130, *D*). The antheridia, however, develop several non-motile male gametes. This peculiarity of the gametes is perhaps due to their exposure to atmospheric conditions, as would be the case when growing upon floating organic matter. The lack of water for the transport of the male gametes is nicely met by a tubular out-



FIG. 130. Features in the life history of the water moulds: *A*, fly infected with *Saprolegnia*, showing the tubular threads of the fungus radiating out from its body. *B*, tip of a tube magnified, showing an early stage in the formation of the sporangium. *C*, sporangium discharging zoöspores. *D*, Reproductive organs of a related fungus, *Achlya*—*o*, oögonium containing four female gametes; *an*, antheridium with tubes penetrating oögonium for transport of male gametes.

growth of the antheridium which penetrates the oögonium when it ruptures, discharging the males in the immediate vicinity of the female gametes (Fig. 130, *D*, *an*). In the majority of the species, singularly enough, the female gametes germinate without being fertilized. The gametospore germinates as in *Vaucheria*. developing readily into new plant.

It should be stated that, in related forms, large multiciliate zoöspores, like those of *Vaucheria*, are formed and also motile male gametes. It is also important to note that there are several

parasitic algae closely related to these alga-like fungi. Their parasitic habit and partial loss of chlorophyll clearly indicate that they are in a transition state from the algae to the fungi. It is therefore safe to state that in this group we have evidence of the derivation of the fungi from the algae.

79. Order b. Peronosporales. Downy Mildews and White Rusts.—These forms are parasites upon the higher plants especially and include two of the most destructive fungi, the potato and grape vine blight. *Phytophthora infestans* causes the potato rot. This disease was widespread in the eastern United States in 1901, causing the entire loss of the crop in some sections. It would be difficult to imagine a pest more perfectly adapted to a destructive career. The tubular filaments of the fungus readily spread through the tissues of the leaf, sending into the cells short lateral branches called haustoria (Fig. 131), which absorb the cell con-



FIG. 131. Hyphae of *Peronospora* extending through the tissues of a plant and absorbing food from the cells by means of haustoria, *h*.

tents and thus supply the fungus with food. In severe cases this produces a withering and decay of the leaf. The hyphae continue their growth into the stems and all parts of the plant, thus causing the black discoloration of the potato and its early decay in bad cases of infection. When the mycelium has become well established in the leaf, numerous branching hyphae extend out through the stomata and form at their tips little sacs or sporangia (Fig. 132, *A*). The formation of the sporangia is effected in a few hours, when they drop off and are carried by the wind to other plants, where they germinate at once, forming a tube that penetrates the leaf and rapidly spreads the disease. If the sporangia chance to fall upon leaves that are wet by dew or rain, the

contents breaks up into several zoöspores (Fig. 132, *B–E*) which finally come to rest and develop the characteristic tubular hyphae of the fungi. This behavior of the sporangium is doubtless a survival of the zoöspore stage seen in the algae. It is equally suggestive that definite changes in the environment, as the dry air, causes the sporangium to germinate as a non-motile spore, whereas the presence of water causes the sporangium to produce several zoöspores.



Fig. 132.

Fig. 133.

Fig. 132. Asexual reproduction of the mildew: *A*, hyphae of *Plasmopara* emerging from a stoma and bearing numerous sporangia.. *B*, enlarged view of sporangium of *Peronospora* germinating on a dry leaf. In this case the sporangium behaves as a spore sending out a hypha that will penetrate the tissues of the leaf. *C*, sporangium of *Phytophthora* germinating in the water and forming zoöspores. *D*, zoöspore enlarged. *E*, zoöspore has come to rest and is forming a tube that will penetrate the tissues of the leaf as in the case of *B*.

Fig. 133. Reproduction of the white rust, *Albugo: A*, asexual stage, showing several erect hyphae forming spores. *B*, sexual stage which is characteristic of the Peronosporales in general—*o*, gametangium containing a single female gamete which is being penetrated by tube from male gametangium, *an*.—After Wager.

Very similar in character is the grape blight, *Plasmopara viticola*. This disease causes a browning of the leaves and the stunting of the fruit, which become brown or gray, and, finally, the death of the infected parts. This pest caused the abandonment of entire vineyards before the method of killing the fungus by spraying the plants with copper salts was discovered. In both of these pests and in allied genera the sporangia-bearing hyphae are produced in such numbers as to cause a downy, mould-like appearance on the leaves, thus accounting for their popular name, Downy Mildews.

In a related genus, *Albugo*, the sporangia are formed by the repeated cutting off of the tips of the hyphae, as shown in Fig. 133, *A*. In this way, not one, but a chain of sporangia, are formed from the end of the hyphae which do not project from the leaf, but grow up in dense masses just under the epidermis, producing glistening white blotches or blisters on the leaves. This growth finally ruptures the epidermis when the spores are scattered by the wind and germinate as in the preceding cases. This fungus, known as white rust, is very common on mustards, pigweed and other plants.

The sexual reproduction of the Peronosporales is suggestive of *Vaucheria*. Gametangia are cut off from the ends of the hyphae, as shown in Fig. 133, *B*. The male gamete gains access to the female gamete, which are usually formed singly, by means of a tube as in *Saprolegnia* (Fig. 133, *B, an*). The thick-walled gametospore, as in many of the algae, tides the plant over the winter, and being set free by the decay of the surrounding tissues it germinates in the spring, starting anew the life of the pest. It may germinate directly (see *Vaucheria*) into the fungus or zoöspores are first produced, as in *Oedogonium*.

These two orders are more suggestive of the algae than any others that we will study and it is well to note the modifications that have been induced in these plants as a result of their change from aquatic to terrestrial conditions. Removed from the water, special root-like organs and haustoria are evolved for the absorption of foods. The absence of water brings about a lack of motility in the male gametes and the formation of a tube to con-

duct them to the female. For the same reason, the zoöspores are reduced to light motionless spores that are developed upon elongated hyphae that expose them to the air currents for distribution. The ability of these spores, or sporangia, to produce zoöspores is doubtless the survival of a trait inherited from their algal ancestors.

80. Order c. Mucorales or Black Moulds.—These are among the most common of the fungi and they are almost sure to appear upon any cooked food or decaying matter that is exposed to the air even for a very short time (Fig. 134). The mycelium has

FIG. 134. Habit of growth of the black mould, *Rhizopus*: *s*, sporangia; *r*, absorbing branches of the mycelium.

practically the same structure as noted in the preceding groups, but the black moulds have lost all motile reproductive bodies and their relationship to any group of the algae is not known. Some of the hyphae of the mycelium creep over the food supply and send into it short branches which serve as organs of absorption, while other rather thicker hyphae grow away from the mycelium and reach up into the air (Fig. 134). The tips of these erect hyphae are cut off by a transverse wall and enlarge into spherical sporangia (Fig. 135). The protoplasmic contents of the sporangia divides into a large number of small bodies or spores, each of which contains a nucleus and becomes covered with a rather thick smoky black wall. The numerous sporangia filled with dark spores are the principal cause of the black color of these fungi. While the sporangium is forming, the transverse wall which separates it from the hypha, becomes convex, owing to the

pressure of the fluid in the hypha, and finally forms a dome-like structure in the sporangium (Fig. 135, *D–E*). The walls of the



FIG. 135. *A, B,* early stages in the development of the sporangium of *Rhizopus*. *C,* the formation of the cross-wall cutting off the sporangium from the hypha. *D,* this cross-wall is being pushed up into the enlarging cavity of the sporangium and in *E* it has become greatly extended, forming a sac-like structure, known as the columella. *F,* the sporangium has ruptured, permitting the scattering of the spores and the columella has formed an umbrella-like structure owing to the loss of its watery contents.



FIG. 136. Sexual reproduction in the black mould: *A,* the meeting of the tips of two hyphae. *B,* later stage, the lower part of the figure shows the cutting off of the tips by transverse walls and in the upper part of the figure the fusion of the contents of the two gametangia thus formed has begun. *C,* mature gametospore.

sporangia readily dissolve in the presence of moisture, owing to their mucilaginous character, and thus allow the spores to float off in the air as an invisible dust. The spores will germinate at

once and produce a new plant under suitable conditions of moisture and food, or they will retain their vitality for months if kept dry. As the spores are disseminated, the dome-like structures in the sporangium reflexes, owing to the loss of its watery contents and assumes an umbrella shape, as seen in Fig. 135, *F*. You can readily understand why these black moulds are so common by counting the number of sporangia on a small bit of mycelium and then estimating the number of spores in a sporangium. So numerous and light are the spores that they are carried everywhere. It is only necessary to expose a bit of moist bread for a few moments to the air and then enclose it in a damp chamber to secure a luxuriant crop of these plants. Under certain conditions the sexual method of reproduction is effected by the union of two club-shaped hyphae, as shown in Fig. 136, *A*. As these hyphae meet, the tip of each branch is cut off by a wall and the



FIG. 137. A fungus, *Empusa*, parasitic upon flies: *A*, fly surrounded by a mass of discharged sporangia. *B*, enlarged view of several hyphae, showing the discharge of the sporangia which are surrounded by a mucilaginous substance.—After Brefeld.

contents of the two tips fuse, forming a thick-walled gametospore (Fig. 136, *B, C*). After a resting period, this gametospore grows directly into a new mycelium. In some of the black moulds reproduction is only effected by the union of hyphae from different plants which must differ therefore in their nature, just as you found to be the case in some species of *Spirogyra.*

81. A Fly Fungus.—An interesting form allied to the mucors is seen in the parasitic fungus *Empusa,* that produces an epidemic among flies at certain seasons when they may be seen dead and clinging to the woodwork and window panes surrounded by a white halo (Fig. 137, *A*). This appearance is caused by the discharge of numerous white sporangia that are found at the ends of the hyphae that project from all sides of the fly's body (Fig. 137, *B*). If, by chance, a fly should come within range of these discharging sporangia, it would become infested and later, branches of the hyphae would project from its body and repeat the process of sporangia formation.

Class B. Ascomycetes or Sac Fungi

82. General Characters.—The Ascomycetes are the largest and most variable group of the fungi. They are illustrated by the powdery mildews which affect the leaves of a variety of plants; the brown and blue moulds that occur on preserves, decaying fruit, old leather, etc. (Fig. 138); the cup fungi and morels (Figs. 156, 158); the black knot of the cherry and plum trees (Fig. 150), and the ergot of rye, etc.

The mycelium resembles in structure that of preceding orders, but its hyphae are composed of numerous cells instead of being tubular as in the Phycomycetes (Fig. 138). The sporangia (Fig. 142) are reduced in size and contain but a single spore each, which is never discharged from the sporangium. This body will be referred to in the future as a spore or conidium (plu. conidia).

The gametangia are often suggestive of those of the downy mildews (Figs. 147, *A*; 157), though in many forms they appear to be lacking. The fusion of the gametes results in the formation of a gametospore which germinates at once, forming in the

simplest cases one or more sacs. These bodies are known as asci (sing. ascus) and they contain more frequently eight spores, called ascospores (Fig. 141), which must be distinguished from the spores or conidia mentioned above. The gametospore germi-

FIG. 138. The green mould, *Penicillium,* one of the most common of the Sac Fungi. The hyphae of the branching mycelium is composed of cells and the spores or conidia are formed in chains that are arranged in brush-like clusters at the ends of the erect hyphae.

nates as a parasite on the mother plant as in the red algae. The sexual process results, in the majority of forms, not only in the formation of asci, but numerous hyphae from the mycelium are also stimulated to growth and become associated with the asci in various ways. As a result bodies of various forms, called ascocarps or perithecia, are developed which often become the conspicuous part of the fungus (Figs. 143, 144). The ascospores germinate immediately or after a period of rest and form a new mycelium. It should be stated that the ascocarps have not been connected, in many cases, with sexual processes, and it is inferred that sexuality has been lost along with other characters as a consequence of the degeneracy resulting from parasitic and saprophytic habits. Only a few of the more important orders will be discussed.

83. Order a. Yeast or Saccharomycetes.—These fungi are unicellular plants and many regard them as reduced ascomycetes. They do not produce a mycelium, but consist of rather oval cells which multiply by a budding process shown in Fig. 139, *A, B.*

The new cells that push out from the side of the mother cell readily drop off, but in rapid growth they may remain attached in chains (Fig. 139, *D*). Under certain conditions, as the ex-



FIG. 139. The yeast plant, Saccharomyces: *A*, single plant. *B*, plant producing three buds. *C*, section of two plants showing buds and nuclear division. *D*, chain of plants due to rapid budding and growth. *E*, formation of ascospores. *F*, germination of an ascospore and the formation of new plants by budding.—After Wager.

haustion of the food supply, the cells become transformed into asci and the contents of each cell rounds off into one or more ascospores (Fig. 139, *E*). The ascospores are freed by the decay of the ascus and when conditions are favorable, grow into the characteristic yeast cells, as shown in Fig. 139, *F*.

(*a*) *Fermentation.*—These microscopic plants must be numbered among those plants that are of the greatest economic value. Their importance is due to the fact that they decompose sugars upon which they feed into CO_2 and alcohol, a change called fermentation. The extensive brewing and distilling industries all over the world are dependent upon the growth and peculiar action of these microscopic plants. When yeast plants are placed in solutions containing sugar in the form of molasses or prepara-

tions of rye, corn, barley, potatoes, etc., and slightly warmed, the growth of the yeast produces a vigorous fermentation. Carbon dioxide rises to the surface, forming a frothy scum while the alcohol accumulates in the fluid. Beers and ales are fermented beverages of this nature, while whiskies, brandies, alcohol, etc., are obtained from the fermented mass by removing a part of the water by distillation. Wines and cider are weak alcoholic beverages formed by the yeast plant from the sugars in the juices expressed from grapes and apples. In this case, the yeast is generally allowed to find its way naturally into the fluids. So minute are these plants that they are distributed in the air throughout the world and no weak sugar solution can be exposed for any length of time without being inoculated by them—a fact that has been apparently taken advantage of by man in all times and places. When these fermented beverages are bottled before the decomposition of the sugar is complete, then a further generation of CO_2 sets up a pressure in the bottle or cask that causes the popping of the cork when the bottle is opened and the sparkle of the fluid owing to the escaping gas.

(*b*) *Bread Making.*—The most important use of the yeast plant is its application to the "raising" of bread. Flour contains, in addition to starch, a little sugar, and this amount is increased in bread making by a ferment, diastase, in the flour which changes starch to sugar. The flour is mixed with water containing yeast plants into a dough and placed in a warm place, when it begins to rise. This means that the yeast plants begin to grow and decompose the sugar into CO_2 and alcohol. The dough prevents the escape of the gas which collects in bubbles in the dough, causing it to swell. Baking further expands the gas and also drives off the water and alcohol, leaving the bread light and porous. There are many forms of yeast plants which differ in their power of producing fermentations and in the flavor which they impart. Consequently, different forms are used for different purposes. Bread yeast is a form that has been selected because of the quickness of its action and the flavor that it imparts to bread. Bread yeast is grown in large vats and put up with starch in cakes, known as compressed yeast. This product is not pure and

will doubtless be greatly improved. It contains bacteria, and if not fresh will seriously interfere with the quality of the bread.

84. Order b. Exoascales or Peach Curl.—This group includes a small number of parasitic fungi that are especially destructive to peach and plum trees, causing distortion of the leaves and fruit, known as leaf curl and bladder plum (Fig. 140). Numer-

FIG. 140.

FIG. 141.

FIG. 140. A branch from peach tree, showing the distortion of the leaves caused by the fungus, *Exoascus.*

FIG. 141. Section of a leaf showing numerous cells rupturing the cuticle and developing into asci, *as.*

ous asci are developed from the mycelium beneath the epidermis, which is finally ruptured by their growth (Fig. 141). The ascospores are discharged into the air by the bursting of the asci and carried by the wind to other plants. The damage in the United States to peach trees alone is estimated at $2,000,000 to $3,000,000 annually.

85. Order c. Aspergillales or Blue-green and Brown Fungi.—This group includes perhaps the most widely distributed and familiar examples of the fungi. They occur as blue-green or brown moulds upon almost any organic matter, forming a delicate mycelium from which are developed numerous erect hyphae. In the common blue mould, *Penicillium,* these erect hyphae are broom like (Fig. 142, *B*) and the spores are formed from the tips of the branches very much after the manner of the budding of the yeast cells. The tip of a branch buds out into a spherical cell that is finally cut off from the stalk, thus forming the spore

(Fig. 142, *C*). This process is repeated again and again just below each successive spore and in this way chains of spores are formed. It will be noticed that many single spores are found instead of a large sporangium which contains many spores as



FIG. 142. Formation of spores or conidia in the Aspergillales: *A*, spore-bearing hypha of *Aspergillus*. *B*, hypha of green mould, *Penicillium*. *C*, one of the terminal branches of *B* enlarged, showing manner of spore formation.

in the case of *Mucor*. These spores are sometimes regarded as sporangia which have become reduced in size and contain but a single spore. In *Aspergillus*, a fungus common upon preserves and upon herbarium plants that have not been sufficiently dried, the spores are developed from very short branches that arise from a bulbous swelling at the apex of the erect hyphae (Fig. 142, *A*). The color of the spores, as in *Mucor*, is the cause of the characteristic blue or brown color of the fungi. These extremely small and numerous spores float off in the air as an invisible dust and quickly germinate under favorable conditions, forming new mycelia.

The male and female gametangia are very similar, short branches that become closely intertwined (Fig. 143, *A*). The fusion

of the gametes is made possible through a dissolution of the separating walls at the tips of the gametangia. The gametospore in germinating, develops an irregular hyphal outgrowth



FIG. 143. Sexual reproduction of the Aspergillales and the formation of the ascocarp or perithecium: *A*, meeting of the male, *an*, and female, *o*, gametangia. *B*, early stage in the development of the ascocarp. The gametospore has formed a series of branches, *s*, which are being surrounded by hyphae from the mycelium. *C*, later stage seen in section. The gametospore has formed a much-branched body, *s*, which is surrounded by a closely interwoven mass of hyphae that appear in section as cells.—After Brefeld.

(Fig. 143, *B*, *s*) which becomes completely overgrown by the hyphae of the mycelium (Fig. 143, *B*, *C*), thus forming a solid body, known as the ascocarp or perithecium (plu. perithecia), that appears to the eye as a minute grain of sand. During this growth numerous lateral branches arise on the hyphae derived from the gametospore and become transformed into asci as shown in Fig. 144, *A*, *B*. The ascocarps finally decay and set free the ascospores which develop a new plant or mycelium. Thus the entire sexual process, with the exception that the male gamete is not set free, is suggestive of the Rhodophyceae.

The truffles are a curious group of related fungi that live for the most part entirely under ground. The mycelium of many forms is supposed to live in contact with the roots of oaks and other trees as a mycorhiza. The fleshy tuber-like ascocarp, often as large as a walnut, is a highly prized delicacy in Europe where

dogs and pigs are trained to locate the truffles by smell. This industry amounts to more than f1,000,000 annually in France and Italy.



FIG. 144. Further development of the ascocarp: *A*, sectional view, showing the branches, *s*, derived from the germinating gametospore, that are forming numerous lateral branchlets. *B*, one of the branchlets enlarged, showing how it divides into cells which round off, forming the asci, *as*. *C*, ascospore. *D*, germinating ascospore.—After Brefeld.

86. Order d. Perisporiales or Powdery Mildews.—This name is given to a common and widely distributed group of largely parasitic fungi that form cobwebby mycelia on the under surface of the leaves of the elm, maple, willow, lilac, rhododendron, Virginia creeper, etc. (Fig. 145), and a great variety of herbaceous plants, as the dandelion and cocklebur. The majority of them do not seriously interfere with the health of the plant, but others produce serious diseases in young cherry and plum trees, hop vines, gooseberry, etc. These parasites are external and obtain their food by means of short branches, haustoria, which dissolve the cell wall and absorb the cell contents as shown in Fig. 146, *h*. The spores are formed in chains (Fig. 146, *c*) from the end of the erect hyphae that project from the surface of the leaf in thick masses, causing the powdery appearance and the popular name of these parasites. These spores germinate quickly and rapidly spread the fungus.

The reproductive organs are formed from short branches as shown in Fig. 147, *A*. The solution of the walls at the point of

Fig. 145.



Fig. 146.

Fig. 145. Appearance of one of the powdery mildews, *Uncinula,* on leaf of elm.

Fig. 146. Enlarged view of the mycelium, ascocarp, etc., of one of the mildews, *Erysiphe: c,* erect hyphae forming spores or conidia; *h,* haustoria penetrating epidermis of leaf; *a,* ascocarp or perithecium.



Fig. 147. Sexual reproduction of a powdery mildew: *A,* meeting of the male, *an,* and female, *o,* gametangia. *B,* fertilization, the male gamete, *m,* is seen approaching the female. *C,* section of young ascocarp showing the early germination of the gametospore, which has become surrounded by hyphae derived from the mycelium. *D,* later stage, the gametospore has developed several cells and the terminal one is being transformed into an ascus, *as.*—After Harper.

contact of these organs permits the male gamete to pass over and fuse with the female (Fig. 147, *B*). The growth of the gametospore forms a limited number of cells, one or more of which develop into asci (Fig. 147, *C, D*). As in *Penicillium,* this growth becomes enveloped by a mass of hyphae that originate from the mycelium. These ascocarps appear at maturity as black specks and in the majority of forms they are provided with hair-like outgrowths that are very regular and characteristic of the genera (Fig. 148). The ends of these hairs are rather



FIG. 148. Forms of ascocarps found among the powdery mildews: *A, Phyllactinia* with needle-like appendages enlarged at the base. *B, Microsphaera,* appendages dichotomous at apex. *C, Uncinula,* appendages coiled at apex. *D, Erysiphe* without appendages and crushed to show escaping asci. *E,* an ascus containing six ascospores.

mucilaginous and may assist in the dissemination of the ascocarps. The ascospores are resting spores adapted to enduring drought and cold as in *Penicillium,* which they resemble in their discharge and germination.

87. Forms with Slightly Opened Ascocarps.—Several orders of the Ascomycetes are distinguished from the preceding group by the formation of ascocarps that are slightly open (Figs. 151; 152, *B*). In this way, provision is made for the discharge of the ascospores without the decay of the ascocarp. Only two of these orders can be noticed.

88. Order e. Sphaeriales or Black Fungi.—This is the largest group of the ascomycetes, over 2,000 species being known in the United States alone. Scarcely a fallen twig or bit of old wood



Fig. 149.

Fig. 150.



Fig. 151.

Fig. 149. A common black fungus, *Hypoxylon: A,* habit of the fungus as it appears on dead branches and logs. The round black bodies are an association of the mycelium, stroma, and numerous ascocarps. *B,* a single ascus enlarged, showing character of the ascospores.

Fig. 150. The black knot, *Plowrightia,* infecting a branch of cherry. At the bottom of the branch is shown the early summer or spore-bearing stage, *c,* and above a black warty mass of ascocarps, *as,* produced the previous season.

Fig. 151. *A,* several ascocarps enlarged, taken from region *c* in Fig. 150. *B,* diagram of an ascocarp as seen in section, showing the asci and the opening for the escape of the ascospores.

can be examined without revealing the minute hard black ascocarps (Figs. 149; 152, *A*). Many of these fungi are quite conspicuous since the ascocarps are found in large compact masses and also because they are often associated with a more or less conspicuous outgrowth from the mycelium called the stroma (Figs. 150; 152, *D*). The majority of the genera are saprophytic upon dead and decaying vegetation, though some of them are destructive parasites. The black knot, *Plowrightia,* the cause of a serious disease to plum and cherry trees, illustrates very well the characteristics of this order. The mycelium grows in the

cambium and cortical regions of the branches, causing the bark to split open in the spring when spore bearing hyphae extend up into the air forming velvety coating (Fig. 150, *c*). By the approach of winter, this mycelium has grown into the familiar black knotty mass in which are developed numerous ascocarp-like bodies (Fig. 150, *as*). The spores from these ascocarps are carried by the wind in the early spring to other branches and probably infect the budding trees. Other conspicuous forms are *Xylaria* and *Daldinia,* which develop an extensive stroma on stumps and trees that contains numerous ascocarps (Fig. 152).



FIG. 152. Other common forms of the Spheriales: *A,* habit of *Hysteriographium* on a dead twig. *B,* ascocarps enlarged. *C,* ascus enlarged, showing character of ascospores. *D, Daldinia. E,* section of the same, showing that the stroma forms a concentric stratum of ascocarps, *as,* each year. *F, Xylaria. G,* the same with branch cut off to show the layer of ascocarps on the periphery of the stroma.

In *Hypoxylon,* the stroma containing the ascocarps breaks through the bark of a large variety of trees and shrubs in the form of spherical or cake-like masses (Fig. 149).

89. Order f. Hypocreales.—These fungi are distinguished from the Sphaeriales by their rather fleshy or membranous ascocarps and stroma, which range in color from white to yellow,

purple, scarlet and brown. Numerous species are saprophytic while others are parasitic upon higher plants, fungi and insects. *Claviceps* is a common example of this group causing the disease

FIG. 153. FIG. 154.

FIG. 153. Head of rye infested by the parasite, *Claviceps,* which has transformed several of the grains into black masses of mycelium known as sclerotia.

FIG. 154. Various phases in the life history of *Claviceps: A,* a young grain or pistil infested with the parasite. *B,* enlarged view of the mycelium as it appears on the surface of the pistil, showing the formation of numerous spores. *C,* the hyphae of a sclerotium growing out and forming several purplish stalks, capped with knob-like clusters of ascocarps. *D,* enlarged sectional view of one of these knobs, showing numerous ascocarps on the periphery. *E,* one of the ascocarps enlarged, *as,* asci. *F,* an ascus containing eight thread-like ascospores.

known as ergot in the flowers of rye and other grasses (Fig. 153). The mycelium at first spreads over the outer part of the pistil, rapidly forming spores (Fig. 154, *A, B*) and exuding a sweet slimy juice, honey dew, which is eagerly eaten by flies. In this way the spores are carried away to infest other plants. The mycelium finally completely absorbs the substance of the grain and grows into a hard blue-black body several times larger than the grain (Fig. 153, *sc*). This body, known as the sclerotium, remains dormant during the winter and in the spring develops a stroma that assumes the form of rose-colored stalks bearing numerous ascocarps arranged in globular heads (Fig. 154, *C, D*). The ascospores germinate in the spring and infest the flowers of the grain. The Chinese wonder, *Cordyceps,* is a related parasite that attacks caterpillars, larvae of beetles and truffles. In the case of the forms living upon insects, the parasite does not usually appear until the spring, when they are in the pupa or

Fig. 155.

Fig. 156.

Fig. 155. The stroma of *Cordyceps* emerging from the pupa of a moth and forming a club-like organ with numerous ascocarps, *as,* in its apical region.

Fig. 156. One of the common cup fungi, *Peziza,* with broadly open ascocarps. Common upon rich humus soil and decaying wood.

cocoon stage. At this time, the mycelium which flourishes in the tissues of the host, sends up club-like bodies (Fig. 155), that bear the ascocarps as in the case of the ergot.

90. Forms with Broadly Opened Ascocarps.—The remaining orders of the Ascomycetes include genera in which the ascocarps

are broadly open and often associated with large masses of mycelium, thus forming a conspicuous stroma.

91. Order g. Pezizales or the Cup Fungi.—These plants are characterized by the formation of fleshy, leathery or gelatinous cup-like ascocarps that range in size from mere specks to forms four or five inches in diameter (Fig. 156). The mycelium lives upon the humus in the ground or on decaying plants and apparently in this and the next order frequently develops the ascocarps directly without a reproductive process. In *Pyronema*, however, Harper has shown that reproductive organs are formed that are more suggestive of those of the red algae than was the case in the blue moulds and downy mildews previously noted. The female gametangium has a long tubular outgrowth that curves over and becomes closely applied to the male (Fig. 157,



FIG. 157. Sexual reproduction of one of the cup fungi, *Pyronema: A*, the flask-shaped female gametangium, *o*, is seen in various stages of fusion with the male gametangium, *an*. *B*, the gametospores, *g*, germinating and forming branching hyphae which bear the asci, *as*. These asci are associated with hyphae or paraphyses, *p*, that arise from the mycelium. The asci and paraphyses constitute the hymenium.—After Harper.

A). The dissolution of the separating walls permits the fusion of the gametes, after which, the gametospore develops a number of erect sac-like hyphae or asci (Fig. 157, *B*). Adjoining hyphae of the mycelium grow up among the asci and around them,

forming a cup-like structure resembling that shown in Fig. 156. A section through one of these cups reveals the asci intermingled with hyphae, also called paraphyses, in the form of a layer or stratum. Such an association of spore-bearing organs and paraphyses is called a hymenium. The spores are discharged to considerable distances, owing to the accumulation of fluids in the asci which finally rupture at their apices. In some forms, the entire ascus is discharged with an audible explosion, owing to the swelling and rupture of the supporting hyphae. The bright scarlet cups of *Sarcoscypha,* common in the spring on decaying sticks and the gray cups of *Peziza* growing upon the ground and decaying wood, are familiar examples of the order.

92. Order h. Helvellales.—These are fleshy forms in which the hymenium is exposed over the surface of a variously shaped ascocarp, as shown in the common genera illustrated in Fig. 158. The morel or *Morchella,* in which the hymenium is spread



FIG. 158. Common forms of the Helvellales: *A,* the morel, *Morchella,* surface view at left and in section at right. The asci and paraphyses form a hymenium over the honeycomb surface. *B, Leotia,* a small gelatinous form of a light, bluish-green color. *C, Geoglossum,* a black tongue-like form. In *B* and *C* the hymenium is confined to the upper enlarged portion of the fungus. Both are common in boggy ground.

over the pitted or honey-combed surface of the ascocarp, is one of the most highly prized of the edible fungi. The members of

this order are largely saprophytic and often attain considerable size. Forms of *Morchella* occasionally reaching the height of a foot and some species of *Gyromitra* weigh over a pound.

Class C. Basidiomycetes or Basidia-forming Fungi

93. General Features.—This class contains two important groups of fungi; one of which includes very destructive parasites, and the other comprises those conspicuous and familiar saprophytes known as mushrooms, bracket fungi, puff balls, etc. The striking feature of the class is the formation of spores on club-like hyphae called basidia (sing. basidium) (Fig. 168, *D*). These organs are often developed in rather complex outgrowths of the mycelium which may be fleshy or woody, as in the mushroom and bracket fungi (Figs. 167; 170, *B*). There is no known sexual reproduction, although it is supposed that these basidia-producing structures may have been formed originally as the result of fertilization. The more important orders are the following:

94. Order a. Uredinales or Rusts.—These parasites are well known by the streaks and blotches of yellow or black rust which they produce on the leaves and stems of a great variety of plants. About 2,000 species are known in the United States. They are among the most destructive parasites, causing great damage to wheat, oats, apples, quinces, roses, carnations, etc. The yearly loss from grain rust alone is estimated at considerably over $18,000,000 in the United States. They exhibit a degree of variation not paralleled among other plants; as many as five different kinds of spores being formed by some species in their life history. This is due, doubtless, to the influence of the climate, the spores varying with the season (spring, summer and fall) and the plants upon which the fungus grows may be a contributing factor to the variation, for one and the same fungus may grow upon different plants, producing one or more kinds of spores on each.

(*a*) *The Life History of Puccinia.*—Several species of this genus infest wheat and illustrate the many forms that may ap-

pear in the life history of a rust. One phase of the life of this parasite appears upon the leaves of the barberry. During May and June the mycelium growing in the leaves forms roundish bodies which rupture the epidermis and finally open out into cups filled with chains of yellowish spores. An examination of Fig. 159 shows that these spores are found at the end of hyphae



FIG. 159. Cluster cups as seen in section of leaf of spring beauty, *Claytonia*. At right one of the cups is ruptured, exposing the aecidiospores. Below a small cup, pycnidium, is discharging pycniospores that are possibly functionless male gametes.

as in the powdery mildews and surrounded by a layer of rather thick-walled hyphae. This stage of the rust is known as the cluster cup or aecial stage and the spores are called aecidiospores. Often smaller spore-bearing cups, known as pycnidia (sing. pycnidium), are associated with this phase of the fungus. These small spores, while capable of germinating, do not appear to enter into the life history of the fungus by producing a new parasite. They have been looked upon as male gametes that originally effected fertilization in a female organ from which developed the spore-bearing cluster cup, the process being similar to that noted in the Red Algae where fertilization resulted in the formation of a spore containing cystocarp. The sexual proc-

ess has apparently entirely disappeared, although Blackman has called attention to indications of reproductive organs in the early development of the cluster cups and also to the fact that there is a fusion of nuclei in certain cells of the mycelium, similar to the fusion of gametes before the aecidiospores are developed.

The aecidiospores are scattered by the wind, and singularly, will only germinate and infest the seedlings and young leaves of the wheat in which they develop a mycelium similar to that of the barberry. However, this mycelium forms in place of cluster cups, groups of erect hyphae at the ends of which single reddish spores are formed (Fig. 160, *C*). This growth ruptures the epidermis, exposing the spores in rusty lines or blotches (Fig. 160, *B*). These yellow-brown blotches account for the popular name of Rusts given to this group of fungi. This is the summer or uredinal stage of the parasite. The spores are known as uredospores. This phase of the fungus is a very destructive one, for the spores are formed in great numbers and provided with thin walls. They are widely distributed by the wind and germinate at once (Fig. 161, *A*) on other wheat plants which soon show the rusty brown streaks of uredospores. In this way, the pest spreads with great rapidity from a single center of infection. This formation of uredospores goes on during the summer, sapping the vitality of the plant and, in severe cases of infection, materially interferes with the maturing of the grain.

Later in the season, this same mycelium forms in the leaves of the wheat quite a different type of spore. They are formed in the same manner as the uredospores but are provided with thick dark walls and from one to several spores are developed at the end of the hyphae (Fig. 160, *D*). Consequently, when the epidermis is ruptured, these spores form rusty black blotches on the leaves. This third stage is known as the telial, since it ends the season's growth. These spores, called teleutospores, are resting spores and tide the fungus over the winter. They germinate in the spring quite independent of any plant and being dependent upon the food stored in the spore, they only form short hyphae which usually become divided into four cells (Fig. 161, *B*). This structure is known as the basidium. Each cell of the basi-

dium sends out a delicate tube into the end of which the cell contents passes, thus forming a small spore, known as the basidiospore. This basidial stage completes the life history of the fungus for the basidiospores are carried to the leaves of the barberry and begin again the life cycle of the parasite by forming the cluster cups.



Fig. 160.



Fig. 161.

Fig. 160. The summer and fall stages of a rust, *Puccinia: A,* rust blotches on leaf of wheat. *B,* portion of leaf magnified, showing rupturing of the epidermis due to the formation of spores. *C,* uredospores or the summer spores which effect a rapid distribution of the parasite during the summer. *D,* teleutospores or fall spores which are dormant during the winter.

Fig. 161. Germination of uredo- and teleutospores: *A,* the thin-walled uredospore sending out hyphae from thin places in its wall. This is effected as soon as it is carried by the wind to a moist leaf. *B,* teleutospore germinating in the spring and forming a short hypha, from the end of which four cells have been cut off that are forming the basidiospores, *b.*

It is not surprising that this story was not unravelled for a long time and that these different stages of the parasite were known as distinct species. The first clue to the relationship was

gained in England where it was observed that the wheat fields to the leeward of the barberry bushes were especially infested with rust. For this reason, a law was passed early in the history of Massachusetts compelling the destruction of the barberry bushes. This suggestion of relationship between the cluster-cup stage of the barberry and the rust of the wheat finally led to the inoculation of wheat plants with aecidiospores and this resulted after a week or more in the appearance of the characteristic rusty streaks on the leaves of the wheat.

Considerable variation characterizes these rusts, not all of them having so elaborate a life history as that outlined above. One of the most common species of *Puccinia* affecting the wheat is perennial in the wheat and possibly in other grasses where it produces uredospores in the spring, thus, the aecial, telial and basidial stages are eliminated. In the apple rust, the uredinal stage is missing. This disease produces on the leaves of various members of the apple family, yellow patches in which are formed tube-like cluster cups (Fig. 162, *A*). The aecidiospores are only

FIG. 162. A rust, *Gymnosporangium*, that infests the juniper and members of the apple family: *A*, cluster cups on leaf of thorn apple. *B*, teleutospore stage on red cedar, juniper.

capable of infesting the juniper, on the branches of which they produce gall-like swellings (Fig. 162, *B*) known as cedar apples

and also bushy outgrowths known as witches' brooms. In the spring, tubular masses of hyphae bearing numerous teleutospores radiate out from these galls and in the early spring rains these strands swell up, forming conspicuous yellow, jelly-like masses. The basidiospores are developed in the jelly and infest the leaves of the apple, thorn, shadbush, etc. In some of the genera of rusts all the stages appear upon the same plant as in the May apple and jack-in-the-pulpit. In the early spring the stems and leaves may often be seen infected with the yellow cluster cups which are followed later by the dark-colored teleutospores. In other cases, teleutospores only are produced, as in the hollyhock, and this was doubtless the original form of the spore. Variations in the character of the spore resulted owing to the stimuli of the seasons, assisted possibly, by the different plants to which the parasite became adapted. It is evident that the formation of aecidio- and uredo-spores which occur in such large numbers and are easily transported by the wind and germinate quickly are all types of variation that would naturally be preserved since they are highly advantageous to the parasite.

95. Order b. Ustilaginales or the Smuts.—These parasites are more primitive than the rusts to which they are doubtless related. They produce but one form of spore that may possibly be compared to the teleutospore. They are common and exceedingly destructive parasites, affecting especially the flowers and fruits of corn and other cereals as wheat, oats and barley (Fig. 163). The damage by smuts to the corn crop of the United States exceeds $2,000,000 annually. In the case of corn, the parasite keeps pace with the growth of the plant without producing serious damage until the flowers appear, when the mycelium increases greatly in the affected ears and "tassels" producing sometimes enormous malformations, especially in the ears, which appear as glistening white blisters or pustules. Later, these bodies change to a sooty black, owing to the transformation of the cells of the mycelium (Fig. 164, *A*) into black greasy spores. These spores are scattered in clouds upon the breaking of the pustules and germinate much after the manner of the teleutospores of the rusts, infesting the young and delicate parts of

other corn plants by means of their basidiospores (Fig. 164, *B*). In other cases, the teleutospores germinate in the spring, infesting the seedlings. This appears to be always the case in the destructive wheat, rye, oats and barley smuts. Infection only

Fig. 163.

Fig. 164.

Fig. 163. A common smut, *Ustilago,* transforming the kernels of corn into sooty black pustules.

Fig. 164. The formation and germination of the spores of a smut: *A,* the formation of the spores from the mycelium in the kernel of corn. *B,* germination of a spore and the appearance of the basidiospore.

takes place when the spores come in contact with the seed. Consequently, in these cases, the fungus can more readily be fought by treating the seed with some fungicide as formaline and copper sulphate.

96. Order c. Agaricales or Mushrooms and Toadstools.— This is the most familiar group of the basidiomycetes and its 12,000 odd species are commonly referred to as mushrooms and toadstools. These fungi are largely saprophytes, living upon the humus in the soil and upon decaying wood. Several of the genera are exceedingly destructive to trees and cause great loss to the lumber industry. In the majority of cases, the fungus appears unable to attack the living portion of the tree and only

thrives upon dead tissues, as the heart wood, which they quickly disorganize and render worthless. As in all the preceding groups of fungi, the real plant body is a delicate mycelium that spreads through the soil or decaying substances, as may readily be demonstrated by splitting open a tree infested with one of the fungi (Fig. 165). The "spawn" that is sold in seed stores for plant-



Fig. 165.



Fig. 166.

Fig. 165. The mycelium of one of the Agaricales forming white masses as it spreads through wood.

Fig. 166. Development of a mushroom: 3, early appearance of the mushroom as a ball of hyphae on the strands of the mycelium. 1, section of one of these spherical masses of hyphae, showing the circular openings in which the gills are developed. 2, a later stage with the gills formed and the velum, *vl*, appearing as a delicate membrane.

ing in, mushroom beds is a dried mass of decaying leaves and straw mixed with earth, in which the mycelium of the mushroom has been allowed to grow. Often the hyphae of the mycelium lose their delicate character and become woven together, forming rather dense woody strands or plates, a development frequently

seen in the timbers of mines and under the bark of decaying trees. In certain species the mycelium is phosphorescent and the cause of the pale light that sometimes appears upon moist decaying wood, the so-called fox wood. The mycelium bears at various places the complex fleshy or woody body commonly known as the mushroom or toadstool. If the mycelium grows in a regular manner, radiating outward in all directions, then the mushrooms will have a similar arrangement. The older central portions of the mycelium will finally die off, while the newer portions continue to radiate outward and produce the mushrooms, which consequently appear in a more or less regular circle. In this way, the fairy rings, which are often held in superstitious awe, come about.

The Mushroom or Toadstool.—The common umbrella type of mushroom consists of a stalk or stipe and a cap or pileus, on the underside of which are located radiating plates or gills (Fig. 167, *A*). This structure originates on a strand of the mycelium as a very small ball composed of a mass of interwoven hyphae (Fig. 166, 3). Soon, however, the hyphae of this mass begin to grow in a very regular manner. At first the growth results in the formation of a cavity in the upper part of the ball that extends completely around it (Fig. 166, 1). Hyphae now grow down into this cavity in regular lines, forming the radiating plates or gills noted in Fig. 167. This development divides the ball of hyphae into an upper part or pileus and a basal region, the stalk or stipe. As this growth proceeds the mass of hyphae extending from the margin of the pileus to the stipe become drawn out and ultimately form a rather thin membrane known as the velum (Fig. 166, 2, *vl*). This entire growth is often enveloped by a sheath-like mass of hyphae, the volva (Fig. 167, *C, D*). At this stage of development, the young mushroom is rather spherical in shape and so small that it is quite concealed in the ground or by surrounding vegetation. When conditions are favorable for further growth, as after a rain, each cell of this miniature fungus absorbs moisture and rapidly expands, thus causing the mushroom to spring up as by magic and reach its full growth in a few hours. As the stipe elongates, the pileus

spreads out like an umbrella, rupturing the veil, a part of which clings to the stalk (Fig. 167, *A, a*) in the form of a ring, the annulus, and a part may also hang in the form of ragged fila-

FIG. 167. Habit of a poisonous mushroom, *Amanita: A,* the mature mushroom—*s,* stipe; *p,* pileus; *g,* gills; *a,* annulus; *v,* volva, a part of which appears in patches on the top of the pileus. *C,* young form of the *Amanita,* the volva beginning to break. *D,* later development, the volva completely ruptured, disclosing the pileus, stipe and velum, *vl.*—H. O. Hanson.

ments from the edge of the pileus. If a volva is formed, this is also ruptured by the elongation of the stipe, forming a cup at the base of the stipe, and usually portions also remain attached to the top of the pileus as scales and patches (Fig. 167, *A, D*). The structure of the mushroom is very simple. The stipe con-

sists of a mass of nearly parallel hyphae that show little modification save at the surface of the stem, where they are sometimes more compactly arranged (Fig. 168, *A, B*). By cutting across the gills, so that we can look into the ends of them, it will be seen that the hyphae extend down the center of the gills and also continually radiate out on either side, forming a compact layer of rather elongated cells on the surface of the gills, known as the hymenium (Fig. 168, *C*). A magnified view of a portion of this hymenium shows that it is composed of paraphyses and basidia (Fig. 168, *D*). The basidia are not divided as in the smuts and



FIG. 168. Structure of a mushroom: *A* and *B*, cross and longitudinal sections of a portion of the stipe, showing the character and arrangement of the hyphae that make up the mushroom. *C*, tangential view of the gills—*p*, pileus; *h*, hymenium appearing as a dark band on the surface of the gills. *D*, a portion of the hymenium enlarged—*b*, basidia; *pa*, paraphyses; *s*, basidiospores.

rusts, but the spores are formed in the same manner at the end of two or four small tubes that grow out from the apex of the basidia. It should be stated that a series of intermediate forms exist that connect the divided basidia with the present form. The spores are mature and begin to drop off as soon as the pileus opens, when they are scattered by the wind and develop under favorable conditions into a new mycelium. If the pileus of a freshly opened mushroom is placed on a sheet of gray paper and tightly covered with a dish, so as to exclude all air currents, the spores will fall directly upon the paper and in a few hours form an exact copy or a spore print of the gill arrangement. There are several very widely distributed and familiar families of the Agaricales, distinguished by the arrangement and distribution of their hymenium.

A. *Thelephoraceae.*—These fungi form membranous, leathery or woody incrustations or shell-like structures or branching bodies on soil or wood (Fig. 169, *A, B*). The hymenium forms a smooth or slightly wrinkled surface on the under side or exposed surface of the fungus.



Fig. 169. Forms of the Agaricales: *A, Stereum,* top view. *B,* underside, showing the characteristic smooth hymenial surface of the Telephoraceae. *C,* a coral fungus, *Clavaria,* with hymenium confined to the tips of the branches. *D,* a prickly fungus, *Hydnum,* with the hymenium on spine-like projection.

B. *Clavariaceae or Coral Fungi.*—This family includes the fairy clubs and coral-like fleshy masses of various colors (Fig. 169, *C*). The hymenium covers the branches.

C. *Hydnaceae or Prickly Fungi.*—These fungi form masses of widely various forms, but provided with spine-like outgrowths upon which the hymenium is developed (Fig. 169, *D*).

D. *Polyporaceae or Pore Fungi.*—This group includes largely leathery, woody or corky fungi in which the hymenium is developed on the surface of pores. They contain some of the most



FIG. 170. Pore forms of the Agaricales: *A, Boletus,* showing fleshy pore-bearing layer, *p*. *B,* top view of a woody, bracket form, *Elfvingia.* The concentric lines represent the annual growth. *C,* section of a similar form, showing three layers of pores that represent three years' growth. *D,* enlarged view of under surface, showing one of the pores with hymenial layer.

destructive of the timber-destroying fungi. Many of the genera of this family form shelf-like outgrowths on trees and are therefore known as bracket fungi (Fig. 170, *B–D*).

E. *Boletaceae or Fleshy Pore Fungi.*—The members of this family are generally characterized by a stalk and a pileus which bears pores, the latter being easily separable as a layer from the pileus (Fig. 170, *A*).

F. *Agaricaceae or Gill Fungi.*—These fungi more commonly assume an umbrella form, although some of them are of the shelving bracket type (Fig. 167). The hymenium is arranged on the surface of gills or plates. Many of them are highly

prized for the table, though they contain comparatively little nourishment, and must be regarded as relishes rather than foods. While the majority of the 5,000 species of the family are edible, some of them contain the most deadly poisons. No rule can be given that will enable the collector to separate these fungi into the two mythical groups of poisonous toadstools and edible mushrooms. Each species must be known individually before it is safe to use them for food. All forms with a volva should be most carefully identified, because this is a feature of the deadly amanitas (Fig. 167), which are among the most poisonous plants known.

The remaining orders of Basidiomycetes are characterized by the concealment of the basidia in cavities and are for this reason collectively known as the Gasteromycetes, meaning stomach fungi. Among the more common orders may be mentioned:

97. Order d. Lycoperdales or Puff Balls.—These familiar fungi are developed, as in the Agaricales, on strands of the mycelium, which often form extensive net-like threads in rotten stumps, logs, sawdust and humus (Fig. 171). The puff balls



FIG. 171. Cluster of common puffballs, *Lycoperdon*. At left three older ones have opened, permitting discharge of basidiospores.

vary in size from a pea to over a foot in diameter. When young, they consist of white cheesy masses of hyphae which form in the interior of the puff ball a series of irregular cavities lined with basidia and on the exterior, a rather firm skin or periderm (Fig. 172). At maturity, the inner hyphae break up, leaving only a dusty mass of spores and in some cases firmer hyphae, the capillitium. The skin ruptures in various ways; often by one or

more pores at the top, and the least touch now causes the spores to sift out in smoke-like puffs, hence the popular name of puff balls. In the earth stars (Fig. 173) the outer layer of the



FIG. 172. Diagram of a section of one of the puffballs, showing the thick skin or periderm and the irregular cavities which are lined with basidia. At the base the larger, sterile cavities of the stipe are shown.

periderm splits into rather regular star-like segments or valves which are hygroscopic. In damp weather these valves roll back, in some species to such an extent as to lift the puff ball from the



FIG. 173. One of the puff-balls popularly known as "earth stars," showing the outer periderm splitting into star-like sections and the inner periderm opening by a pore.

ground, when it may be set rolling by the wind and thus bring about a better discharge of the spores.

98. Order e. Nidulariales or Birds Nest Fungi.—These minute and curious fungi may be found growing upon twigs or upon bare ground in old fields, or upon dried dung (Fig. 174, *A*). They differ chiefly from the puff balls in that the spore-bearing cavities are surrounded by tougher hyphae. Consequently, when

the periderm of these little cup-shaped bodies open, these tougher parts appear as minute eggs in a nest (Fig. 174, *B*).

99. Order f. Phallales or Stink Horns.—These fungi first



Fig. 174. A common bird's-nest fungus, *Crucibulum: A,* habit of fungus on branch of hickory. *B,* one of the cups in section, showing the tough hyphae surrounding the spore-bearing cavities.



Fig. 175. A common form of the Phallales, *Phallus: A,* the so-called egg-stage which in *B,* sectional view, is seen to consist of an outer periderm, *p,* and within is a central stipe, *s,* capped with a pileus. The spore-bearing cavities of the pileus are shaded. *C,* stipe and pileus have emerged from the periderm, which forms an irregular sac about the base of the stipe. *D,* later stage after the spore-bearing layer has dissolved, revealing the cavities on surface of pileus.

appear as egg-like structures on rather coarse strands of the mycelium which traverse decaying vegetation. These bodies consist of a white skin-like periderm which encloses a stipe and pileus (Fig. 175, *A, B*). The basidia are formed in honeycomb-like cavities on the outer surface of the pileus. As soon as the spores are matured the stipe quickly elongates, rupturing the periderm and lifting the pileus into the air (Fig. 175, *C*). In structure and coloration, the Phallales are among the most attractive of the Basidiomycetes, but the majority of the forms are regarded with aversion, since the spore-bearing layer melts down into a slimy mass which emits a carrion-like stench.

Lichens

100. The Nature of Lichens.—These remarkable plants are of almost universal distribution upon tree trunks, rocks, old fences and buildings, and upon the bare earth, where they form variously colored incrustations or leaf-like branching bodies (Fig. 176).



FIG. 176. Common species of Lichens: *A*, an erect branching form, *Cladonia—as*, ascocarps. *B*, a foliaceous lichen, *Sticta*. *C*, *Parmelia* spreading over the bark of tree. The centrally-placed ascocarps are surrounded by smaller pycnidia.

The lichen is one of the most remarkable plants in the vegetable kingdom since it is a union of two separate plants, a fungus and an alga. Naturally the relationship of the lichens to other groups of plants has been a matter of dispute, some regarding them as constituting an independent division, and by others they are looked upon as fungi. The fact that the fungus forms the bulk

of the lichen and lives upon the alga largely as a parasite and is alone capable of forming reproductive spore bodies would lead to the latter position. The members of this curious co-partnership are largely ascomycetes and blue-green algae. Numerous species in each of these two groups of plants have become accustomed to living together. The formation of a lichen comes about through the attachment of a hypha of some fungus to one or more algal cells, as shown in Fig. 177, *B*. In this way, the food manufactured by the algae is absorbed osmotically by the hyphae. In some cases, the fungus obtains its food by means of haustoria which penetrate the algal cells. By the continued growth of the fungus and algae there finally results an interwoven mass of hyphae about the algae. The walls of the fungus are sufficiently transparent to permit the entrance of light and the color of the green algae can readily be seen when the lichen is moistened.

(*a*) *Structure of the Lichen.*—The bulk of the lichen is more usually composed of hyphae which show considerable regularity in their growth and the majority of the algae also are usually confined to a definite zone near the sunned surface of the lichen. Thus, in Fig. 177, *A*, which represents a cross-section of a lichen, it will be seen that the fungus forms a rather firm layer at the top and bottom of the thallus, while the algae are distributed near the upper surface, where they are exposed to the light, and can therefore carry on photosynthesis. Numerous hyphae projecting from the under surface of the lichen serve to anchor it to the substratum and also assist, doubtless, in the absorption of the earth substances. In some of the gelatinous lichens the fungi and algae are more promiscuously arranged.

While the fungus is dependent upon the foods manufactured by the algae, the latter are also benefited to an extent by this arrangement since they are protected by the strata of hyphae, and they are also provided with water and crude material which the fungus readily takes up and holds. This mutually helpful relationship of two organisms is termed symbiosis, meaning living together. The strange feature about this co-partnership is the marked change produced in the nature of the two symbionts. As long as they are independent of each other, very special conditions

are necessary for their welfare, but associated, they form the most resistant plants known. This accounts for their distribution from the equator to the pole and their association upon crystalline rocks, baked earth, bark of trees and other places where no other plant life is possible. Under unfavorable conditions the lichen becomes dry and brittle, in which resting condition it is able to meet any extreme of temperature and drought. With the return of suit-



Fig. 177.



Fig. 178.

Fig. 177. Structure of the Lichen: *A*, section of a lichen, showing the compact arrangement of the hyphae at the top and bottom, also the anchoring fungal threads on the underside and the dark algal cells, *a*, near the top. *B*, enlarged view of the algae to show their relation to the hyphae. *C*, diagram of one of the powdery particles, soredium, appearing upon certain lichens, showing the hyphae and algae. These bodies are scattered by the wind and form new lichens.

Fig. 178. Sexual reproduction of the lichen: *A*, section of an ascocarp, the hymenium appearing as a dark band in the mouth of the cup. *B*, enlarged view of the asci, *a*, and paraphyses, *p*, of the hymenium.

able moisture and heat they become leathery or gelatinous and renew their growth with considerable rapidity. Thus they live on from year to year, but owing to the exposed places in which they are usually found, their growing periods are frequently very short and their total annual growth may not exceed a few millimeters.

(*b*) *Reproduction of Lichens.*—Reproduction of the lichens is brought about by means of fragments that are readily detached when the lichens are dry and brittle, or by means of soredia. These latter bodies consist of a few algal cells intertwined with hyphae, in fact, a miniature lichen (Fig. 177, *C*). In some species the soredia form a rather powdery or granular coating on the upper surface of the lichen, and in other cases they are developed within the lichen. These bodies are easily scattered by the wind when the lichens are dry and under favorable conditions grow into new lichens. Reproduction is also effected by means of spores that are formed in asci since the species of fungi appearing in the lichens are largely ascomycetes. These reproductive parts appear as small open or closed ascocarps which develop a hymenial layer of asci and paraphyes, as already noted in the ascomycetes (Fig. 178). The ascocarps assume a great variety of forms, as seen in Fig. 176, *as.* In two genera the spores are borne on basidia and it is evident that certain species from a widely different group of fungi, the basidiomycetes, have acquired the habit of association with the algae. The origin of the ascocarp from a female gametangium has been observed in a few cases, and it is interesting to note that the male and female organs bear a more striking resemblance to those found in the red algae than has been observed among the ascomycetes. The small spores frequently developed in ascocarp-like bodies, as shown in Fig. 176, *C,* are considered as the male gametes.

The ascospores germinate readily, producing hyphae which, however, soon perish unless they chance to meet an alga with which they can live. It should be noted that these fungi appearing in lichens cannot associate indiscriminately with any species of algae. Each species of fungus is adapted to one or more species of algae and is unable to live with any other form. As may be imagined, the real nature of the lichen was for a long time misunderstood, the algae even being looked upon as spore bodies. The bitter dispute over the question was finally settled when a lichen was produced artificially by bringing together a suitable fungus and alga. It may also be stated that small lichen-like

bodies have been developed on glass plates from fungi alone by feeding them suitable nutrient solutions.

(*c*) *Economic Value.*—Lichens are of considerable economic importance in the world. Many genera live upon the rocks of which they appear to form an integral part. They are usually the first vegetation to appear in such places. The hyphae penetrate the irregularities of the rock and even effect disintegration of its substance. They also serve to collect the dust and other material that is washed down by rains and so prepare a soil for other plants. In many sections they form the characteristic vegetation of the country, as in certain alpine and desert regions and in the northern barrens, where they afford rich pasturage to reindeer and caribou. Their abundant growth in mountainous districts and subsequent distribution by winds and rains accounts for the showers of manna in biblical history. Certain species are still used in northern regions to lengthen out a meager food supply. Litmus, employed in acid testing, and various pigments are derived from lichens, as were also the famous purple and blue dyes of the East.

CHAPTER VII

DIVISION II. BRYOPHYTA. THE LIVERWORTS AND MOSSES

101. Adaptive Features of the Bryophyta.—We have now reached the point in the evolution of the plant where it has left the aquatic condition of its algal ancestors and become largely terrestrial. As has been stated, the fungi do not enter into the scheme of evolution of higher types. They are to be looked upon as a side line, derived doubtless from various groups of the algae, that have undergone a peculiar series of variation owing to their parasitic and saprophytic habits and that are in no way connected with higher plants. In the Bryophyta we take up again the line of ascent. Naturally we would expect that a change from the aquatic to terrestrial conditions would be attended with profound changes, tending to adapt the plants to the terrestrial conditions. The results of the stimulus of the new surroundings are seen in the production of hair-like outgrowths, rhizoids, that anchor the plant to the substratum and bring it into proper relation with the soil. So long as the plant lived in the water any of its cells might serve as absorbing organs, but it is evident that the provision must now be made for procuring the crude materials which are no longer in direct contact with the plant. The exposure of the plant to the atmosphere results also in the formation of a cuticle and of mucilage-producing cells which tend to retain the moisture and so adapt the plant to its drier environment. It will also be noted that modifications appear in the arrangement of the chlorophyll-bearing cells, with the result that they are always distributed in the same favorable positions for photosynthesis, as you have noted in the higher plants. However, the Bryophyta are only imperfectly adapted to terrestrial conditions and the entire plant body often assists in the absorption of moisture and mineral substances. For this reason they are moisture and shade-loving plants, some indeed being still aquatic, and they

remain rather small and inconspicuous, owing to the fact that they have not as yet varied and produced tissues that can supply them with an adequate amount of moisture and protect them against the climatic changes. Owing to their habit of growing together in colonies, they often become conspicuous and cover the bare earth, logs and tree trunks with swards and mats that constitute one of the most attractive features of the forest vegetation, and in northern regions they often form the most characteristic feature of the vegetation over large districts.

102. General Characteristics of the Bryophyta.—Certain groups of the Bryophyta are characterized by thalloid plant bodies, *i. e.,* without distinction of stem, root or leaf, while many have developed into creeping or erect plants with leaf-like organs. All are distinguished by a marked localization of the growing zone, the elongation of the plant being due to the repeated divisions of a single apical cell. The reproduction of the Bryophyta shows such marked departures that we realize that a wide gulf separates even the simplest of them from the Thallophyta. Notable among these departures is the absence of zoöspores. Only in a single genus are nonciliated bodies formed and discharged from certain cells as in the case of zoöspores. In this connection it is worth your time to look back and note the trend of plant life. The lowest forms of the green algae were essentially motile. Then the stationary condition arose and the motile state represented by the zoöspores became less and less conspicuous. In the bryophytes we perhaps have in the spores, referred to above, an interesting example of the last trace of the motile phase of plant life. The suppression by the sexual plant of the spore method of reproduction is due to several causes. They are able to increase their numbers in a vegetative manner. Special branches or ordinary shoots become detached from the parent plant by decay of the older parts or by other means and develop into new plants. Buds, called gemmae, consisting of one or more cells, are also very commonly formed on various parts of the plant and becoming detached grow into new plants (Fig. 186). Possibly also the higher organization of the Bryophyta, which renders them more capable of meeting conditions that would be

fatal to the lower forms, tended to make them independent of spore reproduction. But especially has this change been brought about by the fact that the gametospore in germinating develops an asexual plant or generation with enormous possibilities in the way of spore production, so that the increase and distribution of individuals is now fully provided for by the asexual plant and consequently largely removed from the sexual plant. This shifting of the responsibility for the increase in the number of individuals from the sexual to the asexual generation is one of the most significant departures in the evolution of plants. We will be interested to note how this change came about and how it steadily gains in importance in the remaining divisions.

The sexual reproduction of the bryophytes reveals a series of variations that are only remotely suggestive of the algae, but that indicates very clearly relationships with higher plants. These features will be considered in the discussion of the various groups. There are two classes of Bryophyta: A, The Hepaticae or Liverworts; B, The Musci or Mosses.

Class A. Hepaticae or Liverworts

103. Classification of the Liverworts.—While these plants are the simplest and least conspicuous of the Bryophyta, they are the most interesting, because among them will be found reminders of the algae, and at the same time they present many features that are suggestive of the mosses and ferns. They occur in very moist places and in deep woods, on rotting logs and moist shady banks. A few are aquatic, floating upon still ponds and streams, and some have become adapted to dry conditions. There are three orders: (*a*) Marchantiales or Thallose Liverworts; (*b*) Jungermaniales or Leafy Liverworts; (*c*) Anthocerotales or Horned Liverworts.

104. Order a. Marchantiales or Thallose Hepatics.—The liverworts of this order are characterized by flat, prostrate and rather fleshy bodies which usually grow upon the earth, to which they are attached by numerous rhizoids. Owing to the frequent division of the apical cell of the thallus into two equal parts, there

results two growing points which thus produce the equal forking or dichotomous branching of the thallus, so characteristic of these plants (Fig. 179, *A*, *B*). The appearance of many of these



FIG. 179. Forms of semiaquatic Marchantiales: *A, Riccia,* showing the dichotomous branching of the thallus. *B, Ricciocarpus.* The sexual organs are concealed in furrows that appear as radiating lines in the center of the branches. *C,* diagram of a cross-section of a branch, showing the male gametangia in the bottom of one of the furrows.

hepatics is suggestive of the algae. Especially is this true of the aquatic *Ricciocarpus* and *Riccia.*

(*a*) *Structure of Ricciocarpus.*—An examination of the structure of one of these will show, however, that extensive changes have been induced in even the simplest forms. The new stimuli to which the terrestrial conditions expose them cause a remarkable series of transformations in the cells that are cut off from the apical cell. The upper cells of the thallus, as soon as they are formed at the growing point, are exposed directly to the air and light, and they develop chlorophyll and grow up into vertical rows or plates just as you have already noticed in the palisade chlorenchyma of the leaf (Fig. 180, *A*). At an early period these rows of green cells become separated so that air spaces arise between them, and thus the chlorophyll-bearing cells are brought into direct contact with the atmosphere and enabled to carry on photosynthesis to the best advantage. The terminal cells of these rows enlarge considerably and form a rudimentary

epidermis. It is also interesting to note that the direct contact of the cells with the atmosphere results for the first time in the formation of a cuticle. In most species the air spaces arising among the upper cells of the thallus increase greatly in size and



FIG. 180. Structure of the thallus of *Ricciocarpus: A,* section of the thallus, showing the apical cell, *x,* forming cells that by further division develop into plates of cells separated by air spaces, *j.* At the left the plates thus formed are seen curving over the apical cell. *B,* an older portion of the upper part of the thallus. The air spaces, *j,* are greatly enlarged and the upper cells of the vertical plates have divided, arching over the air spaces but leaving small openings which permit the entrance of air for photosynthesis.—I. D. Cardiff.

the epidermal cells by vertical divisions keep pace with this enlargement and thus arch over the air cavities, forming, however, a small opening in the epidermal layer that permits a free circulation of the air (Fig. 180, *B*). These openings are suggestive of stomata and function in the same way, though they have originated in an entirely different manner. The lower cells of the thallus are compact and doubtless serve as storage cells, being nearly or quite destitute of chlorophyll. The stimulus of the soil causes the lower epidermal cells of the thallus to form numerous smooth or pitted rhizoids that are rather suggestive of the root-hairs of the higher plants, anchoring the plant to the substratum and assisting in the absorption of the earth substances. Plates of cells similar in origin to those occurring on the upper

surface are also found on the under side of the thallus (Fig. 180, *A*). These plates curve up around the growing point and doubtless protect it against drought, in which work they are assisted by the mucilage cells or glands. The development of these glandular cells on the plates, as well as on other parts of the thallus, adapt the plants to the drier terrestrial conditions to which they are exposed, as was the case with the mucilaginous walls of the Schizomycetes and Zygnematales.

(*b*) *Sexual Reproduction of Ricciocarpus.*—The gametes are produced in more complex gametangia than we have as yet seen. These organs are developed upon the upper surface of the thallus and in some species appear as lines radiating from the center of the plant. The male gametangium originates from one of the superficial cells of the thallus, which at first continues to divide transversely after the manner of the vertical plates of chlorophyll-bearing cells (Fig. 181, *A*). The cells of these vertical plates



FIG. 181. The origin and structure of the male gametangia or antheridia: *A*, section of the thallus, showing the apical cell, *x*, and the early stages, *a*, *b*, in the development of the antheridia. *B*, older antheridium with cells dividing vertically. *C*, later stage in which the wall cells are differentiated. *D*, mature antheridium of *Marchantia*, showing the numerous cells that develop the male gametes and the wall cells, *w*. *E*, greatly enlarged view of a male gamete after discharge from the antheridium.

soon begin to divide vertically and thus form an elliptical mass of cells (Fig. 181, *B, C*). As this growth goes on the outer or wall cells become larger than the others and generally develop chlorophyll, while the inner cells divide repeatedly and become very numerous with dense granular contents (Fig. 181, *D*). Each of the latter cells produces a single male gamete that is motile by means of two cilia (Fig. 181, *E*). This complex gametangium is suggestive of the structure we have seen in the *Ectocarpus* (Fig. 119, *B*). The male gametangium is commonly called the antheridium, and the male gametes are often termed sperms or antherozoids. The male gametes are discharged from the antheridium owing to the absorption of water by the walls of the central cells which become mucilaginous at maturity and expanding, rupture the apical cells of the wall and force out the contents of the antheridium. This mucilaginous mass rapidly dissolves in the water and sets free the male gametes or sperms. The female gametangium is also quite different from the single-celled gametangia which we have noticed among the algae and fungi. It originates as in the case of the antheridium, but develops into a flask-shaped body, consisting of a long neck of several rows of cells surrounding a central row, called the canal cells, and an enlarged basal region in which is developed the female gamete (Fig. 182). The female gametangium is commonly known as the archegonium and the female gamete is sometimes called the oösphere or egg. The female gamete presents essentially the same characteristics as seen in many of the algae, as *Vaucheria, Oedogonium,* etc., but it is surrounded by a jacket of cells which constitute the basal portion of the archegonium instead of being contained in a single cell. Possibly the archegonium has come about from a multicellular gametangium, as seen in the brown algae, owing to the sterilization of all but one of the cells. Archegonia, with two or more gamete-like cells, are sometimes found in the mosses and ferns. As soon as the female gamete is formed, the walls of the canal cells become mucilaginous and expanding through the absorption of water, force open the apical or lip cells of the neck, thus leaving an open passageway to the egg cell (Fig. 182, *F*). The male gametes, attracted, it is

supposed, by cane sugar, developed in the archegonium, swim down the canal of the neck, and one unites with the female gam-



FIG. 182. Development of the female gametangium or archegonium: *A–D,* stages in the development of the archegonium from a single cell. *E,* nearly mature archegonium just before the female gamete is formed—*cn,* canal cells. *F,* canal cells have dissolved, forcing open the lip cells of the neck. A male gamete has entered and is seen fusing with the nucleus of the large female gamete.—After Garber.

ete, as shown in Fig. 182, *F,* where the two gametes are seen fusing.

(*c*) *Germination of the Gametospore and Spore Formation.*—The gametospore, resulting from this fusion, becomes surrounded by a cell wall, but does not function as a resting spore as in some of the algae. It germinates at once and by repeated divisions of its nucleus forms a globular mass of cells, the capsule, within the archegonium which keeps pace with its growth (Fig. 183, *A, B*). The wall cells of this capsule are soon distinguishable, owing to their watery contents and shape, from the other cells, which are

rather cubical and densely granular. The growth and division of the granular cells continues until about 400 have been formed, when they round off, become separated from one another and increase greatly in size (Fig. 183, *C*). These large cells, called spore mother cells, form four spores each, as in the tetraspores



FIG. 183. Germination of the gametospore: *A*, basal portion of an archegonium, showing the germinating gametospore in the four-cell stage. *B*, later growth, forming a capsule with wall cells, *w*, which inclose large granular cells. *C*, a portion of the capsule, showing spore mother cells rounding off and floating in the fluid of the capsule. *D*, the mother cells dividing and forming four spores each.—H. O. Hanson.

of the red algae (Fig. 183, *D*). The delicate walls of the capsule break down as soon as the spores are matured, leaving them free in the archegonium, and later they are set free by the decay of the latter organ. These spores in some forms are provided with thick walls and are, therefore, resting spores adapted to

carrying the plant over unfavorable conditions for growth. In other cases the spores have thin walls and germinate at once. The advantage of the transference of the resting stage from a single gametospore to the numerous spores derived from the gametospore is manifest.

(*d*) *Germination of the Spore.*—The germination of the spore is usually indicated by the formation of chlorophyll, and this is



FIG. 184. Germination of the spores: *A*, spore. *B*, first division of the germ tube. *C*, early form of the thallus, due to the formation and subsequent divisions of an apical cell.

followed by the rupture of the outer spore coat and the protrusion of the inner as a delicate papilla or germ tube (Fig. 184). Usually from this tube a small, hair-like outgrowth is soon formed which penetrates the soil as the first rhizoid. The germ tube continues to elongate and often forms a chain of cells by successive transverse divisions, but eventually by oblique divisions an apical cell is formed that develops the characteristic thallus (Fig. 184, *C*). Thus we arrive again at the starting point in the life history of these simple plants.

(*e*) *Noteworthy Departures in the Life History.*—It should be stated that the gametophyte of certain species of the Marchantiales may live for long periods without producing the gametes. In fact, it frequently multiplies by means of buds and branches which become detached and grow directly into new plants. How-

ever, as soon as conditions are favorable the sexual organs and gemetes will appear. There are several features in this history that must be kept clearly in mind. In the first place, the gametospore is not discharged from the plant as in the case of the green or brown algae, but remains permanently in the archegonium, where it continues to be nourished by the plant during its germination and the formation of its spores. Therefore, it develops essentially as a parasite upon the plant, as is the case among the red algae. This retention and nourishment of the germinating gametospore in the plant is perhaps the most important of all the variations that appear in plant life. Owing to this relationship the gametospore attains a larger growth with increased power of spore production. In this way the development of many new plants is made possible by a single fusion of gametes. This is a very significant feature in terrestrial plants, since the fusion of the gametes is effected with more difficulty, owing to the absence of aquatic conditions. Without doubt this change was induced by the transferrence of foods from the sexual to the asexual plant derived from the gametospore. This loss of food gradually prevented the sexual plant from producing spores or other bodies designed to multiply its numbers and we will finally see that the sole work of the sexual plant is limited to the production of gametes. Note also that the gametespore not only has a larger growth but a longer life. This is of the utmost importance because it became exposed in this way to a new series of stimuli that affected it profoundly and that resulted in the evolution of the higher types of plants. Among the green algae the actual germination of the gametospore is limited to a few hours at the most and this is effected in the water, where the conditions are exceptionally uniform. The germination of the gametospore in the case of the liverworts is prolonged over several weeks, and more important still is the fact that this growth occurs on the land where it is exposed to a wide range of stimuli that cause it to vary in a most remarkable manner. So we must bear in mind the character of this rather simple growth derived from the gametospore of *Ricciocarpus* and compare it with the variations that appear in the succeeding forms.

It is also evident that the fusion of the gametes in the simple liverworts results in the production of a cell, the gametospore, that is radically different in its nature and possibilities of growth from any of the cells of the parent plant or liverwort. This was not apparent in the simpler forms of the algae, where the gametospore produced directly a plant like the parent, as in *Spirogyra, Vaucheria* and *Fucus.* In *Ulothrix, Oedogonium* and *Coleochaete,* we saw the first indication of the real nature of the gametospore. It did not develop into a plant like the parent, but produced zoöspores. The reason for this difference in behavior is possibly due to the higher organization of the gametospore. In the lower forms its first divisions result in the formation of cells that are of the same nature as the cells of the parent plant, but in higher forms its composition is more complex and as a consequence it forms a number of cells before cells like those of the parent plant are developed. For example, in the Red Algae, certain of the fungi and in *Ricciocarpus* the gametospore gives rise to a varying number of cells that are different from those of the parent plant, but finally spore mother cells are formed from which are derived spores that are of the same nature as those of the parent plant and that consequently produce new plants like the parent. As has been stated on page 124, one difference between the cells derived from the gametospore and those of the parent plant is to be found in the number of chromosomes which they contain, the former having twice as many chromosomes as the latter. The doubling of the chromosomes occurs when the two gametes unite to form the gametospore and this number is retained during the germination of the gametospore until the division of the spore mother cells, which results in the formation of spores with only one half the number of chromosomes. In the lower algae this reduction must occur in the first divisions of the germinating gametospore, which therefore corresponds to the spore mother cell, but in the Red Algae and *Ricciocarpus* a considerable growth intervenes before the spore mother cells appear and the reduction of the chromosomes takes place. In other words, as we ascend the scale of plant life, the formation of the spore mother cells and the reduction of the chromosomes is preceded by an ever-increas-

ing growth of the gametospore. This postponement in the formation of the spore mother cells, owing to the larger and larger growth of the gametospore, will steadily progress in the following studies.

These simple liverworts, like *Ricciocarpus,* show very clearly two phases or generations in their life history. The thallose plant is the gametophyte or sexual generation because it bears sexual cells or gametes. The capsule is the sporophyte or asex-



FIG. 185. Diagram of the life history of *Ricciocarpus.* The upper portion of the figure represents the sexual generation and the lower portion, the asexual. The former generation begins with the formation of the spores, *sp,* from the mother-cell and ends with the formation of the gametes, *g.* The asexual generation begins with the gametospore, *gm,* and ends with the spore mother cells, *sm.*

ual generation because it can only produce spores. The gametophyte begins with the spore and ends with the formation of the gametes. The sporophyte begins with the gametospore and ends with the division of the spore mother cell (Fig. 185). It may appear to you now as strange to regard the few cells of the capsule, the majority of which become spore mother cells, as a plant. But we shall directly see this microscopic plant assuming larger proportions as a result of its better nourishment and the stimuli to which it is exposed. It will form a larger and larger number of cells and the spore mother cells will not appear until late in its growth and they will be confined to definite regions and only constitute a small part of it. We now call the gametophyte the plant because it is many times larger than the sporophyte, but we will

see the sporophyte become quite as conspicuous as the gametophyte, as in the mosses, and finally it will become much larger, as in the ferns. Then we will call the sporophyte the plant.

(*f*) *Structure and Life History of Marchantia.*—Let us now examine a higher type of the Marchantiales, as *Marchantia,* and note the advances that have been made over simple forms like *Ricciocarpus. Marchantia* (Fig. 186) is of common occurrence

Fig. 186.



Fig. 187.

Fig. 186. Thallus of *Marchantia* bearing three cup-like organs that contain buds or gemmae and also four erect branches that contain the antheridia in their upper surfaces in radiating lines.

Fig. 187. Portion of the surface of the thallus of *Marchantia* enlarged, showing the rhomboidal air chambers and air pores.

in moist places, often appearing in greenhouses on the earth of flower pots, and it also forms luxuriant beds on the damp ground, especially where logs and brush have been burnt. The thallus shows the same general features that we have noticed in *Ricciocarpus,* being rather fleshy and creeping over the ground, to which it is attached by numerous rhizoids. The simple air chambers noted in *Ricciocarpus* are much enlarged and appear as diamond-shaped or rhomboidal plates on the surface of the thallus (Fig. 187). Thin sections across the thallus shows that these air chambers have a complex structure (Fig. 188). They originate in the upper cells of the thallus and as they enlarge they become covered by a well-developed epidermis which forms a chimney-like pore over the center of each cavity. From the bottom of the chamber numerous delicate chlorophyll-bearing cells arise. This arrangement of the tissues is again suggestive of the chlor-

enchyma of the leaf and it is manifestly protective and adapted to photosynthesis. The structure of the lower cells of the thallus



FIG. 188. Section through the center of the thallus of *Marchantia,* showing one of the air chambers and chimney-like air pores in the epidermis—*ch,* palisade-like chlorenchyma arising from bottom of air chamber. The lower cells of the thallus are nearly colorless and filled with watery solutions or mucilage. *r,* rhizoids; *l,* leaf-like plates of cells.

and the distribution of the rhizoids and ventral plates are essentially as in *Ricciocarpus.*



FIG. 189. Diagram of a section of one of the cups shown in Fig. 186—*g,* buds or gemmae associated with small glandular cells.

Marchantia, as in many of the liverworts and some mosses, multiplies extensively by means of buds or gemmae. They are borne in cup-shaped receptacles and consist of small lense-shaped masses of cells which are freed from the thallus by the swelling

of the mucilage secreted by glandular hairs growing among them (Fig. 189). These minute bodies grow directly into new plants and so serve to rapidly multiply the plant.

The reproductive organs appear upon different plants which may therefore be distinguished as antheridial or male plants and archegonial as female plants (Figs. 186, 190). Such a distribu-

FIG. 190. Thallus of *Marchantia* bearing several erect branches with lobed or umbrella-like tops. The archegonia are situated on the underside of the umbrella, between the lobes. *a*, the early appearance of a branch, in which condition fertilization is effected.

tion of the sexual organs is termed dioecious, meaning in two households, whereas *Ricciocarpus* and others are said to be monoecious, because the sexual organs are developed upon the same plant. The antheridia and archegonia are developed, as in *Ricciocarpus,* upon the upper surface of the thallus, but they are confined to special portions of it which appear at first as mushroom-shaped outgrowths (Fig. 190, *a*). These outgrowths, however, are but modified branches of the thallus and bear the sexual organs upon their upper surfaces in lines radiating from

the center of the branches. Fertilization is effected, as in all forms, by means of dews and rains which flood these organs with water. After fertilization these mushroom-shaped branches elongate, raising the sexual organs up into the air, where the antheridial branch assumes the form of a lobed disc, while the archegonial branch terminates in an umbrella-like structure (Figs. 186, 190). Owing to the extended growth of this latter branch, the



Fig. 191.



Fig. 192.

Fig. 191. Section of an archegonial branch similar to *a*, Fig. 190—*ar*, archegonia; *in*, involucre or curtain that hangs down on either side of the rows of archegonia; *ac*, air chambers; *th*, thallus.

Fig. 192. Section of a young antheridial branch: *an*, antheridia sunken in cavities of the branch, which is also provided with air chambers similar to those of the normal thallus. Some of the antheridia have discharged their gametes, as at *x*.

archegonia come to lie on the under side of the structure between the finger-like outgrowths, where they are completely hidden and protected by fringed curtains, the involucre, that hang down from the fingers (Figs. 191, 195, *A*). The antheridia retain their original position upon the branch, where they appear in cavities, as shown in Fig. 192.

The most important and significant departure in *Marchantia* is seen in the germination of the gametospore which divides into an inner and outer cell as in *Ricciocarpus* (Fig. 193, *B*). The inner cell, however, by a series of divisions, forms a rudimentary stalk and the lower part of it, known as the foot, comes into close

contact with the tissues of the plant from which it absorbs nourishment (Figs. 193, 194). The outer cell divides, forming a spherical mass of cells that resembles the capsule of *Ricciocarpus*. Some of these cells of the capsule develop as spore mother cells, while others, known as elaters, elongate greatly and serve to con-



FIG. 193. Germination of the gametospore: *A*, section of a mature archegonium with canal cells dissolved, thus forming a passageway to the large female gamete, *g*. *B*, sectional view of base of archegonium, showing the germinating gametospore in two-cell stage. The perianth, *p*, is seen growing up about the archegonium. *C*, later stage in the growth of the gametospore. The lower cell shown in *B* is forming stalk cells, while the outer cell has produced densely granular cells that will later by further division form spore mother cells and elaters.

duct the foods absorbed by the foot to the spore mother cells, and finally they become spirally thickened (Fig. 194, *C*). During the germination and growth of the gametospore, a delicate membrane (the perianth) grows up about the archegonium and doubtless assists the involucre in protecting it against drying winds (Fig. 193, *p*). When the spores are mature, the cells of the stalk

elongate, rupture the archegonium and push the capsule beyond the curtains of the perianth and involucre, so that it is exposed to the air (Fig. 195). The capsule now ruptures, exposing the spores and elaters to the air. The elaters, a word meaning lift-



Fig. 194. Structure of the nearly mature sporophyte: *A*, continuation of the growth in Fig. 193, *C*, showing the formation of a foot, *f*, stalk, and capsules, *c*, which contain elongated dark cells, the elaters, and the spores. The archegonium has been ruptured by the elongation of the stalk and is not shown in the figure. *p*, perianth. *B*, enlargement of the base of *A*, showing the attachment of the foot (indicated by darker lines) to the tissues of the antheridial stalk. *C*, an elater and spores.

Fig. 195. Archegonial branch with mature sporophytes: *A*, branch with the dark capsules of several sporophytes projecting beyond the curtains of the involucre. *B*, diagram of a branch as seen in section. On the right one of the capsules of a sporophyte ruptured, exposing the elaters and spores. On the left the curtain-like involucre only is shown.

ers, are very hygroscopic; they coil and uncoil with the least change in the humidity of the air and thus doubtless assist in the gradual exposure of the spores to the air currents. The spores germinate as in the lower liverworts. Thus we see that the sporophyte, consisting of a foot, stem and capsule, is more complex and larger than in *Ricciocarpus*. This is doubtless due to the better nourishment which it receives as a result of the development of a more efficient absorbing organ, the foot.

FIG. 196. One of simpler thalloid Jungermaniales, *Pallavicinia*, showing the rhizoidal growth on the under surface of the thallus and a mature sporophyte arising from a cup-like perianth which is surrounded at its base with involucrate leaves. —H. O. Hanson.

105. Order b. Jungermaniales or Leafy Hepatics.—By far the larger number of hepatics belong to this order. They are especially abundant in moist tropical countries, where they often cover the stems and leaves with a rich vegetation, and with us they are of common occurrence on dripping rocks and in deep woods on the moist bark of trees and decaying logs or damp earth. In the lower form the thallus is very simple and delicate (Fig. 196). In other genera there are indications of a lobing in the thallus which becomes more pronounced in some forms and leads by gradual gradations to genera in which the lobes appear as distinct leafy organs arranged in two rows upon a stem-like axis (Fig. 197). Usually a third row of rudimentary leaves, associated with numerous rhizoids, appears upon the under side of the thallus (Fig. 197, *B*). Although the Jungermaniales tend to become highly modified into a leafy body the tissues remain practically unchanged in all forms. Even in the leafy genera, which comprise the majority of forms in this order, there is little evidence of such a differentiation of the tissues as characterize the Marchantiales. Without doubt, the simple thallose Jungermaniales are the most primitive forms of the

hepatics and probably stand nearer the ancestral type than any other form. The Marchantiales represent a line in which the tissues of the thallus have attained a considerable degree of



FIG. 197. One of the leafy Jungermaniales, *Porella: A,* branch of the plant bearing several sporophytes. *B,* under surface of a branch, showing the lobing of the leaves and a row of minute scale-like leaves. *C,* portion of a branch bearing an archegonium surrounded by cup-like perianth with minute involucrate leaves at base. *D,* branch with cone-like antheridial branchlets. At the left a single leaf is shown with globular antheridium in its axis. *E,* section of a branch similar to *C.* The archegonium is seen surrounded by the perianth and below the involucrate leaves. The sporophyte is nearly mature and ready to elongate. It consists of a round capsule containing elaters and spore mother cells that are dividing to form four spores each. Below the capsule is the stalk or seta which ends in a foot buried in the tissues of the branch. At the right is an unfertilized archegonium that shows the original position and size of this organ.—H. O. Hanson.

differentiation, but in the present order, the evolution led to a marked change of form with but slight modification of the tissues. This is apparent in the highest forms where the leaves, consisting of little more than a single layer of chlorophyll-bearing cells, are arranged in two rather oblique rows upon delicate stems. The under surface of these leaves are generally lobed and often form sacs containing water (Fig. 197, *B*). This peculiarity of the leaves, together with the rudimentary third row of leaves that are associated with the rhizoids, make a sharp contrast between the upper and lower surfaces and serves to distinguish them from the mosses, with which they are often confused. In fact, they are often called scale mosses, owing to the moss-like appearance of their dorsal surface and the close contact formed with bark or other surfaces over which they creep. This habit of pressing the leaves against the substratum tends to retain the moisture and also enables the leaves to absorb it directly through their delicate cell walls. The overlapping of the leaves in many genera, as well as their lobing, would serve the same purpose and enable the plants to live under drier conditions than would otherwise be possible. Doubtless, these departures have been of great advantage to the plants and in part account for the common occurrence of these leafy forms. It should be stated that the leafy hepatics are regarded as forms that have been evolved from the simple thallose forms in quite recent geological times and owing to their better adaptation to present conditions they have become the most numerous of all the Hepaticae.

Reproductive Features of the Jungermaniales.—Asexual reproduction is of the same vegetative type as has been seen in the preceding groups. The archegonia and antheridia are also of essentially the same character as seen in Marchantiales. They are borne upon the dorsal surface of the thallus or upon more or less modified branches, the archegonia often arising upon the apex of the branch and the antheridia appearing as rather spherical bodies in the axis of the leaves (Fig. 197, *C, D*). The archegonia are developed in rather conspicuous cup-like or leafy outgrowths of the thallus. This structure,

termed the perianth (Fig. 197, *C*), assumes very characteristic forms in the different genera and is generally associated with modified leaves, the involucre. The gametospore develops after the manner noted in *Marchantia,* forming a capsule with elaters, but its stalk or seta reaches much larger dimensions, owing doubtless to the well-developed foot which penetrates the thallus and absorbs foods, thus making possible a larger sporophyte (Fig. 197, *E*). When the spores have been matured the seta rapidly elongates to several times its original length, rupturing the archegonium and lifting the capsule high in the air (Figs. 196; 197, *A*). The capsule usually breaks open into four valves which are hygroscopic, closing over the spores in damp weather and opening in dry weather to expose them to the wind. These spores germinate and begin the life history of a new gametophyte. As in some of the Marchantiales a filamentous algal-like growth is first formed by the germinating spore before the characteristic plant is reproduced.

106. Order c. Anthocerotales or Horned Liverworts.—This small group of three genera is the most interesting of all the

FIG. 198. One of the Anthocerotales, *Anthoceros,* bearing four pod-shaped sporophytes, *s*. The one on the right, *s'*, has opened at the top and is discharging the spores, but elongation and the formation of spores continues below owing to its basal growth.

hepatics because it presents features that are suggestive of the algae and also of a relationship with the mosses and ferns.

The thallus is of a primitive type, often with simple lobings and therefore, suggestive of relationship with the simpler Jungermaniales (Fig. 198). A peculiar feature of the order is the occurrence of mucilage cavities which communicate with the air through small openings on the under side of the thallus. These cavities are always occupied by one of the blue-green algae, *Nostoc,* which possibly assists the plants in the retention of water owing to their mucilaginous character. Another interesting feature of the order is the occurrence of a single chloroplast in each cell, as in *Coleochaete* and several other genera of the algae. The archegonia and antheridia, while resembling those of the preceding orders, are sunken in the tissues and also originate somewhat after the manner to be noted in the ferns.

The most suggestive variation of the Anthocerotales appears in the development of the sporophyte. We have noted that the asexual generation in forms like *Ricciocarpus* was a microscopic plant consisting of a delicate spore-bearing capsule. In *Marchantia* the sporophyte is still minute but differentiated into a foot, stalk and capsule, and in the Jungermaniales this becomes much more pronounced. In the Anthocerotales the sporophyte assumes a growth and differentiation that is much more extensive and complex and it also presents features that indicate a relationship with the mosses and possibly with the ferns. The germination of the gemetospore gives rise to a rather cylindrical or pod-shaped body (Fig. 199). The basal portion of the sporophyte develops into a massive foot, often provided with rhizoidal-like outgrowths, which serve as a very efficient absorbing organ. The upper portions of the sporophyte present a remarkable series of differentiations. The outer part of it consists of chlorophyll-bearing cells in which, for the first time, genuine stomata appear (Fig. 199, *ch*). Within this zone of chlorenchyma is a dome-shaped layer of spore mother cells alternating with sterile cells which in some genera develop as elaters. In the center of the sporophyte is a mass of sterile cells, the columella, which assist in conducting food from the foot to the forming spores. Directly above the foot is a region of rapidly dividing cells which causes the sporophyte to elongate by basal growth and push out

of the archegonium before the spores are mature. This is a radical departure from the other liverworts in the mode of development of the sporophyte and we will see essentially the same



Fig. 199.

Fig. 200.

Fig. 199. Section of a young sporophyte of *Anthoceros* emerging from the involucre-like outgrowth of the thallus—*sp*, dome-shaped spore forming layer of cells; *ch*, chlorenchyme with stomata; *b*, foot or absorbing region; *c*, region of growth. At right surface view of stoma.

Fig. 200. A common moss, *Funaria: A*, two plants with root-like rhizoids at base and radially arranged leaves. Rising above the leaves are the stalks or setae and capsules of two sporophytes. *B*, magnified view of a plant, showing the early appearance of the sporophyte as a delicate stalk still covered by the enlarging archegonium or calyptra. *C*, a plant bearing antheridia in a rosette of leaves at apex of stem. *D*, enlarged view of the upper portion of the sporophyte, showing the twisting of the stalk that assists in sifting the spores through the fringe of teeth, peristome, that encircle the mouth of the capsule.

manner of growth occurring in the mosses. As the sporophyte develops, the spores in the upper part of it mature and the sporo-

phyte splits into two valves (Fig. 198, *s'*), thus permitting the scattering of the spores. The lower portion, however, may continue to elongate for several months in some species, forming additional spores.

It is evident that this sporophyte would become a self-supporting plant if its foot should reach through the gametophyte and absorb substances from the soil. In comparing the sporophyte in forms like *Ricciocarpus, Marchantia* and the leafy hepatics with *Anthoceros,* we see that it has undergone a gradual evolution and has finally reached a point where it only needs to come in contact with the soil to become an independent plant since it is provided with all the tissues necessary for photosynthesis and absorption. Note also that the growth of the sporophyte is becoming more prolonged, that the spore mother cells appear later in its development and comprise but a small portion of it, and finally, that the sporogenous and sterile tissues alternate, forming a dome-like zone of cells. Structures very suggestive of these features will appear among the mosses and ferns.

Class B. Musci or Mosses

107. General Characteristics.—The mosses are by far the largest group of the Bryophyta and show marked advances over the hepatics in the development of the gametophyte and sporophyte. Particularly noticeable is the size and differentiation of the sporophyte which now becomes among the majority of the genera of equal importance with the gametophyte (Fig. 200). Variations appear in the mosses that have been successful in adapting them to a great variety of conditions, ranging from submerged aquatics to pronounced xerophytes that live upon exposed rocks. As a result, they are of common occurrence everywhere. The mosses, however, do not show any variations that have led to higher types of plants. In fact, they are regarded as a highly specialized group, like the Jungermaniales, that have branched off from the hepatics in recent geological times and become the dominant representatives of the Bryophyta owing to their better adaptation to present conditions upon the earth. This recent

derivation of the mosses is indicated by the remarkable uniformity of structure that characterizes the entire group.

While the plants themselves are relatively small, they often form large swards or mats, owing to the extensive branching and prolonged growth of the stems and their rapid multiplication through the dying off of the older parts and the independent growth of the branches that are thus set free. Owing to this habit of growth and multiplication the mosses often furnish the conspicuous features of the vegetation. Especially is this true in northern regions and in bogs and barrens where great sections of the country are carpeted with them. The mosses are leafy stemmed plants like the Jungermaniales, but they usually possess the decided advantage of having erect stems and radially arranged leaves (Fig. 200). Even in the prostrate forms where but two rows of leaves can be developed, the resemblance to the leafy hepatics is only superficial. The moss leaves are not lobed and the central portion is usually traversed by a strand of cells which serve for conduction like the vascular bundles of higher plants (Fig. 206, *B*). The differentiation of the tissues of the stem also show a marked advance over preceding forms. A central conducting region of elongated cells that may be compared to a rudimentary vascular system and a cortical zone, often with thickened cells and rudimentary epidermis, are frequently to be seen (Fig. 206, *A*). Asexual reproduction is almost entirely confined to the detachment of branches as stated above. In only a few genera have the formation of gemmae been noticed. The sexual reproduction and the development of the sporophyte present some interesting features that will be noticed in the following orders.

108. Order a. Sphagnales. Bog or Peat Mosses.—A single genus, *Sphagnum,* of about 20 species is the sole representative of this order which forms in several respects a transition from the hepatics to the mosses proper. These pale-green mosses (Fig. 201) grow on bogs and moors and other places where they are subject to drainage containing organic matter, as humic acid, derived from the decay of plant and animal life. The majority of plants are unable to endure these conditions, which

are popularly referred to as sour, and as a result, you will always find associated with the bog mosses a rather limited and peculiar variety of plants such as several genera of heaths, sedges, orchids, pitcher plants (*Sarracenia*) and other insectivorous plants like the sundew (*Drosera*). For some reason the ordinary plant is not able to procure its food from these sour bogs and this may explain the common occurrence in such places of insectivorous plants, and of certain trees and shrubs that are associated with mycorrhiza. The sphagnums grow luxuriantly in such places, the lower portions of the stem dying off and the upper portion branching and continuing the growth from year to year. In this way, ponds are gradually covered with a layer of mosses which become rather insecurely bound together by

FIG. 201.



FIG. 202.

FIG. 201. The bog mass, *Sphagnum,* bearing three sporophytes and numerous lateral branches covered with closely overlapping leaves. Note that some of these branches envelop the stem in wick-like strands.

FIG. 202. Structure of the leaf of *Sphagnum: A,* section of leaf, showing its single layer of cells that consists of large empty cells alternating with small chlorophyll-bearing ones. *B,* surface view of the cells, the larger empty cells being marked with spiral bands and often perforated with minute openings.

the subsequent introduction of other plants whose roots or stems traverse the covering formed by the mosses. So frail is this covering at first, that a very slight jar will cause the surface to tremble and for this reason these places are known as quaking bogs. The constant dropping of the decaying vegetation to the bottom of the ponds, assisted by the drainage material, gradually fills them, and in time they may become quite dry. The acid character of these bogs prevents the entrance of those organisms which promote decay. As a result, plants and animals that fall into these bogs may only partly decay and remain for ages in a wonderful state of preservation as is shown by the recovery of skeletons of animals and clothing of men that belonged to a prehistoric period.

(*a*) *The Structure and Reproduction of Sphagnum.*—The leaves of the bog moss are arranged in compact spirals around the stems and consist of a single layer of cells as in the leafy hepatics (Fig. 202, *B*). These cells, however, are of two kinds, large and empty cells with spirally thickened walls which are generally perforated with small pores and very narrow cells containing chlorophyll. This distribution of the cells explains the pale-green color characteristic of the sphagnums. Large and spirally marked cells, similar to those of the leaf, also occur in the cortex of the stem. The closely packed leaves enable the bog mosses to take up water like a sponge and the sphagnums are of considerable commercial value for this reason, being extensively employed by horticulturists to keep plants moist during shipment. Perhaps these large cells may serve as floats, enabling the plant to bridge over ponds and they may enable the plant to endure the acid waters in which these plants grow. In this connection, it is noteworthy that rhizoids are entirely lacking and that water and other substances are absorbed by the outer cells of the stems and by certain branches which hang down in wick-like strands close to the main stem (Fig. 201).

The antheridia and archegonia are developed much as in the leafy hepatics, the former organs appearing as stalked bodies in the axils of the leaves on short cone-like branches and the archegonia originate on the tips of short branches. The gameto-

spore germinates very much as the *Anthoceros* (Fig. 203, 6). It does not, however, have as prolonged a growth, and at maturity consists of a well developed foot embedded in the tissues of the gametophyte and a spore-bearing capsule (Fig. 203, 5*A*).



Fig. 203. The sporophyte of *Sphagnum:* 6, the young sporophyte separated from the archegonium with essentially the same differentiation of parts as noted in *Anthoceros,* Fig. 199. 5*A,* diagram of a later development of the sporophyte in the archegonium. The enlarged foot, *b,* is embedded in the apex of the moss branch, *p,* and the spores, *sp,* form a dome-shaped layer in the upper part of the capsule. 5, the naked stem, *p,* of the moss branch, surrounded at base with spirally arranged leaves, has elongated, lifting the mature sporophyte into the air. The enlargement of the stem, *b,* is due to the growth of the foot region of the sporophyte; *ca,* remains of the ruptured archegonium or calyptra; *o,* lid or operculum of the capsule.—After Schimper.

The stomata and chlorenchyma which were so conspicuous in *Anthoceros* are less perfectly represented, but the spore mother cells are still developed in a dome-shaped zone. This rather minute sporophyte at maturity barely breaks through the archegonium, known in this condition as the calyptra, which covers the capsule as a cap. The absence of a conspicuous stalk or seta in the sporophyte of *Sphagnum* is provided for by the elongation of the upper part of the moss stem which pushes the sporophyte

above the leaves and exposes the capsule to the winds quite as effectually as the seta of the hepatics and mosses (Fig. 203, 5). The spores are freed by the falling off of a circular lid, or,



FIG. 204.

FIG. 205.

FIG. 204. Germination of the spore of *Sphagnum: A,* early growth of the spore. *B,* later development—*sp,* spore; *b,* bud developing into moss plant. *C,* margin of the thallus, showing the origin of the bud.—After Campbell.

FIG. 205. The twisted stalk moss, *Funaria: A,* two sporophyte-bearing plants, the remains of the archegonium or calyptra still attached to the capsule on the right. *B,* enlarged view of plant with the young sporophyte, still enclosed in the archegonium, just emerging above the leaves. *C,* the male plant bearing the antheridia in a conspicuous rosette of leaves. *D,* upper portion of the sporophyte, showing the twisting of the stalk or seta that assist in sifting the spores through the teeth, peristome, that encircle the mouth of the capsule.

operculum, as is the case among the majority of the mosses (Fig. 203, *o*). This is effected by a ring of thin-walled cells, the annulus, that lie in the groove just below the operculum and which are very readily broken as the cells of the mature capsule

dry out. No elaters are associated with the spores in any of the mosses.

The germination of the spores results in the formation of a short chain or filament of cells which soon develop into a flat and rather irregular thallus consisting of a single layer of cells and attached to the ground by numerous rhizoids (Fig. 204). This thallus may produce from its marginal cells other filaments which behave like the original one and thus serve to increase the number of thalli. The archegonia and antheridia are not developed upon these thalli, but there are first formed as in some of the Jungermaniales, bud-like growths which develop into leafy stems of the sphagnum (Fig. 204, *b*). These stems soon become independent plants through the decay of the thallus and finally bear the reproductive organs.

109. Order b. Bryales. The Higher or True Mosses.—The great majority of plants, commonly known as mosses, belong to this order. As a rule, these plants are of a higher type than any of the other Bryophyta (Fig. 205). The stems are usually erect with radially arranged leaves and show considerable differentiation of tissues, as is apparent in the cross-section of the stem of one of the higher forms, shown in Fig. 206, *A*. Particularly noticeable are the thin-walled cells located in the central strand of the stem which serve to conduct the fluids and are therefore, comparable to the tracheal tissue of the higher plants. A similar conducting tissue is often seen in the central strand that appears in the leaves of the majority of the Bryales (Fig. 206, *B*). In connection with this higher development of the plant, attention should be directed to the more efficient organs of absorption. The rhizoids that spring from the base of the erect stems or from the under side of forms with creeping stems are often larger than those noted in the hepatics, and generally they are multicellular and often twisted together into root-like strands (Fig. 206, *C*). These features doubtless account in part for the larger size that is attained by many of the Bryales. Some of these mosses are short lived and many are perennial, continuing their apical growth from year to year. Some can live only in the wettest of places

or as submerged aquatics, while others are capable of enduring almost complete desiccation and revive quickly when moistened. An interesting device of service in adapting these plants to



FIG. 206. Structural features of the moss plant; *A*, cross-section through the basal region of a moss stem, showing rhizoidal outgrowths from epidermis, cortical and central conducting tissues. *B*, leaf of *Mnium* with midvein of elongated conducting cells. *C*, rhizoids twisted together into root-like strands. *D*, cross-section of a leaf of *Polytrichum*, showing the partial folding in of the margins of the leaf to protect the delicate plates of chlorophyll-bearing cells against drought.

periods of drought appears in the leaves of some of the higher forms, as the hair cap moss, *Polytrichum*. Plates of green cells parallel to the midvein project from the upper surface of the leaves, thus greatly increasing the surface of the leaves for photosynthesis and also serving as water reservoirs (Fig. 206, *D*). During drought, this delicate apparatus is protected by the coiling of the leaves into needle-like rolls which result in the exposure only of the thickened epidermal cells.

The sexual organs are developed on the apices of the main or lateral branches and they are protected by modified leaves which are often colored and form a more or less conspicuous bud or

FIG. 207. Male reproductive organs: 1, male plants of *Polytrichum* bearing antheridia in cup-like buds. 6, section of apex of stem of *Funaria* —*an*, antheridia; *p*, paraphysis. 7, male gametes, the one at left still enveloped in mucilaginous remains of mother cell wall.



FIG. 208. Female reproductive organs: 8, longitudinal section of apex of stem, showing arrangement of archegonia. 9, archegonium enlarged, the canal cells just beginning to dissolve—*g*, the female gamete.

cup (Fig. 207, 1). This is especially true of the antheridial buds, the plants being monoecious or dioecious. The antheridia and archegonia are essentially of the same structure as noted in previous groups and they are usually associated with modified leaves known as paraphyses (Figs. 207, 6; 208). The germination of the gametospore and the development of the sporophyte, while presenting many features in common with the hepatics and *Anthoceros* in particular, shows a remarkable series of variations that are of decided advantage to the plant. The gametospore in its early growth forms a spindle-shaped mass of cells, the basal portion of which reaches down into the stem of the moss plant and forms a well-developed absorbing organ or foot, while the upper portion elongates by means of an apical cell (Fig. 209, 10). Later, the growth becomes basal as in *Anthoceros*. For a time, the archegonium keeps pace with the elongation of the young sporophyte (Fig. 209, 11), but finally it is ruptured and lifted up as a cap, called the calyptra, on the apex



FIG. 209. Germination of the gametospore: 10, base of the archegonium in which the gametospore has germinated, forming a mass of cells with apical growing cell, *x*. 11, later growth of the gametospore. The sporophyte, *spr*, still enveloped by the archegonium, *ar*, appears as a cylindrical mass of cells with foot, *b*, penetrating the stem of the moss plant. At right an unfertilized archegonium. See Fig. 205, *B*.—After Sachs.

of the young sporophyte. This is one of the most essential differences between the mosses and hepatics. In the latter group, with the exception of the Anthocerotales, the spores were matured in the archegonium. In the Bryales, the spores are usually not formed until the seta has elongated considerably. As the sporophyte elongates, the upper part enlarges and finally forms a complex capsule which remains covered for a varying length of time by the calyptra.

The capsule and calyptra assume various forms and positions. In some cases the capsule is quite erect and completely covered by the calyptra (Fig. 210, 1), and again it may be more or less inclined or even pendulous and the calyptra laterally placed like a cap (Figs. 205, *A*; 211, 5). Removing the calyptra you notice that a lid or operculum is situated on top of the capsule while at the bottom is a somewhat enlarged region, the apophysis, provided with stomata that communicate with the interior regions of the capsule (Figs. 210, 2; 211, 5). When the operculum is removed a circle of minute teeth-like structures, the peristome, appear that more or less completely close the mouth of the capsule (Fig. 205, *D*). In some mosses a delicate membrane, the epiphragm, lies just below the peristome, which assists in closing the mouth of the capsule (Fig. 210, 2*B*, 4*A*). We are now interested to know the meaning of these structures and to learn how they have come about. By examining a longitudinal section of the nearly mature capsule that has been freed of its calyptra you will see that the spore-producing tissue is relatively small and appears as a hollow cylinder instead of a dome-shaped zone of cells (Figs. 210, 3; 211, 6). This sporogenous tissue is usually surrounded on the outside, and less commonly on the inside, by loose chlorophyll-bearing cells which extend down to the base of the capsule where chlorenchyma and stomata are developed. This lower region of the capsule, the apophysis, is often enlarged and provided with air spaces and stomata, thus constituting the chief photosynthetic apparatus of the sporophyte (Fig. 210, 3*A*). The entire surface of the capsule is protected by a well-developed epidermis but at the top a lid or operculum is cut off by a ring of rather delicate mucilage-bearing cells,

the annulus, which swell at maturity and so assist in casting off the operculum (Fig. 210, 3, *r*). Below the operculum a layer of cells, known as the peristome, extends across the capsule and



FIG. 210. The mature sporophyte of *Polytrichum:* 1, moss plant bearing sporophyte. 1*A*, calyptra removed, showing capsule, which has curved to one side. 2, capsule, the calyptra, 2*A*, removed, showing the operculum or lid and the enlarged apophysis, *a*, at base. 2*B*, capsule with operculum removed, showing the teeth-like peristome and the epiphragm, which has been lifted up at one side. 3, section of nearly mature capsule—*sp*, cylinder of spore-forming cells that is surrounded on its inner and outer sides by loosely arranged chlorophyll cells; *o*, operculum; *r*, annulus; *p*, peristome; *e*, epiphragm; *a*, apophysis. 3*A*, stoma of apophysis. 3*B*, cross-section of capsule—*sp*, spore-forming cells. 4*A*, portion of mouth of capsule, showing position of peristome and epiphragm in dry weather. 4*B*, peristome pressed down on epiphragm, closing the capsule in damp weather. —H. O. Hanson.

becomes greatly thickened on its outer and usually on its inner walls. This layer of cells is firmly attached to the walls of the capsule by rather short thick-walled cells (Figs. 210, 3, *p*; 211, 6–8, *p*). When the spores are mature, the more delicate cells break down, leaving little more than a loose mass of spores and the peristome within the capsule. The cells of the peristome break apart into teeth-like segments (Fig. 211, 7) and being



FIG. 211. Structure of capsule of *Funaria:* 5, capsule with calyptra, 5*A*, removed. 6, section of nearly mature capsule—*sp*, spore-forming cells surrounded on outside by loosely arranged chlorophyll-bearing cells; *o*, operculum; *r*, annulus; *p*, peristome; *a*, apophysis. 8, magnified view of a portion of the capsule, showing the annulus, *r*, and the thick-walled cells of the peristome, *p*, which are attached at their base to the epidermis by a double row of cells; *sp*, spore-forming cells. 7, the cells shown in 8, *p*, have split apart, thus forming the inner and outer teeth-like segments of the peristome.—After Sachs.

very hygroscopic, assist in forcing off the operculum. After the removal of the operculum the mouth of the capsule is covered by the peristome which now appears as a fringe of teeth radiating out from the walls of the capsule (Fig. 210, 4). The teeth of the peristome are always in multiples of two and form one or more, usually two layers, according as only the outer or both surfaces of the peristome cells are thickened. These teeth



FIG. 212. Germination of the spore: 3, early stage in the germination. 4, character of the branching, algal-like filaments that are finally developed from the spore—*r*, rhizoids which penetrate the ground; *b*, bud which will develop into a leafy moss plant.

are so variously marked and fashioned that they furnish one of the most characteristic features of the various genera of the mosses. The peristome closes the mouth of the capsule in wet weather, but on dry days they are constantly twisting and probably assist in the sifting out of the spores somewhat after the manner of the elaters of the hepatics. The dispersal of the spores is also brought about by the contraction and folding of the capsule and also by the seta which is very elastic and often hygroscopic (Fig. 205, *D*).

The spores germinate and produce profusely branching chains or filaments of green cells, called the protonema (Fig. 212). In a few instances, thalloid structures are developed as in *Sphagnum*.

The protonema absorbs the earth substances through delicate rhizoids and spreads over the ground, appearing to the eye as a green mould. Buds finally appear on these branches at numerous places and develop the leafy stems of the moss plant (Fig. 212, *b*). This in part accounts for the dense colonies so characteristic of the mosses. As soon as the young plants are established in the soil by means of rhizoids that spring from the basal parts, the protonema withers away, although in some genera it is perennial and continues to produce new plants from year to year.

110. The Relationship of the Bryophyta.—It appears probable that there have been three lines of variation or evolution among the Bryophyta that have had their origin in some simple thalloid form most nearly related to the lower Jungermaniales. One line has resulted in the highly differentiated thallus of the Marchantiales with its simple sporophyte. Another, the Jungermaniales, retaining the simple structures of the primitive thallus has modified its form into a leafy plant and developed a more complex sporophyte, while the variations of the third line, or Anthocerotales, have resulted in slight alterations of the primitive type of the thallus but in profound modifications of the sporophyte. The variations of the sporophyte in the latter group appear to have been very advantageous to the present conditions upon the earth and doubtless lead to such forms as the sphagnums and other mosses. The important feature in this evolution, already noted among the algae, is the increasing importance of the sporophyte. Among the green algae it was dependent upon the amount of food stored in the gametospore. In the Bryophyta, it is better nourished, owing to its essential parasitic habit. This led to a larger growth and more complex development and it is especially important to note that the sporophyte became more profoundly modified by the stimuli of external conditions than did the gametophyte. For example, we see in the sporophyte of *Anthoceros* and of the mosses, a plant that is practically dependent upon the gametophyte only for water and the earth substances. So the way is prepared for a study of the next division where the sporophyte finally comes to absorb its crude materials directly from the soil and thus becomes an independent self-supporting plant.

CHAPTER VIII

DIVISION III. PTERIDOPHYTA OR FERNS

111. Nature of the Ferns.—The members of this division differ greatly in character and live under a wide range of conditions, although, like the Bryophyta, they usually frequent moist and shady places, as glens, ravines and damp woods, or in tropical countries they often cover the moist rocks and tree trunks with a luxuriant and attractive vegetation. The most noticeable fea-

FIG. 213. The Christmas fern, *Polystichum*, with young leaves uncoiling in the spring at the tip of the stem and further back the fully developed leaves of the past season. The older portion of the stem is covered with the petioles of leaves that have died off and numerous roots are seen arising from the stem and bases of the petioles.—H. O. Hanson.

ture in the fern is the large development and complex character of the plant (Fig. 213). This is due to a more efficient absorbing and conducting apparatus, the root and vascular bundle, which are essentially of the same character as in the higher plants. Furthermore, this fern plant does not correspond to the moss plant. It is the sporophyte which has become independent of the gametophyte. The sexual generation is a small thallus, commonly called the prothallium, that is quite as primitive as the simplest Jungermaniales. Its chief function is the formation of gametes (page 254) and it is capable of nourishing the sporophyte only for a short time in the majority of ferns (Fig. 214). The gametophyte of the Bryophyta never attained any considerable proportions owing to its inability to vary and produce adequate absorbing organs and conducting tissues. This naturally limited the development of the sporophyte which was parasitic upon it. In the Pteridophyta, however, the sporophyte is not at such a disadvantage since it actually becomes independent of

FIG. 214. The thalloid gametophyte or prothallium, *p*, of a fern from which a young fern plant is being developed. This sporophyte has already developed two roots, *r*, that penetrate the soil, and two leaves, *l*, are appearing, thus making the fern at an early age independent of the gametophyte, *rh*, rhizoids.

the gametophyte owing to the development of a root which puts it in communication with the earth substances. This radical de-

parture in its mode of life acted as a profound stimulus and coöperated in inducing marked variations that were so beneficial in character as to cause the sporophyte to assume large proportions and become differentiated into a highly organized plant. The stem elongates through the repeated division of a single apical cell, as in the Bryophyta, but the stem and also the leaves and roots in addition contain vascular bundles and an arrangement of tissues already noticed in the higher plants.

The gametophyte in many of the Pteridophyta is very suggestive of the simpler thalloid hepatics. Owing to the fact that it is no longer permanently burdened with the nutrition of the sporophyte, it becomes greatly reduced in size and length of life. In fact, in several of the more specialized ferns, the gametophyte may be reduced to a few cells and the entire development may take place within a day. The reproductive organs and the germination of the gametospore are suggestive of the corresponding features noted in the Bryophyta. The more important orders of the Pteridophyta are: 1. Ophioglossales or Adder Tongue Ferns. 2. Filicales or Common Ferns. 3. Equisetales or Horsetail Ferns. 4. Lycopodiales or Club Moss Ferns.

Order 1. Ophioglossales or Adder Tongue Ferns

112. General Characters of the Sporophyte.—This order contains three genera of very primitive ferns which are probably but a remnant of an earlier and widely distributed group. Only two, *Ophioglossum* and *Botrychium,* are of common occurrence (Fig. 215). They are of unusual interest because they present many features suggestive of the liverworts and also of the more specialized ferns and seed plants. The sporophyte consists of a short upright stem with thick fleshy roots that are associated with mycorrhiza. The leaves are simple or divided and usually appear singly, ensheathing the apex of the stem. One of the most remarkable features about these ferns is the development in the stems, roots and leaves of the same tissues that we have noted in the higher seed plants. The vascular bundles are usually collateral and in the larger species of *Botrychium* are arranged

in the stem in a circle and provided with a cambium, so that there is a striking resemblance to the dicotyledonous stem with its pith,



FIG. 215. The two common genera of the Ophioglossales: *A*, the adder-tongue fern, *Ophioglossum*, with single leaf ensheathing the short stem and producing a spore-bearing spike. *B*, section of the spike—*sp*, the spore-forming cells arranged in groups or sporangia. The spores are exposed by the breaking apart of the cells between the dark lines. *C*, stoma from epidermis of spike. *D*, the grape fern, *Botrychium*. The leaf is much divided and also forms a branched spore-bearing organ. *E*, two sporangia, showing the manner of opening for discharge of spores.

ring of xylem, cambium and phloem, medullary rays, cortex, cork, etc. (Fig. 216).

The spores are borne in peculiar modified outgrowths of the leaves that assume a cylindrical shape in *Ophioglossum* and become more or less branched in the species of the *Botrychium* (Fig. 215). The structure of these organs presents many features that recall the sporophyte of *Anthoceros* (Figs. 215, *B*; 199). The outer portion of the cylinder consists of chlorophyll-bearing cells which communicate with stomata in some of the

forms at least. The spore mother cells are also distributed as in *Anthoceros,* the essential difference being that they are separated by a larger number of sterile cells and a larger number of



Fig. 216. Cross-section of a stem of *Botrychium: p,* pith; *x,* xylem; *m,* medullary ray; *c,* cambium; *ph,* phloem; *e,* endodermis; *cr,* cortex.—After Jeffrey.

spore mother cells are also grouped together, forming rather conspicuous sacs or sporangia (Fig. 215, *sp*). The spores are formed from the mother cells as in the Bryophyta and are discharged



Fig. 217. Arrangement of the sporangia of an allied order, Marattiales: *A,* leaflet of *Archangiopteris* with sporangia or surface of leaf and arranged in groups or sori. *B,* magnified view of a portion of the leaflet. *C,* section of leaf, showing two sporangia, the left-hand one in section.

from the sporangia through a transverse cleft (Fig. 215, *E*). It is interesting to note in a closely allied order occurring in the tropics, that the sporangia are developed directly upon the leaves (Fig. 217) instead of upon special branches and that they are sometimes associated together in groups, known as sori (sing.

sorus) and provided with thickened cells, the annulus—characters that will become conspicuous in the next order. These sporangia originate, however, as in *Ophioglossum* and open by transverse clefts which operation is promoted by the unequal drying of the thick- and thin-walled cells of the annulate forms.

(*a*) *The Gametophyte.*—The gametophyte or sexual generation is not often met with, owing to the fact that, like the sporophyte, its development is associated with mycorrhizal fungi and consequently it is usually completely buried in the soil and usually quite devoid of chlorophyll. The various stages in the germination of the spores have never been observed, but the mature gametophyte of *Botrychium* was found a few years ago by Jeffrey and carefully studied. It appeared as a rather tuberous body that is provided with numerous rhizoids and a growing apical cell as in the Bryophyta (Fig. 218, *A, B*). The archegonia and antheridia are usually borne upon the upper surface of the gametophyte and in origin and appearance are rather suggestive of *Anthoceros.* The antheridia consists of a number of mother cells quite buried in the tissues of the thallus, being covered by one or two layers of cells which are destroyed or break open when the male gametes are mature (Fig. 218, *C*). These gametes are quite different from any yet seen, being spirally coiled and provided with numerous cilia (Fig. 218, *D*). The archegonia are similar to those noted in *Anthoceros,* consisting of a rather short neck with two central canal cells and a swollen base sunken in the tissues of the gametophyte (Fig. 218, *E*). The female gamete is formed at the base of the canal cells which finally become mucilaginous and force open the lip cells of the neck, thus forming an open passage-way for the entrance of the male gametes. It has been shown in the common ferns that malic acid is present in this mucilaginous substance which strongly attracts the male gametes so that they crowd into the canal, often completely choking it. So we see that the gametophyte of this fern (though differing in form) is essentially of the same character as in the Bryophyta, its subterranean character and absence of chlorophyll being due to its association with fungi. It should be stated that lobes are produced from the end of the tuberous gametophyte

of one of the species which become green on reaching the surface of the soil and in a closely allied order of tropical ferns, the sexual generation is normally a green thallus strikingly like some of the simpler Jungermaniales.



FIG. 218. The gametophyte and young sporophyte of *Botrychium: A,* tuberous appearance of the gametophyte—*em,* a young sporophyte or embryo developing in one of the archegonia. *B,* a section of gametophyte—*ar,* archegonia; *an,* antheridia. *C,* two antheridia in section. *D,* male gamete. *E,* archegonium before dissolution of canal cells. *F,* gametospore in two-cell stage of germination. *G,* young sporophyte with roots and first leaf developed but still attached by foot to the round gametophyte, *gm. A–F.*—After Jeffrey.

(*b*) *The Germination of the Gametospore.*—The early development of the sporophyte resembles *Anthoceros* in many respects. The gametospore divides into two cells (Fig. 218, *F*), from the lower of which a large foot is formed, and from the upper cell the short stem and root arise. Later, the first leaf or cotyledon is formed from the stem. This development of the sporophyte

goes on very slowly within the archegonium where it lives as a parasite for a long time. Eventually the root ruptures the archegonium or calyptra and comes in contact with the soil. The cotyledon now grows upward and makes its appearance above the ground as the first green leaf (Fig. 218, *b*). The sporophyte thus becomes a self-supporting plant, although it probably remains in part dependent upon the gametophyte for several years. At this stage of development the young sporophyte is a very simple type of fern, but gradually more efficient roots are developed and each year the stem sends up a larger and larger leaf until the adult size is reached when the spore-bearing branch is formed (Fig. 215), thus making possible again a new series of gametophytes.

(*c*) *Comparison of the Adder Tongue Ferns with Preceding Groups.*—The most noteworthy difference between these simple ferns and the Bryophyta is the larger development of the sporophyte and its final independence of the gametophyte. This is doubtless due to the development of true roots which made possible a continuous and abundant supply of the crude materials from the soil. This change acted as a stimulus which promoted variations in the sporophyte while the light and various climatic factors also assisted to a very marked degree. Among the algae, the gametophyte is the dominant generation, the sporophyte being represented often by the gametospore. This is essentially the relationship among the majority of the Hepaticae where the gametophyte performs all the work of food construction and the sporophyte is a minute parasitic plant upon it. In *Anthoceros* and the mosses, the sporophyte is longer lived and more highly organized, but the gametophyte is still the more important generation as it performs the major portion of the work. Among the ferns, this relationship and the distribution of labor is completely turned about. The sporophyte becomes the larger plant and the principal center of photosynthesis, while the gametophyte remains small and gradually becomes very short lived.

There is also to be noted the longer postponement in the formation of the spores. In certain of the algae the gametospores produce the spores directly on germinating while in *Ricciocarpus*

a small mass of cells is first formed and certain of these cells soon become spore mother cells. In *Marchantia,* a larger number of cells are formed by the gametospore and the spore mother cells originate after a few weeks in a definite region of the sporophyte. These features are more noticeable in the Jungermaniales and especially in the long-lived sporophyte of *Anthoceros.* Recall also the considerable growth of the sporophyte of the Bryales and the final development of the spore mother cells in a definite and restricted region in the capsule. In the Pteridophyta, the sporophyte not only becomes larger and more highly differentiated, but the formation of the spores is often deferred for years and the spore mother cells are localized in special organs. This postponement of the spore formation may appear at first sight as a disadvantage since the sporophyte is exposed to many dangers during the long period of preparation for its work. However, the delay is more than compensated for by the large number of spores that are finally formed and also by the fact that the sporophyte does not entirely perish as in previous cases, but lives on, sending up annually new spore-bearing leaves. It would appear that a point had at last been reached in the evolution of the sporophyte where it is so well organized as to ensure its existence. It can consequently with safety defer the formation of spores until well-developed roots, stems and leaves have been formed. The independent existence of the sporophyte brings out very clearly the two phases in the life history of the plant which we call the alternation of generations. The formation of the spores mark the beginning of the gametophyte or sexual generation, which ends with the formation of the gametes. The formation of the gametospore through the fusion of the gametes starts the sporophyte or asexual generation which ends with the division of the spore mother cells.

Why has not the gametophyte varied as well as the sporophyte? Perhaps the simplicity of its structure has become fixed by inheritance from a long line of algal ancestors. The occurrence in the Bryophyta and Pteridophyta of motile male gametes and the consequent necessity of water for fertilization, points to the inheritance of this peculiarity from aquatic ancestors. It is evident

that there can be no considerable modification of the gametophyte, as long as this feature is retained, without greatly decreasing the possibilities of fertilization. Consequently the gametophyte must remain of necessity a primitive structure. The sporophyte, on the other hand, was accustomed from the first to terrestrial conditions, or to such conditions as were unfavorable to the growth of the gametophyte. Among the algae we have noticed that it may carry the plant over the more or less complete drying up of the water or unfavorable temperatures. Perhaps these exposures to a variety of stimuli which the gametophyte never experienced or in a lesser degree enabled the sporophyte to respond with more profound variations when it became parasitic upon the gametophyte and exposed directly to the light and air as was the case among the Bryophyta. Certainly we know that abundance of foods, light, temperatures, etc., are among the important stimuli in causing variations. The simple sporophyte of the Bryophyta was exposed to just such forces as these and steadily gained in complexity until in the Bryales it nearly equaled in importance the gametophyte. The sporophyte of the Pteridophyta, owing to the development of the root, experiences a new stimulus, *i. e.*, that of the soil which, coöperating with the stimuli of the light, etc., causes it to assume much larger proportions than the gametophyte and become the dominant generation in the life history of the plant.

Order 2. Filicales or Common Ferns

113. General Features.—The great majority of plants popularly known as ferns belong to this order. They are a highly specialized group that have branched off from the primitive ferns in recent geological times and owing to their variations being highly adaptive to present conditions upon the earth, they have become very numerous and widely distributed. In temperate climates the majority live upon the ground in moist and shady regions and some are aquatic or xerophytic. They attain their greatest abundance in the mountainous district of tropical countries where they occur in astonishing profusion and variety upon the moist rocks and trunks of trees, as well as upon the earth.

The leaves are the most striking feature of the Filicales. They are usually large and divided and are characterized by being coiled when young (Fig. 219). This is due to the stronger

FIG. 219. Christmas fern, *Polystichum*, with prostrate stem bearing at the tip young coiled leaves covered with chaffy scales and further back large leaves of the previous season. The older portions of the stem are covered with the petioles of dead leaves.—H. O. Hanson.

growth of the outer cells which causes an inward rolling of the leaf. The growth of the leaf is very slow, often requiring three years for its formation in temperate regions. During the season preceding its expansion, the petiole and blade are completely formed and appear in crosier-like coils more or less covered with chaffy scales (Fig. 220, *c*). This development enables the leaves to expand with surprising rapidity in the spring when the more rapid enlargement of the cells on the inner side of the leaf cause

it to uncoil. The tissues of the leaves are differentiated into an epidermis, stomata, chlorenchyma and vascular bundles as in the higher plants (Fig. 223). The stems are more usually pros-



Fig. 220. Stem in early spring freed from all of its leaves save the young ones, *c*, near the tip: *r*, roots; *v*, vascular bundles in base of petiole; *x*, region from which the cortex has been removed to show the vascular bundles from the leaves uniting to form a network.

trate, creeping rhizomes that bear a few large leaves near their growing tips. The stems branch sparingly and so gradually give rise to colonies of ferns. One of the most attractive fea-



Fig. 221. Cross-section of fern stem: *v*, concentric vascular bundle, the large cells of the xylem being surrounded by the phloem. Each bundle is surrounded by a compact layer of cells, the endodermis; *st*, stereome; *cr*, cortical region.

tures of certain tropical districts are the tree ferns with erect stems of palm-like appearance which lift their great crowns of leaves 30 to 50 feet in the air. The vascular bundles are of

the concentric type (Fig. 221) and usually form a hollow network in the stem (Fig. 220, *x*). Certain cells of the cortex and pith often become modified into strengthening cells of stereome (Fig. 221, *st*). Roots arise near the base of the leaves, and in some of the tree ferns form a thick mat-like covering on the stems. They originate from the endodermis of the bundles and possess a root cap and radial arrangement of the vascular bundles as in higher plants.

(*a*) *Structure and Character of the Sporangia.*—The sporangia, instead of being produced in the tissues of special branches as in *Ophioglossum,* are borne in curiously constructed capsules, usually situated on the under surface of the ordinary green leaves (Fig. 222). The sporangia-bearing leaves are usually

FIG. 222. Arrangement of the sporangia: *A,* lobe of leaf of *Dryopteris* with sporangia grouped in circular sori, *s*. *B,* sorus enlarged, showing the shield-like membrane or indusium, *in,* covering the sporangia, *sp*. *C,* lobe of leaf of *Asplenium* with elongated sori, *s*.

called sporophylls, meaning spore-bearing leaves. In some cases the sporophylls are highly modified, being entirely given up to spore production and therefore quite different from the green leaves (Fig. 227). The sporangia are usually associated in groups or sori (sing. sorus) on the vascular bundles and pro-

tected by a membranous outgrowth of the epidermis, known as the indusium (Fig. 222, *B*). Each sporangium originates usually from a single epidermal cell, which by repeated divisions

FIG. 223. Section of a leaf of *Woodwardia*, showing two sori: *i*, indusium; *sp*, sporangia arising from the epidermis and in various stages of development. Note the epidermis, stoma, *s*, palisade and spongy chlorenchyma, and vascular bundles, as in higher plants.

(Fig. 223) produces a capsule or sporangium that varies considerably in structure in the various genera. In the simplest forms, it consists of a uniform layer of wall cells inclosing a mass of cells, the majority of which become spore mother cells,



FIG. 224. Character of sporangia: *A*, simple type of sporangium of *Osmunda* with rudimentary annulus, *an*, of a few thickened cells. *B*, common type of sporangium—*an*, annulus; *l*, lip cells.

producing four spores each. In other cases, a few of the wall cells are thickened, forming a rudimentary annulus (Fig. 224, *A*). but in the majority of our common ferns, the sporangium con-

sists of a delicate stalk, supporting a rather spherical spore-bearing capsule (Fig. 224, *B*). The walls of this capsule consist of a single layer of thin-walled cells except for a row of thickened cells, the annulus, that extend from the stalk over the capsule nearly to the opposite side (Fig. 224, *B, an*). At the latter point, the annulus ends in a few rather weakened cells, two of which, the lip cells, are conspicuous for their larger size (Fig. 224, *B, l*). It is to be noticed that the walls of the annulus are thickened on their inner and radial walls while the outer walls remain comparatively thin. When the spores are mature and lie loose in the capsule, the cells of the annulus begin to dry out. It is evident that the thin outer walls will be the first to yield and their contraction pulls open the sporangium at the weak lip cells, the split extending back through the lateral cells of the capsule which is finally drawn back to such an extent that the annulus becomes nearly straight. As soon as the tension is relaxed through the more complete drying of these cells the elasticity of the thick walls causes the capsule to snap back to near its original position with a quick jerk, thus throwing out the spores to a considerable distance. This motion of the capsule can be approximated by mounting sporangia that have been soaked in a drop of water and watching them under a microscope while a drop of glycerine placed in touch with one side of the mount is drawn in with a filter paper applied to the opposite side. The denser glycerine will draw the water out of the cells by osmosis and set up a tension in the cells of the annulus just as did the evaporation of the water.

The form of the sorus and its relation to the indusium and particularly the structure of the annulus are important characteristics to be observed in identifying the ferns. In the flowering ferns, *Osmunda,* the large slightly-stalked sporangia are provided with a rudimentary annulus of a few thickened cells located at one side of the capsule (Fig. 224, *A*). The entire leaf or certain leaflets are completely covered with the sporangia, little more than the vascular bundles remaining (Fig. 225, *A*). *Osmunda* belongs to the lowest group of the Filicales and shows certain analogies in the arrangement and structure of its col-

lateral vascular bundles with the Ophioglossales that suggest the derivation of the Filicales from this source. The majority of ferns in temperate climates are characterized by sporangia of the type shown in Fig. 224, *B*, and they are distinguished by the form and arrangement of the sori and indusia. The shield fern, *Dryopteris*, has a rather circular or curved sorus covered by an indusium that is laterally attached at a single point (Fig. 222, *A*). The Christmas fern, *Polystichum*, has a circular sorus with centrally attached indusium (Fig. 225, *C*). The spleenwort

FIG. 225. Common forms of the Filicales: *A*, the flowering fern, *Osmunda*, showing below two green leaflets and above two sporangia-bearing leaflets. At left a cluster of sporangia magnified. The first leaves in the spring only bear sporangia; those appearing later have only green leaves. *B*, chain fern, *Woodwardia*. *C*, Christmas fern, *Polystichum*. *D*, bladder fern, *Felix*. *E*, hay-scented fern, *Dennstaedtia*—*s*, sorus enlarged. *F*, *Woodsia*.—After Sprague.

fern, *Asplenium*, is characterized by elongated sori arranged obliquely to the midrib upon the upper side of the veinlets. The indusium is attached on one side of the sorus along its entire length (Fig. 222, *B*). The chain fern, *Woodwardia*, differs

from *Asplenium* in having the sori arranged in chain-like rows parallel to its midrib (Fig. 225, *B*). In several genera of ferns the indusium is partly or entirely inferior. Thus in the bladder fern, *Filix,* the partly inferior indusium covers the circular sorus like a hood (Fig. 225, *D*), while it is wholly inferior in the hay-scented fern, *Dennstaedtia,* forming a cup (Fig. 225, *E*) and in *Woodsia* the indusium is roundish or star-like (Fig. 225, *F*). Several ferns are distinguished by false indusia that are formed by the more or less modified margins of the leaf. In the bracken, *Pteridium,* the entire membranous margin of the leaf curves over the closely crowded sori (Fig. 226, *A*), and in

FIG. 226. Ferns without indusia or possessing false ones: *A,* leaflet of bracken fern, *Pteridium. B,* maiden-hair fern, *Adiantum. C,* polypod fern, *Polypodium.*—After Sprague.

the maiden-hair fern, *Adiantum,* the sporangia are at the ends of the veins and covered by reflexed portions of the leaf (Fig.

226, *B*). The indusia are lacking in some forms, as in the beech fern, *Phegoteris,* and in the polypody, *Polypodium* (Fig. 226, *C*). These genera are distinguished by the fact, among others, that the leaves of *Polypodium* drop off, leaving a scar as in our deciduous trees. The sporophyll of the sensitive fern, *Onoclea,* bears a striking outward resemblance to *Botrychium,* but the sporangia-like bodies are really leaf bodies rolled up and each bears several round sori on its inner side (Fig. 227). In this



FIG. 227. The sensitive fern, *Onoclea: A,* portion of normal green leaf. *B,* a spore-bearing leaf. *C,* two views of one of the round lobes of *B,* showing the sori on inner side of the lobe.—After Bailey.

and several other genera, see *Osmunda,* the work of photosynthesis is given over to large green leaves that do not produce sporangia.

(*b*) *The Gametophyte.*—In the majority of ferns the spore germinates by rupturing the outer coats and producing a germ tube from which one or more delicate rhizoids are cut off. The germ tube elongates, forming a short chain of cells which soon develop by apical growth into a flat thalloid structure, commonly called the prothallium, that is attached to the ground by numerous rhizoids (Fig. 228). This gametophyte often becomes heart-shaped (Fig. 228, *C*), owing to the more rapid growth of the cells that are cut off from the apical cell. In some genera branched filamentous or narrowly thalloid growths are developed that resemble the protonema of the mosses or the thallose hepatics. The gametophyte usually lives but a few months, although

in some species they may endure for years, multiplying extensively by gemmae, and so form conspicuous mats upon the moist trunks and rocks. The archegonia and antheridia are usually borne upon the same gametophyte. Some genera, however, are strictly dioecious, producing small antheridial or male gametophytes and larger archegonial or female gametophytes. Small male gametophytes occur not uncommonly among any of the genera, owing doubtless to their poorer nourishment. The sexual organs are developed upon the under side of the gametophyte (Fig. 228, *C*), probably because this position is of advantage in keeping them in contact with any water that may fall upon the



FIG. 228.



FIG. 229.

FIG. 228. Gametophyte of the Filicales: *A*, germination of the spore. *B*, early appearance of the thalloid structure of the gametophyte owing to the formation of an apical cell, *x*. *C*, mature gametophyte—*an*, antheridia; *ar*, archegonia; *r*, rhizoids; *v*, apical cell or growing point.

FIG. 229. Structure of reproductive organs: *A*, antheridium as seen in section, just before the discharge of the gametes. *B*, male gamete. *C*, sectional view of archegonium—*g*, female gamete.

earth and thus preventing their drying out. The antheridia develop before the archegonia, often appearing while the gametophyte is quite small. They usually project from the surface of the thallus as spherical bodies covered with a single layer of chlorophyll-bearing cells which inclose usually from thirty-two to sixty-four gamete-bearing cells (Fig. 229, *A*). The male gametes are discharged as in the Bryophyta—the swelling of the antheridium causes the rupture or throwing off of the apical cell and the extrusion of the inner cells as a mucilaginous mass. The gametes are large spirally-coiled bodies provided with numerous cilia and the larger posterior coils enclose a delicate sac containing the remains of the nourishment stored in the mother cell (Fig. 229, *B*). The archegonia are developed on the older prothallia, usually just back of the growing joint (Fig. 228, *C*). The neck of the archegonia, consisting of four rows of cells, project from the under surface of the prothallium and is usually curved back from the growing point, while the basal portion containing the female gamete remains buried in the tissues of the prothallium (Fig. 229, *C*). When the female gamete is mature the canal cells become mucilaginous and effect the opening of the archegonium as in the mosses.

(*c*) *Germination of the Gametospore.*—The gametospore germinates in a week or more after fertilization, dividing very regularly first by a vertical and later by a transverse division into four cells. Each of these cells form a primary organ of the young sporophyte. Of the two outer cells, the one nearest to the growing point of the prothallium (Fig. 230, *A, c*) by repeated divisions produces the first leaf or cotyledon, while the other one forms the first root (Fig. 230, *A, r*). Of the two lower cells. the one directly below the leaf cell gives rise to the stem (Fig. 230, *A, s*) and the other one to the foot (Fig. 230, *A, f*). The growth of these four cells results at first in the formation of a rather globular sporophyte, but soon the growing point of the young root grows through the calyptra, turns downward and penetrates the soil (Fig. 230, *B*). This is followed by emergence of the cotyledon, which, however, curves upward and spreads out its blade to the light (Fig. 231). The young sporo-

phyte, which has up to this time drawn its nourishment by means of the foot from the gametophyte, is now in a position to care for itself. The prothallium soon withers, leaving the sporophyte



FIG. 230. Germination of the gametospore: *A*, section of archegonium after fertilization, showing the four-celled stage of the germinating gametospore. The cell *r* by repeated division forms the first root, *c* forms the first leaf, *s* forms the stem and *f* the foot; *x*, apical cell of prothallium. *B*, later stage. The young sporophyte rupturing the archegonium or calyptra, lettering as in *A*.

an independent and self-supporting plant. The first leaf is small and usually bears little resemblance to those of the mature plant,

FIG. 231. Older sporophyte that has developed two roots, *r*, and is unfolding the second leaf, *l*, but still attached to the withering gametophyte or prothallium, *p*; *rh*, rhizoids.

but as the stem elongates, new leaves are formed which gradually become larger and each succeeding one resembles more closely the adult form. The foot disappears with the withering of the gametophyte, and this is soon the fate of the primary root, but numerous secondary roots are formed along the stems as it continues to creep along on or near the surface of the soil. This growth goes on from two to several years before the plant is prepared to develop sporangia upon the leaves, page 254.

Order 3. Equisetales. The Horsetail Ferns

114. General Characters.—This small group of ferns, comprising but a single genus of about twenty-five species, is but a remnant of an extensive group of plants that flourished in the coal period and that formed conspicuous features of the vegetation at that time. The structure of these earlier forms has been so perfectly preserved in fossil remains that they give a better idea of the relationship of the group than could be obtained from the living plants. Forms allied to *Equisetum* doubtless formed during the coal period of the earth large forests and attained a height of sixty to ninety feet and perhaps three feet in diameter. The species that survive to-day are rush-like plants that rarely exceed a foot in height, though a single tropical form supports its delicate stem upon other vegetation and so attains a length of over thirty-five feet. Species of *Equisetum* are of common occurrence in nearly all countries, living in shallow ponds, swamps, and marshes or drier soils.

At first sight they show little suggestion of fern relationship (Fig. 232) and impress one as being singularly out of harmony with our common plants, as though they were indeed relics of a past age. The large leaves, which were so characteristic of the Filicales, are reduced to minute papery scales in this order and take little or no part in photosynthesis. These scale leaves are arranged with great regularity at the nodes of the stem, and owing to their close association they grow together, forming a papery sheath with teeth-like points about the stem (Fig. 232, *l*). It is noteworthy that in allied fossil forms large chlorophyll-bearing leaves occurred.

The stems are of two kinds, subterranean rhizomes that branch extensively through the soil, and aërial stems that arise as branches from the rhizomes. The aërial stem is simple or branched and is characterized by nodes made conspicuous by the



FIG. 232. A common horsetail fern, *Equisetum arvense: a,* the green branching plant that lives through the summer—*l,* scale leaves; *r,* rhizome or underground stem with tuberous storage organs. *b,* early spring shoot that bears a spike or strobilus of modified spore-bearing leaves, *sp.* This stem is of a light brown color and withers after the spores are shed.—H. O. Hanson.

sheathing teeth-like leaves and strongly furrowed internodes. In such species as branch, these organs originate with great regularity in the axils of the leaves and perforating the sheathing leaves

produce a bushy symmetry that caused the name of *Equisetum* or horsetail to be applied to these plants. The epidermal cells are hard and rough, owing to the abundant deposit of silica in the cell walls. For this reason certain species were used in early times for scouring purposes and so they became popularly known as scouring rushes. Complete silicious casts of the epidermal walls and stomata can be obtained by treating the tissue as noted in the diatoms. The arrangement of the leaves and branches and the distribution of the vascular bundles and other tissues presents a mathematical regularity unexcelled in any plant. As in the Bryophyta, the elongation of the stem is effected by a single apical cell which cuts off with extreme regularity the cells from which the various tissues and organs of the plant are formed. The vascular bundles, which are of the collateral type noted in



FIG. 233. Cross-section of a portion of the stem showing its grooved character and stereome confined to the ridges: *a*, air spaces; *e*, endodermis, inside of which are shown three collateral bundles.

the Ophioglossales and *Osmunda,* are arranged around a large pith (Fig. 233). They rarely show secondary growth through the activities of a cambium, although this was a marked feature of many of the extinct species. These bundles lie directly below

each ridge of the stem and run downward through the internode to the next lower node, where they divide into two equal parts. One of these branches joins a similar branch from the adjacent bundle and continues straight down to the next node, where the branching is repeated. The rudimentary bundles of the leaves also join on to the bundles of the stem at the node. Owing to this regularity of branching, the leaves and vascular bundles alternate in each succeeding node and the vascular system assumes a cylindrical form composed of oblong six-sided figures. Thick-walled stereome cells are developed in the ridges and constitute the principal mechanical tissue of the stem, while the stomata and chlorophyll apparatus are largely confined to the grooves. The stems are very light, owing to absorption of the larger part of the pith and certain regions of the cortex (Fig. 233, *a*).

(*a*) *The Spore-bearing Leaves or Sporophylls.*—The sporangia are only formed upon special leaves, sporophylls, that are arranged in a compact cone or strobilus (plu. strobili) at the end of the stem (Fig. 232, *sp*). In some species the strobilus is borne upon a special branch that withers away as soon as the spores are shed (Fig. 232, *b*) and in other cases the strobilus appears at the tip of the ordinary green shoot. The development of the sporangia upon specialized sporophylls that are grouped together at definite points on the plant is one of the significant departures that appears in the evolution of the plant. This distribution of labor that is seen in the setting aside of certain groups of leaves for spore production will appear in all the succeeding groups; and the arrangement and structure of the sporophylls will steadily become more and more complex until a point is reached where they are popularly called a flower, although this term can be just as correctly applied to the strobilus appearing in *Equisetum* and succeeding groups. The sporophylls of *Equisetum* originate at the nodes, as in the case of the scale leaves, but as they enlarge the apical portion spreads out like a shield. The internodes do not elongate and separate the sporophylls to any considerable extent, consequently the shields become six-sided through contact with the adjacent sporophylls.

(*b*) *Character of Sporangia and Spores.*—The sporangia ap-

pear as rather elongated sacs on the under side of the shields (Fig. 234, 2) and at maturity open by a longitudinal cleft. The structure of the spores is of especial interest. The outer wall of these spores is thickened in spiral bands, and owing to the dissolution of the thin portion of the wall separating these bands, the outer coat uncoils and appears as four bands, with spoon-like ends, attached to the spore at one point (Fig. 234, 3). These



Fig. 234.

Fig. 235.

Fig. 234. Sporophylls and spores of *Equisetum:* 2, sporophylls viewed from outer and inner side, showing form and attachment of sporangia and the central stalk attaching the sporophyll to the strobilus. 3, spores with elaters expanded in *a* and partially coiled in *b*.

Fig. 235. Female gametophyte of *Equisetum* bearing several archegonia and leaf-like lobes. At right male gamete.—After Sadebeck.

bands or elaters are very hygroscopic and their movements assist in rupturing the sporangium, but their special significance is seen in the fact that the elaters become entangled and so several spores are carried away together by the wind. The meaning of this arrangement will appear directly.

(*c*) *Germination of the Spore.*—The gametophyte produced from these spores (Fig. 235) is an irregularly lobed thallus more suggestive of the irregular thallus of an hepatic or the leaves of a moss plant, or the lobed gametophyte of certain species of the Ophioglossales than of the prothallium of the Filicales. The most important feature about the gametophytes is the fact that they are as a rule dioecious and of two sizes, the smaller ones bearing only antheridia and the larger only archegonia. This difference in the nature of the gametophytes is largely due to

nutrition as noted among the Filicales, the well-nourished ones being female. In fact antheridia may appear upon the female gametophyte as a result of insufficient nourishment during its later development. Thus we see that the germination of these spores, which are apparently exactly alike, is controlled by a definite stimulus, just as was the case in the formation of zoöspores and gametes among the lower green algae (page 165). It is noteworthy that some of the extinct species of this group actually stored more food in certain spores than in others, so that they came to differ somewhat in size. This habit is well established in some of the living Filicales and will also be noted in the next order. As a consequence of this tendency, the nature of the gametophyte developed from the spore is no longer a matter of chance. The larger spores, called megaspores, by reason of the more abundant food produce female gametophytes, while the smaller spores, microspores, form small gametophytes bearing only antheridia. The significance of this tendency to produce megaspores and microspores will be seen in the discussion of the fourth order, but attention is called to it here because the Equisetales show very clearly how such a condition came about. It would appear probable that the spores of *Equisetum,* although exactly alike as far as we can see, must have already undergone some physiological change which predisposes them to develop either male or female gametophytes. If this is not the case it would be difficult to explain the common occurrence of dioecious gametophytes in *Equisetum* and the rarity of such an occurrence among the Filicales. It would appear reasonable to suppose that the living substance in the spore of the Filicales is not so highly organized and therefore not so readily influenced by external conditions, while the composition of the spore of *Equisetum* is of such a nature that the amount of food placed at its disposal profoundly effects its germination and development. Perhaps this physiological differentiation of the spores lead to their appropriating different amounts of food during their formation in the sporangium, and so they finally came to be distinguishable as large and small spores, as noted above. The antheridia and archegonia present essentially the same features as were noted

in the lower ferns and Filicales, and fertilization is effected in the same manner.

(*d*) *The Germination of the Gametospore.*—The most noteworthy departure in the germination of the gametospore is seen in the limited growth of the stem. The early stages are similar to those of the Filicales, but after the stem has formed a few nodes with three leaves each, it is replaced by a stem that develops at its base. This second stem attains a somewhat larger size, but is finally replaced in the same manner by a third shoot. In this way several stems are formed until finally one is developed that penetrates the ground and forms the characteristic rhizome of the mature sporophyte. These plants with their scale leaves reduced to protective organs, sunken stomata and chlorenchyma confined to the grooves of the internodes, present an extreme form of xerophytic structure and form a sharp contrast with the large and usually thin-leaved Filicales. While they occur in dry localities, to which conditions their structures admirably adapt them, they are of more common occurrence in moist and wet places. This peculiar distribution of the species of *Equisetum* has not been explained. It is evident that these plants, like the rushes and sedges of our marshes and shallow ponds, are often exposed to intense heat and light, which would cause an excessive transpiration. Possibly these plants are not able to absorb water rapidly owing to the limited amount of conducting tissues and to the exclusion of the atmosphere by the water which surrounds the roots (see pages 40, 46), consequently they are at the same disadvantage and require the same protective devices as plants living in arid localities.

Order 4. Lycopodiales. The Club Moss Ferns

115. General Characters.—The members of this group are popularly known as the club moss ferns owing to their small moss-like leaves and the arrangement of the spore-bearing leaves into club-like strobili or cones (Fig. 236). The gametophyte or sexual generation is also suggestive of the mosses. Particularly is this true of the male gamete, which is small and apparently biciliate (Fig. 240).

The Lycopodiales are largely tropical, though a small number of forms are of common occurrence in temperate regions. The stems are erect or creeping and rather small, but owing to their

FIG. 236. A common club moss fern, *Lycopodium annotinum,* with creeping stem and erect branches covered with small moss-like leaves: *s,* strobilus; *r,* roots.—H. O. Hanson.

prolonged growth and extensive branching they often form conspicuous and attractive colonies. The tissues of the stem do not materially differ from those of the common fern, though the vascular bundles are centrally placed and often appear as radiating plates (Fig. 237). The work performed by the leaves, as in the Equisetales, is usually of two kinds, namely, photosynthesis by the green foliage leaves and spore production by the leaves that usually form strobili at the tips of certain branches. But a single sporangium is associated with each of these sporophylls (Fig. 239).

The club moss ferns are a much larger group than the Equisetales, but like the latter group, they are a remnant of a highly developed and widely distributed race. Fossil remains indicate that the ancient allies of these plants were conspicuous features

FIG. 237. Cross-section of a portion of the stem of *Lycopodium,* showing the centrally arranged vascular bundles.

of an earlier vegetation, with palm-like trunks 100 feet in height and three feet in diameter, and bearing a crown of long narrow leaves that attained a length of three feet. They reached their greatest abundance in the coal age and thence gradually declined, being crowded out by the more specialized seed plants. There are two important families of the Lycopodiales: 1, Lycopodiaceae; 2, Selaginellaceae.

116. Family 1. Lycopodiaceae.—With but one exception the members of this family belong to the genus *Lycopodium,* commonly known as the club moss fern, ground or running pine or ground fir, etc. (Fig. 236). These plants are very well represented in our open woods, where they often form conspicuous colonies owing to the extensive branching and prolonged growth of the stems which creep over or through the ground, sending

up numerous, erect branches and giving off roots that branch with great regularity. Various species of *Lycopodium* are extensively gathered for decorations since the aërial portions of the stems are thickly clothed with small moss-like leaves. The second genus of the family (*Phylloglossum*) contains but one

Fig. 238.

Fig. 239.

Fig. 238. *Phylloglossum Drummondi.*—After Pritzel.

Fig. 239. Strobilus and sporophylls of *Lycopodium:* 2, strobilus. 3, a leaf or sporophyll enlarged from the strobilus with attached sporangium. 4, a spore greatly magnified.

species, which is found in New Zealand and Australia. It is of interest because it is the most simple fern known (Fig. 238), consisting of a few narrow leaves, rudimentary strobilus and poorly developed tissues.

The sporangia of the Lycopodiaceae are large considering the size of the leaf, and usually appear on specialized leaves that are arranged in strobili on the ends of erect branches (Fig. 239). The sporangia open by a longitudinal cleft and the spores are produced in such quantities in some of the forms that they are of commercial importance, being used for flashlight effects and sold as lycopodium powder.

(*a*) *The Gametophyte of the Lycopodiaceae.*—The most interesting feature about the club moss ferns is the sexual generation, because it indicates that these ferns may have been derived from the same ancestry as the mosses or at least from a very primitive stock. In certain species the spores produce a rather cylindrical erect gametophyte which terminates in a number of radially-arranged green leaf-like lobes, among which the archegonia and antheridia are produced (Fig. 240). This structure



Fig. 240. Gametophyte of *Lycopodium* as seen in section: *an,* antheridia; *ar,* archegonia. The young sporophyte or embryo developed in one of the archegonia consists of a foot, *f;* a root, *r;* and a stem, *st,* bearing a leaf on either side; *s,* the suspensor, which has pushed the sporophyte early in its development into the tissues of the gametophyte. Above a single male gamete is figured.—After Bruchmann.

has been compared to a miniature leafy moss plant. This suggestion of relationship is strengthened by the resemblance of the male gametes to those of the mosses, the sperms, so far as they have been observed, being small, almost straight bodies, and provided with two cilia (Fig. 240). The archegonia may also be more primitive than those of the ferns in possessing several neck canal cells and five instead of four rows of neck cells. The gametophyte appears in all species examined to be associated with symbiotic fungi, as in Ophioglossales, and this has led in many forms to its subterranean and chlorophylless development.

(*b*) *The Germination of the Gametospore.*—The germination of the gametospore is characterized by new departures that will

continue on through the seed plants. The gametospore enlarges and divides into two cells by a transverse wall. The outer of the two daughter cells takes no part in the formation of the sporophyte, though elongating and often dividing several times. These cells are termed the suspensor (Fig. 240, *s*) and function in pushing the lower daughter cell down into the nourishing tissue of the gametophyte. The lower daughter cell, by a series of divisions that at first resemble the common ferns, forms the young sporophyte. This consists of a stem with one or two cotyledons, a massive foot and finally, at a late period in the development of the sporophyte, of a root (Fig. 240). The tardy development of the root has been cited as an indication of the origin of the lycopods from very primitive ancestors, in which the formation of the root had not become established. It may also be due to the abundant food stored in the gametophyte and hence the development of the root might well be delayed until this store is exhausted. The growth of the sporophyte is very slow and it remains as a parasite upon the gametophyte for a long time, even for years in some of the subterranean forms (see Ophioglossales). The elongation of the stem and root is effected by the division of several cells rather than by one apical cell, as in previous cases, a feature to be noted in the following division.

117. **Family 2. Selaginellaceae.**—This family includes but a single genus, *Selaginella,* of over 600 species. Only a few forms occur in the temperate regions, the majority being confined to tropical countries, where they often form one of the most attractive features of the forest vegetation owing to the symmetry of their branching and the rare delicacy of their foliage (Fig. 241). For these reasons they are extensively cultivated and familiar objects in conservatories and florists' shops. The so-called resurrection plant, *Selaginella lepidophylla,* lives in the very arid sections of the southwestern United States, and during drought reduces its surface to a nest-like ball by rolling up its branches into tight coils. In this condition it appears as a brownish dead mass. When moistened, the absorption of water causes the branches to quickly uncoil and also renders the tissues translu-

cent, so that the green color of the chloroplasts can be seen. These reactions occur even in the dead plants and so create the impression that they have returned to life. Many of the species of *Selaginella* have a creeping habit like that of *Lycopodium,*



FIG. 241. A common cultivated *Selaginella:* 1, habit of the plant—*s,* strobilus; *b,* branch-bearing roots. 1*A,* portion of stem, showing leaf arrangement.—H. O. Hanson.

and the aërial stems are covered with leaves arranged in several rows, very frequently in four rows of two large and two smaller leaves (Fig. 241, 1*A*). Usually but a single chloroplast appears in each cell, as already noticed in several algae and in the Anthocerotales. The roots are frequently developed from the end of naked branches that extend from the stems to the ground (Fig. 241, *r*).

(*a*) *Sporangia and Spores.*—The sporangia are borne on the stems in the axils of the leaves which form a strobilus, as in *Lycopodium.* However, they differ radically from those of the

Lycopodium, in that two kinds of spores are formed, small spores or microspores and larger ones or megaspores. The formation of two kinds of spores, or heterospory, is not confined to the Lycopodiales. It occurs among certain genera of the Filicales and, as has been stated, characterizes some of the fossil Equisetales. The megaspores are generally formed at the base of the strobilus and the microspores occupy the upper sporangia (Fig. 242). The sporangia are called microsporangia and megaspor-



FIG. 242. Sporophylls and spores of *Selaginella: 2*, strobilus with lower sporophylls separated for spore dissemination. 3, megasporophyll with sporangium containing four megaspores. 4, microsporophyll. 5, megaspore enlarged. 6, microscope equally magnified. 6*A*, more enlarged view of the microspore. The triangular surfaces of the spores show that these spores have been formed in tetrads from a spore mother cell as in previous groups.

angia accordingly as they contain small or large spores and for the same reason the leaves may be designated as micro- and mega-sporophylls. The two kinds of spores originate in the same manner as previously noted and the difference in size is due to the amount of food which they receive. In the case of the microspores, the numerous mother cells of the sporangia form four spores each in the usual manner; but in the megasporangia only one of the mother cells divides in this manner, and in some cases only two spores are formed. The other mother cells do

not develop and are ultimately consumed in nourishing the megaspores. As a result of the large amount of food transferred to the megasporangium, both it and especially the two or four spores become much larger than the others (Fig. 242, 3–6).

(*b*) *The Germination of the Spores.*—The most important and suggestive feature in the life history of *Selaginella* appears in the germination of the spores. The spores germinate in the sporangia and not after being shed, as in other ferns. The microspore forms two cells while in the sporangia, a small one



Fig. 243. Gametophyte and young sporophyte of *Selaginella:* 7, section of a microspore that has nearly completed its germination—*g*, a single cell that is possibly a remnant of the large gametophyte or prothallium of previous groups of ferns; *an*, antheridium consisting of a layer of wall cells which enclose the gamete mother cells. 7*A*, a male gamete. 8, section of a megaspore, showing the stage of germination that is usually attained in the sporangium. 9, mature female gametophyte that has ruptured the wall of the megaspore, thus exposing the archegonia, *ar*, one of which has developed a young sporophyte or embryo with root, *r;* stem bearing two leaves, *st;* foot, *f*, and the suspensor, *s*. 10, young sporophyte with root, stem and leaves emerging from the gametophyte.

often compared to a remnant of the male gametophyte or prothallium and a large cell which ultimately forms a single antheridium. At about this stage of germination the microsporangium opens by a vertical cleft, permitting the scattering of the

spores. The larger cell of the microspore now forms a central mass of gamete mother cells which are surrounded by a single layer of wall cells (Fig. 243, 7). The male gametes are of the same character as noted in the *Lycopodium* and are set free by the disintegration of the wall cells. The megaspores begin to germinate even before they have reached their full size. The nucleus of a megaspore divides repeatedly, forming numerous nuclei which become arranged about the walls of the young spore. Subsequently walls are formed about the nuclei at the apex of the spore (Fig. 243, 8) and by further division a mass of cells results, in the outer part of which the first divisions of the archegonia arise. At this stage of development, or, earlier, the megasporangium opens by a vertical cleft permitting the discharge of the spores. It must be borne in mind that this origin of the gametophyte is of a radically different nature from any case previously noted. Heretofore the gametophyte always had an independent existence and the sporophyte for varying periods of time was a parasite upon it. In *Selaginella* you note the beginning of a reversal of this relation, for the sporangia remain green and continue to nourish the spores during their germination; in other words, during the early stages in the development of the gametophyte. We see that the gametophyte is becoming parasitic upon the sporophyte (see page 252). After the spores are shed they complete the growth of the gametophyte by forming the archegonia and filling the space within the spores with a solid mass of cells. This growth ruptures the spore walls at the apical regions, thus exposing the archegonia to the male gametes (Fig. 243, 8). The archegonia are rudimentary structures consisting of but two neck cells, and a single canal cell leads to the female gamete.

It is evident that these variations are of great advantage in ensuring the perpetuation of the species. Especially is the nourishment of the young gametophyte by the highly organized sporophyte a distinct gain. In previous types the formation of the gametophyte was dependent upon favorable conditions, such as moisture, light, temperature, etc., and it should be said that only in rare cases do the spores fall in such favorable situations and

meet with such suitable conditions as to enable them to mature archegonia and antheridia, and so provide for a new sporophyte. In *Selaginella* the formation of the gametophyte is ensured by the food supplied to it by the sporophyte. Thus the germination of the spore is practically independent of its surroundings and it can complete its growth under a variety of conditions, even in the dark. It should also be noted that this relationship of the two generations results in a marked reduction in the size of both the male and female gametophyte, since they are no longer burdened with the work of food construction. This reduction will proceed very rapidly as we advance through the remaining groups.

(*c*) *Germination of the Gametospore.*—The germination of the gametospore is very much like that of the *Lycopodium*. The suspensor is much longer and the young sporophyte consists of a stem bearing two cotyledons, a well-developed foot and a root that is developed after the other organs (Fig. 243, 8). The foot continues to absorb food from the gametophyte after the stem and root have emerged, and in this condition the relation of the sporophyte to the gametophyte is strikingly suggestive of a sprouting seed (Fig. 243, 10). In this connection it should be remembered that the young sporophyte of certain species remains in the gametophyte during the winter or during a drought in a resting or dormant condition, thus resembling very closely the seed structure to be seen in the next group. It may also be stated that fertilization and even the formation of the young sporophyte may begin before the spores are discharged from the sporangium, thus helping us to understand how it came about that the megaspores were permanently retained in the sporangium, as is the case among seed plants. In one of the large groups of ferns that have become extinct, it is noteworthy that both the gametophyte and young sporophyte were retained in the sporangium so that the seed-like structures were formed.

CHAPTER IX

DIVISION IV. SPERMATOPHYTA OR SEED PLANT

118. General Characters.—This is the largest group of the vegetable kingdom and includes all our common trees, shrubs and herbs. The relationship of these plants is very imperfectly understood and the various classes, orders, etc., into which they are divided are in part artificial and do not therefore represent completely the alliances of the groups. You have noticed that the sporophyte is the most important feature in the life history of the fern and that the gametophyte, subordinate from the start, becomes very inconspicuous in the higher forms. This inequality of the two generations is more noticeable in the seed plants where a progressive series of variations result in a more and more complex external and internal differentiation of the various organs of the sporophyte and in a steady reduction of the gametophyte. Heterospory that appeared in several groups of ferns, becomes a constant characteristic of the seed plants. The spores originate much as in the ferns but the megaspore is nourished and permanently retained in the sporangium where it not only forms the female gametophyte (see *Selaginella*) but also develops the young sporophyte. At this point, the most characteristic feature of the Spermatophyta is seen. The young sporophyte or embryo ceases to grow while still in the gametophyte and passes into a resting or dormant condition and the sporangium which has become variously modified for the purpose of protecting the embryo is discharged from the sporophyte. This sporangium containing the embryo is called the seed. Contrast with these features two important differences that appeared in *Selaginella*. First: The spores of these ferns are retained and nourished in the sporangia only during the early development of the gametophytes. Second: The young sporophyte of *Selaginella*—it may also be called the embryo—does not normally pass into a resting state, but steadily grows on into the mature sporo-

phyte. The seed plants are divided into two subdivisions based upon the relation of the megasporangia to the sporophylls: 1. Gymnospermae, with sporangia, hence with the seed on the surface of the sporophyll. 2. Angiospermae, with sporangia, hence with the seed, enclosed by the sporophyll.

Subdivision 1. Gymnospermae. Plants with Naked Seeds

119. Origin of the Gymnospermae.—This group is principally represented today by the cone-bearing trees. In very ancient geological times, several groups of primitive gymnosperms, now extinct, flourished and formed an extensive and varied forest vegetation. Fossil remains of these various groups have been wonderfully well preserved and show that they were undoubtedly related to the ferns. The modern group of gymnosperms have been derived from these ancient forms and but a remnant have survived the changes occurring upon the earth and the competition with the more highly specialized Angiospermae. A few genera, however, are well represented today and they are of great commercial importance as lumber. The sporophylls of the gymnosperms are arranged in a strobilus as in the Equisetales and Lycopodiales, but the microspores and megaspores are not associated, being developed in separate strobili. Attention will be directed to only two of the more important orders: The Cycadales or Cycads and the Pinales or cone-bearing trees.

120. Order a. Cycadales or Cycads.—These plants are strictly tropical, though two genera are subtropical, *Zamia* or coontie in Florida, and *Cycas,* often called the sago palm, in China and Japan. Many of them are extensively cultivated in green-houses owing to their peculiar growth and attractive foliage (Fig. 244). The cycads include less than 100 species of what was in geological times, a very extensive alliance. They are of special interest as showing unmistakable fern characters and because they are the most primitive of our extant seed plants. The stems of these plants are rather tuberous, though certain species are decidedly palm-like in appearance, with stems 12 to 60 feet high. The vascular bundles are collateral and arranged around a large

pith as noted in *Botrychium*. A slight enlargement of the stem is brought about owing to the weak growth of the cambium of these bundles, but the principal increase is effected by the formation of new bundles outside of those first formed. Concentric bundles also occur in the leaves and portions of the stem of certain species. The foliage leaves are large and leathery and form a rosette, alternating with scale leaves, at the apex of the stem. The bases of these leaves form an armor-like plate over the surface of the stems. In certain genera, the young leaves are coiled as in the ferns.

(*a*) *The Sporophylls and Sporangia of the Cycads.*—The sporangia are borne on more or less modified leaves arranged in large terminal strobili and either scattered over the sporophyll or sometimes arranged in groups suggestive of the sori of the ferns, in some forms even showing a rudimentary annulus. In

FIG. 244. *Zamia,* a cycad common in southern Florida, with strobilus of megasporophylls.—H. O. Hanson.

origin and development, they are strikingly like the lower ferns. There is a considerable variation in the form of the sporophylls and the distribution of the sporangia. For example, in *Cycas,* the megasporophylls are loosely associated and only slightly

modified, the sporangia being developed on their margins (Fig. 245, 3*B*). The microsporophylls are small and more compactly arranged, the sporangia being associated in sori on the lower



Fig. 245. Sporophylls and sporangia of the Cycads: 2, strobilus of *Zamia*. 2*A*, cross-section of strobilus of *Zamia*, showing arrangement of microsporophylls. 2*B*, microsporophyll enlarged, showing sporangia arranged in sori. 2*C*, sorus of three sporangia that have opened. 2*D*, microsporophyll of *Cycas*. 3, cross-section of a strobilus of *Zamia*, showing arrangement of megasporophylls. 3*A*, megasporophyll enlarged with two sporangia. The one on the right shown in section: *ms*, megaspore; *i*, integument; *sp*, sporangium; *m*, micropyle. 3*B*, megasporophyll of *Cycas* with laterally arranged megasporangia.—H. O. Hanson.

surface of the sporophylls (Fig. 245, 2*D*). In *Zamia*, the strobili and sporophylls are quite suggestive of *Equisetum*, the sporangia being developed on the inner side of shield-like sporophylls (Fig. 245, 2–3*A*).

The microspores originate and are discharged from the sporangium very much as in the lower ferns. The megasporangia, however, present several new departures that must be borne in mind. A rather thick coat or integument covers the sporangium save for a small opening called the micropyle (Fig. 245, 3*A*). The integument will be a feature in all the succeeding groups and

it has been compared to a highly modified indusium. It originates, however, from the base of the sporangium rather than from the epidermis as in the common forms. As in *Selaginella,* only one of the mother cells takes part in the formation of spores and but one megaspore (instead of four) is formed. This megaspore does not escape from the sporangium, owing to the large number of cells surrounding it and consequently it does not form a thick protective coat. It has been suggested that these cells enveloping the megaspore are due to the transformation of the



FIG. 246. Section of a megasporangium of *Zamia,* showing the female gametophyte just before fertilization: *i,* integument differentiating into a fleshy outer and hard inner coat; *m,* micropyle; *p,* cavity in which three microspores are shown germinating; *g,* female gametophyte which has developed from the single gametospore. At the micropylar end of the gametophyte are two archegonia opening into the archegonial chamber, *ar.* Two neck cells lead to large female gamete, *o; sp,* remains of the sporangium not yet consumed by the growth of the gametophyte. At the right a few of the jacket cells which supply the female gamete with food are enlarged.—After Webber.

spore mother cells, which make up the bulk of the fern sporangium, into nourishing cells. This structure of the megasporangium is characteristic of all the spermatophyta and the arrangement is of decided advantage since it provides the megaspore

during its formation and germination with an abundance of food and moisture.

(*b*) *The Gametophyte of the Cycads.*—The megaspore germinates as in *Selaginella.* Numerous free nuclei are first formed that arrange themselves around the walls of the spore and by the later formation of walls a tissue is developed that gradually fills the enlarging spore. Archegonia now appear at the micropylar end of this gametophyte and become sunken in pits, called the archegonial chamber, owing to a continued growth of the adjacent cells of the gametophyte (Fig. 246). While this female gametophyte is very suggestive of the one noted in *Selaginella* it should be observed that the archegonia are more simple in structure, usually possessing but two neck cells and that the central canal cells of the neck are entirely lacking. The female gamete is also much larger (Fig. 246, *o*) than any heretofore noticed and it is nourished by specialized cells, the so-called jacket cells, that surround it and supply it through minute pores with an abundance of food.

It is evident that the buried position of the archegonia must necessitate some new departure in order to bring the male gametes to the archegonia. The microspores have already begun their germination at the time when they are discharged from the sporangium and being very light, they are carried by the winds to strobili containing megasporangia. When the megasporophylls are mature, they spread apart slightly, permitting the microspores to rattle down to the sporangia where they are caught, at least in some forms, by a mucilaginous secretion from the micropyle. This substance in drying contracts and pulls the spores in large numbers through the micropyle into a chamber formed in the upper part of the sporangium (Fig. 245, *p*). As has been stated, the microspore begins to germinate in its own sporangium and by the time that it reaches the cavity of the megasporangium its nucleus has already divided twice, forming a rudimentary male gametophyte that may be compared with that of *Selaginella* (Fig. 247, *A*); the cell (*g*) representing the small cell, and the cell (*a*) corresponding to the large antheridial cell of the gametophyte of *Selaginella* (Fig. 243, 7), while

the cell (*t*) called the tube cell, represents a new departure in the evolution of the male gametophyte. The tube cell grows out into the tissues of the megasporangium forming a tubular structure and often branching extensively and absorbs food for the nourishment of the antheridial cell (Fig. 247, *B*). This latter cell finally divides, forming a rudimentary antheridium



FIG. 247. Male gametophyte of *Zamia: A,* stage of germination of the microspore attained in the sporangium. See text for explanation of figures. *B,* formation of tube for absorbing of food from megasporangium. *C,* spore end of gametophyte showing the antheridial cell dividing into a body cell, *b,* and a single-wall cell, *w*. *D,* the body cell has divided, forming two cells, which become the male gametes. *E,* spore end of male gametophyte, showing the spirally ciliated male gametes.—After Webber.

consisting of but a single wall cell (*w*) and a body cell (*b*), which by division forms directly two huge male gametes with spirally arranged cilia (Fig. 247, *C–E*). Owing to the growth

of the numerous tube cells, the tissues above the archegonial chamber become disorganized and absorbed so that the end of the gametophyte containing the gametes hangs down into the cavity thus formed and comes into close proximity to the archegonial chamber (Fig. 246). At this stage of development the archegonial chamber has become filled with water. Owing to the swelling of the cells (*g*) and (*w*) assisted also by the accumulation of fluids in the tube, the male gametophyte is ruptured at its basal or spore end and the male gametes are discharged into the archegonial chamber, where they swim about and finally enter the archegonia, as in the case of ferns.

The most noticeable departure in the sexual generation of the cycads is the very considerable reduction of the male gametophyte which consists of a delicate tube containing a few cells from one of which the male gametes are formed directly by division and not as in the ferns, through the reorganization of the contents of the cells into gametes which are set free through the mucilaginous modification of the walls of the mother cells. You also notice that the gametophytes have become entirely parasitic upon the sporophyte, the male gametophyte at first upon the sporophyte which bore it and later upon the megasporangium. Attention may be called to a suggestive departure in the development of the female gametophyte that has been observed in one of the cycads. In case fertilization is not effected, the gametophyte ruptures the sporangium and projects as a green tissue, as in certain heterosporous ferns.

(*c*) *Development of the Sporophyte.*—The germination of the gametospore differs in two important respects from the ferns. Its nucleus gives rise by repeated divisions to numerous nuclei which arrange themselves around the walls of the gametospore, becoming especially numerous at its lower end, and finally form cell walls (Fig. 248, *A*). This structure is called the pro-embryo. From the lower cells of the pro-embryo are organized a massive suspensor (Fig. 248, *B*), which pushes the outermost cells of the pro-embryo deep into the tissues of the gametophyte where they develop the young sporophyte or embryo, which consists of a rudimentary root, stem and two cotyledons (Fig. 248, *C*). In

the ferns the gametospore divided into two cells which by further division formed the sporophyte. The second departure now appears. The sporophyte or embryo ceases to grow and it is admirably protected during its resting period by the integument which has become differentiated into a stony inner coat and often into a fleshy outer coat. This completed structure, consisting of the modified integument, the embryo, and the gametophyte,



FIG. 248. Development of the sporophyte or embryo of *Cycas: A,* section of archegonium, showing a large number of cells lining the walls of the germinating gametospore. This is the pro-embryo stage. *B,* later stage in which the pro-embryo has developed a suspensor, *s,* which pushes the embryo-forming cells, *e,* into the nourishing tissues of the gametophyte. *C,* still later development, showing the greatly coiled suspensor and the young sporophyte or embryo, *e,* with two cotyledons. *D,* section of a seed of *Zamia: i,* integument which is fleshy without and stone-like within; *m,* micropyle; *g,* gametophyte; *e,* dormant sporophyte or embryo.—*A–C.* —After Treub.

which has increased in size until it has absorbed all the tissues of the sporangium, we call the seed (Fig. 248, *D*). At this stage of development, the seed falls from the strobilus and may remain in a dormant condition for years.

The formation of the seed is the most characteristic feature of the Spermatophyta and the most important advance that is to be noted in the evolution of plant life. This variation has given the seed plants an advantage over all other terrestrial groups and made them the dominant plants upon the earth. Not

only are the male and female gametophytes better provided for than in *Selaginella,* owing to their complete parasitism on the well developed sporophytes, but the formation of the young sporophyte or embryo is also ensured by the abundance of food placed at its disposal. The most important advantage, however, possessed by the seed plant appears in the modification of the integument which renders it impervious to gases and fluids. This change so effectually seals up the embryo as to protect it for years in some cases against conditions unfavorable for growth. The modification of the integument was doubtless the chief factor that led to the seed habit, since it cuts off the supply of oxygen and fluids from the sporophyte or embryo and thus stops its growth. When conditions are favorable for growth (see page 115), the integument becomes permeable to fluids and gases and the embryo continues its development. The cotyledons remain in the seed to absorb the food stored in the gametophyte, but their basal portions elongate, pushing out the root which soon becomes established in the soil, while the stem which is carried out with the root extends up into the air, and develops the leaves (Fig. 249). This renewal of growth of the



FIG. 249. Renewal of growth of the sporophyte or embryo in the seed: 7, section of the seed, showing the base of the cotyledons extending from the seed, thus pushing out the stem and root, the latter organ (shown in part) curving down into the ground. The free end of the cotyledons are enlarging as they absorb the food stored in the gametophyte. 6, seedling six months old with first normal leaf.—After Sachs.

embryo, or sprouting of the seed, is often erroneously termed the germination of the seed.

The maidenhair tree, *Ginkgo biloba,* a native of China and Japan and now extensively cultivated throughout the world, is the sole survivor of another ancient order of gymnosperms that have many points in common with the cycads and ferns. The tall shaft of the stem, wide-spreading branches and the arrangement and character of the tissues is more suggestive of the next order, the cone-bearing trees. The leaves, however, are fern-like in appearance and are deciduous, unlike most of the Gymnospermae. The arrangement of the sporangia, the development of the gametophytes, and fertilization are of the same nature as in the cycads. It is noteworthy that fertilization may be effected after the megasporangia have fallen, thus approaching the condition seen in *Selaginella.*

121. Order b. Pinales or Cone-bearing Trees.—This order is by far the largest group of gymnosperms and includes our well-known evergreen or cone-bearing trees. They are largely confined to temperate regions and are more numerous in the northern than in the southern hemisphere. The shores of the Pacific are especially favorable for their development and on our western coast they attain dimensions equalled only by the giant *Eucalyptus* of Australia. Though less ancient in origin than the cycads, they have declined in the same way as the latter order and only a remnant of this very extensive group survive today. However, their variations have been more beneficial than in the case of the cycads and as a result nearly 300 species are adapted to present conditions upon the earth and constitute an important part, and commercially the most valuable part of our forest vegetation.

(*a*) *Structural and Adaptive Features of the Pinales.*—One of the variations that is of great benefit to many of these plants is their habit of forming buds (see page 97), which appear in this order in more perfect form than any of the preceding. By means of the scale leaves of the buds all the organs that are to be produced each season are protected against drying winds. The terminal bud of the tree is frequently the most vigorous and as a result spire-like stems with very symmetrically arranged

branches are often formed. The growth of the Pinales is usually prolonged and vigorous, the age of some of the giant trees of California being estimated at over 5,000 years, and they may attain the height of over 300 feet and a diameter of 30 feet. The elongation of the stem is effected by a group of actively dividing cells at the apex, while the increase of girth is brought about by the activities of a cambium, as in the dicotyledons (see page 75). In fact, all the features seen in the cross-section of the stem as the cork, cortex, phloem, cambium, xylem with its annual rings of growth and medullary rays and the pith are suggestive of the dicotyledons, save that the tissues are more simple. This is particularly noticeable in the xylem, which is composed



FIG. 250. *A*, radial section of the xylem of *Pinus*—*t*, tracheids; *p*, border pores or thin places in cell wall to promote transfer of fluids; *m*, medullary ray of two cells accompanied by tracheids, *mt*. *B*, tangential section, showing medullary rays, one cell broad and from three to nine cells high; *p*, bordered pores. *C*, cross-section of the outer cells of pine leaf, showing heavy epidermal cells, *e*; well-developed cuticle, *c*; sunken stoma, *s*; strengthening stereome cells, *st*, and chlorenchyma, *m*.

of very regularly formed tracheids with peculiar bordered pores on their radial walls to promote the rapid transfer of fluids to and from the medullary rays (Fig. 250, *A*, *B*). The characteristic uniform and even-grain of pine, spruce, etc., is due to the simplicity of the tracheids which compose the wood. This struc-

ture of the wood explains the ease with which the timber is worked and the common name of "soft wood" that is applied to it. The leaves are xerophytic in character and usually rather small and often needle-like. The strongly thickened epidermis is strengthened with strands of stereome that give them a leathery character (Fig. 250, *C*). The stomata are sunken and the centrally-placed vascular bundles communicate with the rather irregular chlorenchyma cells by means of tracheids, and thin-walled parenchyma cells instead of small veins as in the higher plants. The leaves in the majority of the genera remain on the branches for several years, and for this reason the members of the Pinales are commonly called "evergreens."

These plants generally flourish in localities where there is an abundance of moisture and at first sight the modifications of the leaves noted above are not in harmony with such surroundings, being decidedly characteristic of plants growing in arid regions. The minute tracheids which constitute the conducting system of the Pinales cannot, however, transport the fluids absorbed from the soil so rapidly as the large cells noted among the dicotyledons. Consequently smaller amounts of water are placed at the disposal of the leaves, and were it not for the fact that they are admirably adapted to lessen transpiration these plants would doubtless have disappeared from the earth long ago. The ability of the leaves to diminish the loss of water may also account for the common occurrence of the cone-bearing trees in northern and mountainous districts where the soils are cold and root absorption is therefore lessened and where the vegetation is especially exposed to the drying winds of winter (see page 34).

Nearly all the Pinales are characterized by the presence of resin passages or ducts in the stems and leaves. Whenever the stems are cut or injured these passages pour out a thick resinous liquid that effectually heals the wound. This doubtless explains in part the freedom of these trees from the attack of insects and wood-destroying fungi. It may possibly account for the inability of many of these trees to sprout from the stump when cut. Hardwood forests will very generally perpetuate themselves by means of sprouts that arise from the stumps, but our evergreen

forests are rapidly disappearing, because with few exceptions the trees are only able to develop from seeds. Possibly the rapid covering of living cells of the stump by the resinous substance excludes the atmosphere so effecutually as to prevent any further growth (page 100). This resinous liquid is obtained in large quantities from certain species of pine by removing the bark and a little of the wood from the side of the trees when the liquid exudes and is caught in a pan placed at the bottom of the cut surface or in a pocket made in the tree. This substance is removed from time to time and distilled, yielding turpentine and resin. Balsam is a resin obtained from the balsam fir (*Abies*).

(*b*) *The Sporangia of the Pinales.*—The sporangia resemble those of the cycads in origin and structure and are developed upon sporophylls that are usually grouped in strobili (Figs. 251,



FIG. 251. Microsporophylls of the pine: 1, branch of pitch pine with leaves spirally arranged in fascicles of three and also with several strobili, *st*. At tip of branch the bud, *b*, is shown elongating and bearing numerous fascicles of leaves that are still enclosed by paper bracts. 2, surface and side view of microsporophyll, showing arrangement of sporangia, *sp*. *2a*, microsporophyll of juniper bearing three sporangia.—H. O. Hanson.

252). The microsporangia are formed on rather small, delicate and often brightly colored sporophylls that perish as soon as the spores are scattered (Fig. 251, 2). The microsporophylls bear one or several sporangia on their under surface and are hygro-

scopic, curving away from each other when the sporangia are ripe, thus permitting the opening of the sporangia and the gradual distribution of the spores by the wind and closing when moistened to protect the spores against wetting. In some of the genera the outer coat of the microspore is lifted away from the inner coat in such a way as to form two sac-like outgrowths which renders them more buoyant and adapted to distribution of the wind (Fig. 254). The microspores are produced in such enormous numbers that the ground in the vicinity of the evergreen forests is often covered with the yellow spores, the so-called "sulphur snow." This extravagant formation of microspores is a necessity owing to the small chance of any one of the microspores being carried to the megasporophylls and so ensuring fertilization. Were it not for the fact that the microspores need be carried only comparatively short distances, this order



FIG. 252. Strobili of megasporophylls: 1, a branch that has developed in the spring from a bud. The fascicles of leaves are just emerging from their papery bracts and at the right is shown a strobilus, *st*, consisting of minute spine-tipped scales. 2, growth of strobilus shown in 1, twelve months later. 3, appearance of strobilus shown in 2, six months later.—H. O. Hanson.

would doubtless become extinct. The gymnosperms, however, live in colonies and the strobili of mega- and micro-sporophylls usually occur upon the same plant, thus necessitating but a short transfer of microspores. The megasporophylls usually endure for a year or more and become quite large, forming the leathery or woody scales of the cone or strobilus (Fig. 252). The megasporangia are usually developed on the upper side of the sporophyll (Fig. 253, *A*), the number formed varying in the different genera. In the case of many of the Pinales, the sporangia appear upon curious outgrowths of the sporophyll which become large and form the conspicuous scales of the cone, while the sporophyll proper appears as a small bract just below it, as in spruces, pines, firs, etc. (Fig. 253, *B*). The origin and structure of the megasporangium and megaspore are essentially as noted in the cycads. Several mother cells may form megaspores, but only one of these megaspores ever develop and germinate.

(*c*) *The Gametophyte of the Pinales and Fertilization.*—The female gametophyte is formed as previously noted in the cycads. By the repeated divisions of the nucleus of the megaspore numerous free nuclei are formed that become arranged around the walls of the megaspore. Later cell walls are developed about these nuclei, and by further division the entire space within the enlarging spore is filled with tissue (Fig. 253, *C*). A few or a large number of archegonia are developed at the micropylar end of the gametophyte. The archegonium consists of two or several neck cells, and a large female gamete which is surrounded by nourishing cells as in the cycads. You notice that there is no chamber formed above the archegonia, and it would be a natural inference that the male gametophyte must be of a somewhat different character from that of the cycads in order to meet this new departure. The microspores which have already begun to germinate when discharged from their sporangia are carried by the wind to the megasporophylls which are slightly spread apart at this time, permitting the microspores to rattle down to the megasporangia. The microspores fall into the micropyle either by reason of the position and construction of the megasporangia, which may be so placed that the spores naturally roll down into

the micropyle, or the microspores may be drawn through the micropyle by mucilaginous excretions as in the cycads. The early stages in the germination of the microspore are essentially as in *Cycas*, though more reduced. The microspore on reaching the

FIG. 253. Megasporangia of pine: *A*, scale-like outgrowth of sporophyll bearing two sporangia. *B*, section of *A*, taken through one of the sporangia, showing the sporophyll, *s*, and large scale-like outgrowth which bears the sporangia, *sp*. The sporangium is surrounded by an integument, *i*, and contains a single megaspore, *m*, as in the Cycads. *mc*, microspores that have been drawn through the micropyle to the sporangium, into which their tube cells are growing. *C*, section of megasporangium enlarged and diagrammatic, showing the female gametophyte at the time of fertilization —*g*, female gametophyte, showing two archegonia with large female gametes, *fg*. At the left the tube of the microspore, *mc*, has entered the neck of the archegonium and the two male gametes have been discharged into the large sac of the female gamete. On the right the tube cell with male gametes is seen approaching the neck of the archegonium. *i*, integument; *sp*, remains of sporangium not disorganized by the gametophyte; *t* shows the disorganization caused by the tube cells as they branch and grow towards the archegonia.

megasporangium has germinated, forming a tube cell and an antheridia cell, but the one or two cells, which correspond to the cell (*g*) of the gametophyte noted in the cycads, are quickly disorganized and appear as faint lines (Fig. 254, 8). These cells quickly disappear because they are no longer of service in setting free the gametes. The tube cell develops as in the cycads, but it is directed in its growth so that one of its branches finally



FIG. 254. Male gametophyte of the pine: 6, section of microspore, showing the two air sacs, *s*, formed by the lifting up of the outer wall of the spore. 8, stage of germination of the spore at time of discharge from its sporangium—*t*, tube cell; *a*, antheridial cell, above which are seen two black lines, the remains of cells formed by earlier division of the nucleus of the microspore. 9, continuation of germination of the microspore after reaching the megasporangium—*t*, tube cell forming a branching tube that disorganizes and absorbs the cells of the megasporangium. The antheridial cell (*a*) of 8 has divided into a wall cell, *w*, and a body cell, *b*. 10, end of tube cell as it approaches the female gamete—*m*, *m*, male gametes. The nuclei of the tube cell and wall cell are also seen (somewhat disorganized) in the end of the tube.—After Coulter and Chamberlain.

reaches one of the archegonia, when it pushes aside the neck cells and fuses with the cell membrane of the large female gamete (Figs. 253, *C*; 255, *A*). In the meantime the antheridial cell has divided into a wall cell and body cell (Fig. 254, 9), and two motionless male gametes are formed from the body cell. It is

now apparent why the tube cell extends to the female gamete and also why the archegonial chamber is not developed. The male gametes and the wall cell, now quite unattached, are carried down to the end of the tube (Fig. 254, 10) by the cytoplasmic currents and pass into the cytoplasm of the female gamete through an opening that is formed in the end of the tube. Chemical attraction now draws one of the male gametes to the nucleus of the female and their fusion results in the formation of the gametospore. The remaining cells of the male gametophyte are apparently disorganized (Fig. 255, *A*). The functions performed by the tube cell in the Pinales will be a constant characteristic of all the members of the next subdivision or Angiospermae, and it should be noted that this important modification of the male gametophyte probably arose at first as an absorbing organ to supply the antheridial cell with food, as is the case in the cycads. In the more advanced types the archegonia become more effectually enclosed in the sporangial tissues and the tube cell assumed in addition the function of a conduit for the male gametes which lost their motility as a result of this new method of transport to the female gamete.

(*d*) *Development of the Sporophyte.*—The germination of the gametospore of the pine will illustrate the more important features in the process. The nucleus of the gametospore divides, forming four nuclei (Fig. 255, *B*), which pass to the upper end of the spore and arrange themselves in a plate. By successive divisions these cells are increased until four plates of four cells each are formed which become surrounded by cell walls save in the case of the lower plate, which is in direct contact with the rich food of the spore (Fig. 255, *C–D*). This meagre growth corresponds to the pro-embryo of *Cycas* (Fig. 248, *A*). In this connection it should be noticed that only a portion of the gametospore, as in the cycads, is used in forming the embryo. In all the preceding groups, as the mosses, ferns, etc., the gametospore in germinating behaves as an ordinary cell, dividing into two cells which continue the process. In the Pinales the larger portion of the gametospore serves as a storehouse for food, cell formation being confined to the upper end of it (Fig. 255, *B, C*).

With one exception this is the only group characterized by this peculiar germination which is more suggestive of the growth of the animal egg where a large portion of the egg is reserve food



FIG. 255. Development of the pro-embryo of pine: *A*, section of an archegonium at time of fertilization. One of the male gametes, ♂, is seen fusing with the female, ♀. The second male gamete, ♂′, the tube nucleus, and the wall cell are also shown near the neck of the archegonium. *B*, the gametospore has germinated, forming four cells, which are passing to the upper end of the sac. *C*, the cells shown in *B*, arranged at end of sac and in process of division (only two cells in this sectional view). *D*, later stage, the pro-embryo of four plates of cells. *E*, the second plate or suspensory cells, *s*, of the pro-embryo elongating, thus pushing the embryo-forming cells, *e*, into the tissues of the female gametophyte.—*A* after Ferguson; *B–E* after Coulter and Chamberlain.

for the nourishment of the embryo. The nourishment of the female gamete by the jacket cells is also suggestive of certain animals in which the egg cells are formed in a similar manner. The embryo is developed from the uppermost plate of cells which are pushed into the cells of the gametophyte by the elongation of the suspensory cells just below them (Fig. 255, *E*). Each of these four cells may form an embryo (Fig. 256, 2), but usually only one develops. The mature embryo consists of a stem bearing two or several laterally-placed cotyledons and a root with

root cap (Fig. 256, 3). Attending this formation of the embryo, pronounced changes occur in the sporangium. It steadily



Fig. 256. Development of the young sporophyte or embryo: 2, diagram of a section of a megasporangium, showing the formation of elongated suspensory cells and numerous embryos, *e; i,* integument; *sp,* sporangium nearly consumed by the growth of the gametophyte; *g,* gametophyte, the central portion of it disorganized by growth of embryos, *e.* 3*A,* section of a nearly mature embryo; *s,* suspensor; *r,* root cap; *c,* cotyledons; *st,* stem. 3*B,* external view of embryo.

increases in size and the integuments become modified into a hard coat, or the outer layer may be pulpy, as in some of the cycads. The gametophyte also increases in size, and as the em-



Fig. 257. Sectional view of pine seed: *i,* hard integument; *g,* gametophyte, often called the endosperm, which has completely consumed the sporangial tissues. The embryo consists of a root, *r,* ensheathed in a large root cap, cotyledons, *c,* and stem, *s.*

bryo matures it absorbs all the tissues of the sporangium, thus filling the space within the integument (Fig. 257). The cells of the female gametophyte are often called the "endosperm." By these growths the seed is formed and prepared for its dormant state, as in the preceding order. In some cases the integument forms a membranous outgrowth, which assists in the distribution of the seeds (Fig. 258, *C*). It is interesting to note



FIG. 258. *A*, mature strobilus of pine with open scales to permit the scattering of the seeds. *B*, scale from strobilus showing the winged seeds developed from the two sporangia. *C*, a seed with wing-like outgrowth, as it escapes from strobilus.

that the seeds are so attached to these wings that they rotate in falling to the ground, thus retarding their fall and making possible a wider flight. The stimulus resulting from the fusion of the gametes also produces extensive changes outside of the sporangia. The sporophylls or their outgrowths often become greatly enlarged, forming the hard, woody scales of the cones (Fig. 258, *B*), or they may become fleshy and fuse, forming a berry-like fruit, as in the juniper (Fig. 260, 7). When this growth has been completed, the woody sporophylls or scales of the strobilus dry out, and becoming hygroscopic, they spread apart on dry days, thus permitting of the scattering of the seeds by the winds (Fig. 258, *A*). Many of the fleshy fruits are eaten by birds and the hard nut-like seeds are distributed in this way.

The stages in the development of the seed, outlined above, are very much prolonged in many of the pines. The microspores on

reaching the megasporangia, in the spring, develop only a short tube cell during the first season and not until about the first of July of the following season are the male and female gametes mature and ready for fertilization. The seeds are matured during the two following seasons, over three years after the appearance of the strobilus. Accordingly, three stages in the development of the strobilus of megasporophylls may be seen on certain species of pine in the early summer—very small ones that received the microspores in the spring, larger ones in which fertilization and the formation of the embryo is being effected and older strobili in which the seeds are approaching maturity (Fig. 252).



Fig. 259. Renewal of growth of the embryo. At the left a seed has been cut across to show the relation of parts during the early growth. The integument has been ruptured by swelling of the seed and the protrusion of the root which is curving down into the soil. The cotyledons, *c*, remain in the seed absorbing the food from the gametophyte or endosperm, *g*. At the right a later growth with the cotyledons partially withdrawn from the seed after the absorption of its food.

When the conditions are favorable the embryo renews its growth. The root is pushed out through the micropyle and bends down into the soil, the hard integument often being cracked open by the swelling of the cells. The cotyledons remain within the seed until they have absorbed all the food from the gametophyte, when they are withdrawn and become erect (Fig. 259).

(*c*) *The Principal Genera of Pinales.*—The more important genera of Pinales may be distinguished as follows: *Pinus* or pine, leaves long and needle-like, borne in fascicles on short stems that are quite concealed by papery sheathing scales (Fig. 251, 1). *Larix* or larch, short needle-like leaves clustered in tufts on short lateral branches. This is the only northern member of the order with deciduous leaves. *Picea* or spruce, leaves angled or four-sided, radiating from all sides of stem, petioles remaining on branchlet after leaves fall, thus causing the rough appearance of the branchlets (Fig. 260, 9). *Tsuga* or hemlock, leaves flat in



FIG. 260. Common examples of the Pinales: 5, *Thuja* or arbor vitae. 6, Strobilus of *Chamaecyparis* or southern white cedar. 7, strobilus of *Juniperus* or red cedar with fleshy scales fused into a berry-like fruit. 8, branch of *Taxus* or yew. The seeds are produced singly in the axil of a leaf on short lateral branches and nearly enveloped by a thick fleshy cup that becomes bright red. 9, *Picea* or spruce.

two rows, the petiole remaining on the stem after fall of leaf; the strobilus is ovoid with rather leathery scales. The spruce and hemlock have pendent strobili. *Abies* or fir, leaves flat in two rows, the stems being smooth after leaf fall, strobili erect on the branches. *Taxodium* or bald cypress, leaves flat in two rows and deciduous, strobilus globose with thick, spirally-arranged and shield-like scales. *Thuja,* arbor vitae, or northern white cedar, leaves scale-like, closely appressed to the branches in four rows, strobili ovoid with flat tough scales (Fig. 260, 5). *Chamaecyperus* or southern white cedar, leaves resembling *Thuja,* but cones globose with shield-like opposite scales (Fig. 260, 6). *Juniperus* or juniper, pointed scale-like leaves, opposite or in whorls on stem, scales of strobili becoming fleshy and fusing to form a berry-like fruit (Fig. 260, 7).

CHAPTER X

SUBDIVISION 2. ANGIOSPERMAE. PLANTS WITH ENCLOSED SEEDS

122. Origin of the Angiospermae.—These plants, like the Gymnospermae, produce seeds, but they differ so essentially from the latter group in the character and structure of the sporophyte and gametophyte as to justify their separation into a distinct division. They are retained here as a group coördinate with the gymnosperms out of deference to common usage. It has been noticed that the seed habit arose quite independently of the Spermatophyta (see page 319) and it appears more probable that the Angiospermae have been derived from some extinct seed-producing group of plants and have developed independently of the gymnosperms. Though less ancient than the latter class, their origin is still uncertain and the meagre evidence points to their derivation from some primitive line of ferns rather than from the higher pteridophytes or gymnosperms. The variations of this modern group of plants have been many and so successful that they have crowded out many of the lower forms and become the dominant plants upon the earth. They exceed all other groups combined in variety and number of species. Approximately, 125,000 species are known that are adapted by their variations to practically all conditions that will sustain life, ranging from aquatics to xerophytes, from terrestrial to epiphytes and from photosynthetic to saprophytes and parasitic plants. The important features of the plant body have been considered in the opening chapters of the book.

123. The Sporophylls of Angiosperms.—The flower is often considered as one of the most characteristic features of the angiosperms, but this structure contains as its essential organs one or more sporophylls and the term flower could be applied quite as well to the association of these organs in the ferns and gymnosperms.

The sporophylls, like all organs of the plant, show a wide range of variation and they are generally associated with more or less modified leaf-like organs which serve to protect them. These leaf-like organs are known as the floral envelope or perianth and doubtless arose in many forms through the sterilization and modification of the sporophylls (Fig. 261*A*, 1). The microsporophylls, often called stamens, usually consist of a stalk or filament and a four-lobed spore-bearing part, the anther (Fig. 261*A*, 2). In cross-section, the anther is seen to consist of four sporangia in which the microspores originate in tetrads as in the



Fig. 261*A*. Flower and sporophylls of Angiosperms: 1, flower of *Sedum* with leaf-like perianth, *p;* microsporophylls, *s;* megasporophylls, *c.* 2, microsporophyll of the buttercup, showing four-lobed anther and filament. 3, diagram of a cross-section of an anther, showing the breaking down of the tissue about the four sporangia and the beginning of the opening of the anther. 4, one of the sporangia from a young anther, as seen in cross-section—*m,* spore mother cells. At the right a mother cell forming four microspores, the upper one being characteristic of monocotyledons and the lower of dicotyledons.

preceding groups (Fig. 261*A*, 3, 4). At maturity, the two sporangia on each side of the anther usually merge into one cavity, owing to the breaking down of the intervening tissue. The anther opens in a variety of ways, as by slits and pores, permitting the scattering of the spores (Fig. 261*A*, 3). The



FIG. 261*B*. Megasporophyll and megasporangia: 1, megasporophyll of buttercup—*s*, stigma; *st*, style; *o*, ovary. 2, longitudinal section of megasporophyll—*mg*, megasporangium; *m*, megaspore, the mother cell having formed four spores in series, but the end one only develops; *i*, integument; *mi*, micropyle.

microsporophylls are often externally quite suggestive of those noted in certain of the Pinales, but the megasporophylls are essentially different from any other group considered, in that the sporangia are inclosed in a cavity formed in the sporophyll. This peculiarity is one of the important characteristics of the angiosperms, as the name indicates. The megasporophyll, often termed the pistil or carpel, is usually a rather elongated, flask-shaped organ with a hollow swollen base and consists of a stigma, style and ovary (Fig. 261*B*, 1) within which are produced the megasporangia or ovules. Such a structure would result if the edges of the leaf-like megasporophylls of previous groups were inrolled so as to form a closed organ. The mega-

spores originate in these sporangia as already noted in the gymnosperms and usually but one megaspore is developed in each sporangium (Fig. 261*B*, 2).

124. Development of the Flower of Angiosperms.—The sporophylls are variously associated in groups that are commonly called flowers. In its simplest form, the flower may be defined as a minute branch or receptacle bearing one or more sporophylls. Such a type is illustrated in the cat-tail (Fig. 262, *B*, *C*)



Fig. 262. Forms of primitive flowers: *A*, inflorescence of *Typha* or cat-tail—*mi*, region bearing only flowers with microsporophylls; *mg*, flowers with megasporophylls; *b*, bract. *B*, flower consisting of two microsporophylls which are sessile on a short stalk that has numerous hairs. *C*, flower consisting of one megasporophyll—*s*, stigma; *o*, ovary surrounded with hairs. *D*, early appearance of the inflorescence of *Salix* or willow. *E*, inflorescence bearing only megasporophylls. *F*, flower of a single megasporophyll with forked stigma—*b*, bract; *n*, nectar gland. *G*, inflorescence bearing only microsporophylls. *H*, flower of two microsporophylls.

where the flower consists of one or a few sporophylls associated with hairs, and also in the willow where the sporophylls are developed in the axil of a minute bract (Fig. 262, *F*, *H*). These

simple types of flowers are often developed in large numbers upon an elongated stem (Fig. 262, *A, D–G*) and are rather suggestive of a strobilus although the individual flower is more comparable with a strobilus as will be seen especially in those types where the sporophylls are numerous and arranged spirally upon the receptacle of the flower (Fig. 264). A group or cluster of flowers is called an inflorescence in contradistinction to the solitary flower developed at the end of a branch or stem. It is noticeable in these primitive angiosperms that the micro- and mega-sporophylls are usually borne in separate flowers or inflorescences; compare Pinales, which are developed on separate plants as in the willow or on different parts of the same plant as in the cat-tail. This arrangement is probably associated with the fact that the advantages of crossing or the transfer of the microspores of one flower to the megasporophylls of another is effected by the wind. In higher types, which include the great majority of angiosperms, the micro- and mega-sporophylls are developed in the same flower which is therefore said to be perfect since it contains both of the organs essential for seed production (Fig. 261*A*, 1). A type like the willow is termed imperfect because the flower lacks one kind of sporophyll. Crossing is effected in the perfect type of flower by the earlier ripening of the micro- or mega-sporophylls and often also, by the arrangement of the organs of the flower which is of such a nature that the microspores cannot readily reach the mega-sporophylls of the same flower. Insects are usually the agents for the transport of the microspores in such cases. Flowers in which the microspores mature and are shed from the anthers before the stigma of the megasporophylls are ready to receive them, are called protandrous, meaning that the microspores which develop the male gametes, are the first to mature. If the mega-sporophylls become receptive before the anthers open, the flower is said to be protogynous, meaning that the pistil matures first. It must also be borne in mind, although the devices for effecting a crossing are almost universal among the various groups of angiosperms, that there are equally elaborate provisions for the transference of the microspores of perfect flowers to the stigmas

of their own flower. This is called autogamy and would appear to be a provision for setting seed in case crossing fails.

Following the development of the type of flower noted in the cat-tail and willow, there appeared as the next advance in the flower, minute outgrowths about the sporophylls, known as the perianth. In its simplest form this consists of a few scales as in the sweet flag, oak, etc. (Fig. 263, *B*), but in higher forms,

FIG. 263. Development of the perianth: *A*, inflorescence of *Quercus* or oak—*mi*, inflorescence with flowers bearing only microsporophylls; *mg*, inflorescence with flowers bearing megasporophylls. *B*, flower of oak enlarged, consisting of several microsporophylls and a perianth of minute scale-like organs. *C*, flower of *Erythronium* or fawn lily. The perianth of six conspicuous leaf-like organs. *D*, flower of *Melandryum* or day pink —*ca*, calyx of green sepals; *c*, corolla of five delicate petals.

the perianth appears as the conspicuous leafy portion of the flower as in the lily (Fig. 263, *C*). Finally flowers appear in which the leaves of the perianth become differentiated into an outer calyx composed of several green sepals and a corolla of larger, more delicate and often brightly-colored leaves, called the petals (Fig. 263, *D*). We have now reached a point where the flower

is said to be complete, consisting of all the organs that are normally associated in the flower. The receptacle or region upon which the sporophylls and perianth are arranged, is often somewhat elongated in the simpler types of flowers, thus giving room for the separate attachment of the organs. In such cases the various parts are spirally arranged and very numerous (Fig. 264). This spiral type of flower is characteristic of a large number of the lower orders of angiosperms and, as stated above, is suggestive of a strobilus.

The shortening of the receptacle is one of the most important variations that appear in the evolution of the flower. This is brought about by the checking of the apical growth of the receptacle and is often associated with a more or less extended growth of its basal region. These changes affected the flower



Fig. 264.

Fig. 265.

Fig. 264. Flower of strawberry with elongated receptacle bearing numerous spirally arranged sporophylls: *A*, open flower. *B*, flower in section, showing arrangement of parts upon receptacle, *r*, which forms a shallow cup at base bearing the perianth and microsporophylls. *C*, the fruit or enlarged receptacle bearing the minute spirally-arranged megasporophylls.

Fig. 265. Flowers with shortened receptacles: *A*, flower of *Pyrola* with calyx and corolla arranged in whorls or cycles. *B*, section of flower, showing all the organs in cycles. *C*, flower of geranium. *D*, flower with corolla removed to show the coherence of the five megasporophylls that results from the shortening of the receptacle.

in a most profound way. A crowding resulted, organs were reduced in number and their spiral arrangement upon the receptacle became so flattened that the various sets of organs appear to arise in whorls or cycles (Fig. 265, *A, B*). These cyclic flowers are characteristic of all the higher orders of angiosperms. The crowding of the organs on the receptacle also caused them early in their development to come in contact with one another laterally and to cohere, forming a united structure. This tendency is especially noticeable in the megasporophylls,



FIG. 266. Forms of adhesion that result from shortening of receptacle: *A*, flower of rose. *B*, section of flower, showing the lower portion of receptacle forming a cup about the megasporophylls, *mg*, and bearing the other organs of the flower. *C*, inflorescence of comfrey, *Symphytum*. *D*, flower enlarged in section to show adhesion of microsporophylls, *mi*, to the tubular corolla.

which very frequently cohere into a compound megasporophyll (Fig. 265, *C, D*), and in the same way the sepals and petals become united, forming a more or less tubular calyx and corolla (Fig. 266, *C*). The crowding also led to the adhesion of the organs of adjacent sets or whorls. This is often seen in the case of the petals and microsporophylls, the latter organs appearing to arise from the corolla owing to the union of the lower portion of the filament with the base of the petals (Fig. 266, *D*).

More frequently the receptacle plays an important rôle in all cases of adhesion. As stated above, its apical portion ceases to elongate at an early period while the basal part continues active, forming a cup about the ovaries. Sometimes the sepals, petals, and even the microsporophylls are so associated with this basal growth of the receptacle that they appear to arise *en masse* about the ovaries. This growth may be so slight that a careful examination of a section of a flower is required to detect it or a conspicuous cup-like structure may be formed, as in the rose (Fig. 266, *A, B*). This type of flower is termed perigynous, meaning that the receptacle and other organs form a more or less conspicuous cup about the ovaries. In simpler types, as the spiral flowers, it can be seen that the megasporophylls arise at the top of the receptacle and that each of the other organs arise at a point just below the next inner one. Such flowers are called for this reason hypogynous, meaning below the ovaries. Very frequently the basal growth of the receptacle also results in an adhesion to the ovaries which becomes partially or completely overgrown, so that the other organs of the flower arise from the sides or from the top of the ovaries; the flowers being partially or completely epigynous, meaning that the organs of the flower are developed upon the ovary (Fig. 267, *B, D*). Often in completely epigynous flowers the receptacle elongates very slightly and the basal portion grows up around the apex forming a cup-like cavity which is roofed over by the megasporophylls. The sporangia or ovules usually arise from the walls of the cavity thus formed and not from the walls of the megasporophylls at the top of the cavity (Fig. 267, *D*). The calyx, corolla and microsporophylls arise from the top of the structure thus formed and these organs may be developed separately or there may be varying degrees of cohesion and adhesion.

Another feature to be noted in connection with the evolution of the flower is its symmetry. In the lower types, the organs of a set are alike and radially arranged about the center of the flower (Fig. 267, *A*). This is the regular or actinomorphic type of flower, meaning radially symmetrical. In many of the orders of angiosperms one or more members of a set are different from

the others, thus destroying the radial symmetry (Fig. 268). This is the irregular or zygomorphic type of flower, meaning yolk-form. Such flowers can be cut into two similar halves in but one plane.

It must not be understood that evolution of the flower has progressed steadily through the various changes outlined above



Fig. 267.

Fig. 268.

Fig. 267. Adhesion due to basal growth of receptacle: *A,* flower of saxifrage. *B,* section of flower, showing megasporophyll partially inclosed by receptacle. *C,* inflorescence of red currant. *D,* section of flower, showing the receptacle forming a sporangial cavity that is covered at the top by the megasporophylls. The other organs arise from the top of this structure, the microsporophylls adhering to the corolla.

Fig. 268. Irregular or zygomorphic flower of honeysuckle: *ca,* calyx; *c,* corolla of five unequal cohering petals; *s,* stigma.

and that consequently in the following lessons we can begin with the most primitive type and proceed by regular steps to the highest forms. The various orders of angiosperms have doubtless been derived from several distinct stocks and they have not only varied in different degrees but especially will it be noted that some orders have a tendency to emphasize certain forms of these variations, while in other alliances, the variations will

proceed along quite different lines. These various modifications of the simple type of flower have been retained very largely because they are mutually beneficial to the plant and to insect life. In the lower forms, the microspores are scattered by the wind as in the gymnosperms. Such flowers are called anemophilous, meaning wind loving, and are small, very simple in structure and produced in large numbers to ensure the formation of seed. In higher types of flowers, odor and nectar glands appear and bright colors which serve to attract insects. Such flowers are termed entomophilous, meaning insect loving. In this way the spores are carried from one flower to the other by the insect visitors with greater certainty than in the case of the anemophilous plants. The variation in the form of the flower and insect went on hand in hand and so it came about that types of flowers finally appeared that were adapted to special kinds of insects, sometimes to a single species, all other forms being unable to enter the flower owing to its peculiar shape. This was of the greatest benefit to both plant and insect because the insect will confine its visits to such flowers owing to the food which it alone can secure and the plant is equally fortunate, since the microspores will only be carried to flowers of the same kind. So we will expect to find in the following lessons that the highest types of flowers are adapted to special kinds of insects and it would be a natural inference that such plants will be very abundant provided of course that they are also adapted to the present conditions upon earth.

125. The Gametophyte of the Angiospermae.—The angiosperms have developed along quite distinct lines, but they show such a remarkable uniformity in the development and character of the gametophyte generation that this feature of their life history may be considered at this point as it applies to all forms. The megasporangium, also called the ovule, originates in the cavity of the ovary at various points known as the placenta (Fig. 269, *A*). The structure of the sporangium and the formation of the megaspore is very similar to that of the gymnosperms. More often two integuments are formed and the sporangium or the stalk which supports it becomes curved so that it very fre-

quently is turned completely over (Fig. 269, *B*). The germination of the megaspore results in a female gametophyte which is very much more reduced than in the case of the gymnosperms. As the megaspore enlarges, disorganizing the cells of the sporan-



FIG. 269. Development of the megasporangium and megaspore: *A*, sectional view of the pepper-grass, *Lepidium*. This flower has two cohering sporophylls, only a portion of the right-hand one being shown. *s*, stigma with protruding cells to receive the microspores; *mg*, megasporangium containing a single spore mother cell, *mc*. Two integuments, *i*, are growing up about the sporangium. *B*, later stage of development, the megasporangium, *mg*, becoming inverted and completely covered by the integument. The mother cell of *A* has found four daughter cells in series and not in tetrads as in the Pteridophyta. The innermost cell of the series, *ms*, only matures as a megaspore; *m*, micropyle; *f*, stalk or funiculus of sporangium.

gium, also called the nucellus, its nucleus divides and the daughter nuclei move to each end of the spore (Fig. 270, *A*). Each of these nuclei divides twice, forming four nuclei at either end of the spore (Fig. 270, *B*, *C*). A nucleus from each of these

groups, called the polar nuclei, now approach each other and fuse, forming a single large nucleus that is usually called the endosperm nucleus. This rudimentary growth represents the female gametophyte. The three outer or micropylar cells are not pro-



FIG. 270. Germination of the megaspore: *A*, first division of the megaspore. *B*, second division of the nuclei. *C*, final division of the nuclei.

vided with walls and consist of a rather lower-lying gamete and two nourishing cells, the synergids or helpers (Fig. 271). The inner group or antipodal cells usually have walls and they are either soon disorganized and absorbed by the enlarging gametophyte or they may remain as permanent features of the gametophyte for a long time and even increase greatly in number, serving to nourish the gametophyte by absorbing food from the sporangium. The endosperm nucleus plays a very important rôle in the development of the sporophyte, for as soon as fertilization has been effected it forms by repeated division a mass of cells that completely fill the entire space within the enlarging spore, thus providing food for the nourishment of the sporophyte. It would appear impossible to compare the various cells of this peculiar gametophyte with the tissues of the female gametophyte of the gymnosperms or ferns. It has been suggested that the female gamete and the synergids are the remains of three archegonia,

only one of which is usually capable of being fertilized, and that the antipodal cells are a remnant of the numerous vegetative cells of the gametophyte. The peculiar formation of the endosperm nucleus through the fusion of the two polar cells is looked upon

FIG. 271. Section of a megasporangium of lily, showing the mature female gametophyte: ♀, female gamete, below which are two synergids; *p*, the two polar nuclei uniting to form the endosperm nucleus; *a*, antipodal cells; *mi*, micropyle; *i*, integuments; *f*, funiculus in which a vascular bundle, *v*, has been formed to transport foods to the sporangium.

as a nourishing device to give the endosperm nucleus the power to grow and form a tissue that supplements the antipodal cells and so takes the place of the nourishing gametophyte of the gymnosperms. According to this view, the completion of the gametophyte in the angiosperms is delayed until the fertilization is effected. However, the origin and behavior of the various

parts of the female gametophyte of the angiosperms is so different from any of the preceding cases as not to warrant a comparison in the present state of our knowledge.

The male gametophyte presents several features suggestive of the gymnosperms. The germination of the microspores usually begins within the sporophylls, and by the time that they are shed and carried to the stigma, their nuclei have already divided once and each spore consists of a large tube cell and an antheridial cell (Fig. 272, *B*). There is very rarely a trace of the vegetative



FIG. 272. Germination of the microspore: *A*, mature microspore of lily. *B*, first stage of germination—*t*, tube cell; *a*, antheridial cell. *C*, final division, in this case effected while in the microsporangium—*t*, tube cell; *a*, antheridial cell forming directly to male gametes. *D*, diagram showing the formation of the tube which grows down the style and finally reaches the female gametophyte. The two male gametes, *g*, are shown passing down the tube; *t*, tube nucleus.

cells of the gametophyte, as in the cycads and pines, since the necessity for these cells no longer exists. The stigma is generally provided with minute outgrowths or papillae derived from the epidermal cells which serve to hold the microspores (Fig. 269, *s*). These cells of the stigma usually secrete a sugary solution which nourishes the microspores and causes a continuation of their germination. It is noteworthy that these microspores may be made to germinate by placing them in sugar solutions, but the approximate strength of the solution on the stigma must be determined

in order to prepare a solution suitable for their development. The tube cell, stimulated by the secretion of the stigma, ruptures the outer wall of the spore, which is provided with one or more thin places to favor this growth, and protrudes as a delicate tube. This tube, owing to the fact that it is repelled by the oxygen of the air, grows down into the tissues of the style which are really a continuation of the stigma. These tissues of the style are usually looser and provided with abundant foods which are deposited in the cells just ahead of the elongating tube to nourish and direct it in its growth. In this way the tube is directed down the style to the cavity of the ovary where, owing to the attractive influence of the organic substances in the sporangium, possibly in the synergids, it usually turns out into the cavity of the ovary, enters the micropyle and works its way through the sporangium, and, unlike the Pinales, enters the female gametophyte alongside of one of the synergids (Fig. 273). The antheridial cell usually



FIG. 273. Section of the micropylar end of the megasporangium, showing the process of fertilization. The tube, *t*, has passed through the micropyle, entered the female gametophyte and ruptured, discharging the male gametes. One, ♂, is shown fusing with the female gamete, ♀, and the other one, ♂', is uniting with the two polar nuclei, thus making a triple fusion in the formation of the endosperm nucleus; *s*, one of the synergids; *i*, integuments.

divides, forming directly two motionless male gametes during the elongation of the tube, and in other cases the gametes are already formed when the microspores are discharged from their sporangia. In none of these cases is there any indication in the division of the antheridial cell of the formation of a wall cell as in the gymnosperms, so that the male gametophyte of the Spermatophyta presents a very regular series of reductions from the cycads to the pines and thence to the angiosperms, where it consists of a tubular growth containing three naked cells (Fig. 272, *D*). This development of the male gametophyte requires from a day to several months and is quite independent, apparently, of the distance that it has to traverse in reaching the female gametophyte. The male gametes are often somewhat elongated and even spirally coiled and carried to the end of the tube, as in the Pinales.

126. Fertilization.—The end of the tube finally ruptures, owing to the tension of the fluids that gradually accumulate in it, and the male gametes are forcibly expelled into the sac-like cavity of the female gametophyte (Fig. 273). One of the male gametes passes over to and fuses with the female gamete, thus forming the gametospore. In a constantly increasing series of cases the other male gamete has been observed to fuse with the endosperm nucleus, thus making a triple fusion of cells in the formation of this nucleus. While this process is probably in the nature of a reinforcement, enabling the endosperm nucleus to perform its work, it is noteworthy that the qualities of the male parent are transmitted to the endosperm cell just as though this fusion were a sexual process.

127. The Germination of the Gametospore.—After fertilization the endosperm nucleus divides repeatedly, and usually the resulting nuclei become arranged about the walls of the sac-like gametophyte which may now be called the embryo sac (Fig. 274). Later the endosperm cells develop walls and by further division completely fill the embryo sac with cells. This mass of cells is called the endosperm and the method of its development is exactly similar to that of the female gametophyte of the gymnosperm, and it serves the same purpose, namely, to nourish the young sporophyte or embryo. Note that it has had an entirely

different origin from the gametophyte of the gymnosperms, which is unfortunately also often called the endosperm.

The germination of the gametospore and the formation of the embryo vary so greatly that only a very general statement can be made. Following fertilization, the gametospore becomes sur-



FIG. 274. Sectional view of megasporophylls of *Lepidium* shortly after fertilization (see Fig. 269): *en,* early development of endosperm cells about wall of embryo sac; *p,* young sporophyte developing from gametospore; *mi,* micropyle; *s,* stigma; *f,* funiculus.

rounded by a cell wall and attached to the wall of the embryo sac (Fig. 275, *A*). The first division of the gametospore is transverse, resulting in the formation of an outer cell, *i. e.,* facing the micropyle, and an inner daughter cell. The outer cell divides several times, usually in the same plane, forming a row of cells (known as the suspensor) which push the inner daughter cell (called the embryo cell) into the cavity of the embryo sac. The suspensor and embryo cell constitute the pro-embryo (Fig. 275, *B*). The synergids, which are partly consumed during the entrance of the tube cell and the process of fertilization, usually become entirely disorganized and absorbed during these early stages of germination. The embryo is formed by the repeated

divisions of the embryo cell of the pro-embryo, assisted to a varying extent by one or more of the adjoining cells of the suspensor (Fig. 275, *C–F*). The remaining cells of the suspensor



FIG. 275. Stages in the germination of the gametospore of *Lepidium*, sectional view: *A*, micropylar end of embryo sac, showing the enlarging gametospore provided with cell wall and attached to wall of sac. *B*, later growth—*s*, suspensor; *e*, embryo cell; *en*, endosperm cells. *C*, pro-embryo after first division of embryo cells. *D*, further divisions of embryo cell, showing formation of an epidermis and a central stem region. *E*, later growth, two growing regions, the cotyledons, appearing on the sides of the stem. *F*, micropylar end of the embryo sac in which the embryo cell has formed a small plant or embryo, consisting of two cotyledons, *c*; stem, *st*, which terminates in the root, *r*; the endosperm cells, *en*, are being absorbed by the enlarging embryo; *s*, suspensor.

are ultimately disorganized or they may increase greatly in size and themselves become the principal means of absorbing food and transferring it to the embryo. The structure of the mature embryo varies greatly. In some genera, as in the orchids, Indian pipe, etc., it remains rudimentary, consisting of only a few cells.

Among the monocotyledons, the embryo cell frequently produces the single cotyledon, while the next underlying cell of the suspensor forms the root and the laterally-placed growing point of the stem (Fig. 279). The pro-embryo of dicotyledons is frequently a filament of cells of varying length and the embryo cell, by a regular service of divisions, gives rise to the stem, two laterally-placed cotyledons and all of the root, save the tip, which is formed from the cells adjoining the embryo cell (Fig. 276, *A*).



FIG. 276. Seed structure: *A*, section of a nearly mature seed of *Lepidium*. The stem of the embryo is differentiated below into a hypocotyl, *hy*, and above into an epicotyl, *pl*, commonly known in the seed as the plumule. *r*, root with root cap; *c*, the two cotyledons, which are bent over, lying one upon the other; *v*, vascular bundles extending through the stem into the cotyledons, where they form a network of veins; *en*, remains of endosperm. *B*, section of seed of water lily (after Conard)—*e*, embryo, surrounded by a layer of endosperm cells; *mg*, cells of the megasporangium; *i*, integument.

128. The Fruit and Seed.—Various changes occur in the sporangium during the growth of the embryo. More frequently, perhaps, the embryo sac enlarges, absorbing all the cells within the integument, and it becomes filled with endosperm cells. In a case like this, the embryo either remains small and embedded

in the endosperm (Fig. 279), or the embryo may entirely consume the endosperm, the food in this case being stored in the cotyledons (Fig. 276, *A*). Less commonly, the embryo sac absorbs only a portion of the sporangial tissue (often called the perisperm) and consequently the embryo is associated with a varying amount of endosperm, which in turn is surrounded with sporangial cells or perisperm, as in the water lily (Fig. 276, *B*). The integument usually undergoes pronounced changes during this growth, becoming hard and tough, to protect the parts within and often developing appendages of various kinds, as hairs and wings, to promote the distribution of the seed (see gymnosperms, page 341).

The stimulus of fertilization extends beyond the changes wrought in the sporangium. This is particularly noticeable in the megasporophyll and often in adjacent parts which keep pace with the growth of the sporangium and often undergo remarkable transformations. The result of this total growth is called the fruit, while the term seed is restricted to the modified sporangium with its integument and embryo. The megasporophyll may form a firm coat that is closely attached to the seed, as in the corn and other grasses. Such a fruit is called a grain. Again the tough walls of the megasporophyll are free from the seed, as in the buttercup, forming a fruit known as the akene. The sporophylls may become papery or hard and split open to scatter the seed. Where a single megasporophyll behaves in this way and opens by two valves, the fruit is called a pod or legume, example the bean, or if by one valve the fruit is termed a follicle, example the peony. Frequently wing-like processes develop from the megasporophyll, as in the maple and ailanthus, which are of service in distribution and in other ways, as in the manufacture of foods, these organs often being green during the development of the embryo. Such fruits are known as key fruits or samaras. In other cases the sporophyll becomes fleshy, forming a berry as in the currant or the inner layer forms a pit or stone, while the outer layer forms the pulp and skin, as in the cherry and peach, fruits known as drupes. In many fruits the receptacle becomes fleshy and forms the major portion of the fruit, as in the fig,

apple, strawberry, etc., and in the pineapple, the axis of the inflorescence, as well as parts of the flower, become fleshy. These various devices serve primarily to preserve the life of the embryo and some of them also assist in distributing the seed.

When conditions are favorable for growth, which may not be till several years after the seeds have been scattered, the embryo renews its growth. The region below the cotyledons, the hypocotyl, or base of the cotyledon itself, elongates and pushes out the root which grows down into the soil, and by a later growth the stem tip, with or without the cotyledons which may remain in the seed, elongates and develops the characteristic stem, branches, leaves and sporophylls of the parent plant. The structure and variations of the mature sporophyte have been presented in the first part of the work, pages 7–122, and will be further considered in the following studies. The terms sporangium, microsporophyll and megasporophyll have been retained up to this point in the work in order to keep clearly before you the progress and relationship of the variations that have attended the evolution of plant life. Now that we have arrived at the highest and largest group of plants, where the sporophylls have become highly modified and curiously associated in clusters called flowers, more familiar names will be used in discussing them, *i. e.,* ovule for the megasporangium and its integument, stamen for microsporophyll and pistil for megasporophyll. The terms "carpel" and "pistil" are synonymous when these organs are distinct—*i. e.,* free from one another; but when they cohere to form a compound pistil the term "carpel" is employed to indicate the number of pistils that are united, as in the willow the pistil is composed of two carpels, in the lily of three carpels, etc.

While the sexual generation, mode of reproduction, and the character of the tissues are very similar in all angiosperms, two very distinct lines of variation or classes are to be noted: A, the Monocotyledones, represented by the grasses, lilies and orchids; B, the Dicotyledones, represented by our common trees, roses, mints and daisies.

Class A. Monocotyledones

129. General Characteristics.—This group of angiosperms contains about 25,000 species that constitute a very natural alliance, owing to the uniformity and simplicity of their structures. These features may be due to the uniform conditions under which these plants live. The majority of Monocotyledones are moisture-loving plants and are therefore exposed to very constant conditions, which would naturally result in less stimulation and consequent variation than in the case of plants exposed to the varying conditions of drier soils. Some, to be sure, have become adapted to dry and even arid regions, owing to peculiar modifications of their leaves or stems, as in the grasses, certain bulbous plants, etc.

The leaves are smooth and simple, lance-shaped or linear in outline, sessile and often attached to the stem by sheathing bases. The veins do not end in free branches on the margins of the leaves, and as a result of this closed venation the leaves are usually entire and destitute of teeth or lobed margins. In many instances the principal veins are quite parallel, but whatever the arrangement, the prominent veins are connected by very minute veinlets that form an inconspicuous network or reticulation throughout all parts of the leaf (Fig. 277). The stems are composed largely of parenchyma, through which are scattered numerous veins or vascular bundles derived entirely from the leaves (Fig. 278). These bundles do not develop a cambium (Fig. 60) and consequently the stem does not increase materially in diameter, and usually it is columnar in appearance. In many instances the stems are reduced in size and subterranean, the bulb and rhizome being common forms of stems which send up annually short-lived aërial branches. As a rule the stems do not branch, owing to the failure of the buds in the axils of the leaves to develop.

The flowers are also of rather simple and uniform structure, in the simplest cases being imperfect and consisting of one or more naked stamens or of one or several pistils. In the higher types spiral flowers with perianth and both kinds of sporophylls appear, and these give place to forms in which the organs are

arranged in successive whorls of three parts each, the latter type of flower being especially common. Perigynous, epigynous and irregular flowers are of less common occurrence. The most conspicuous feature of the monocotyledons is the embryo, which has a single terminal cotyledon and a laterally developed stem tip (Fig. 279). Generally the root and stem tip are set free from

Fig. 277.



Fig. 278.

Fig. 277. Leaf of Solomon's seal with closed venation, entire margins, and sessile upon the stem, *i. e.*, without petiole.

Fig. 278. Cross-section of stem of rush: *v*, vascular bundles; *st*, stereome.

the seed by the elongation of the basal portion of the cotyledon, while the cotyledon remains in contact with the endosperm as a food-absorbing organ, being often highly modified to perform this work. An examination of a few of the more important orders will give a better idea of the characteristics and variations of the monocotyledons.

130. Pandanales, the Cat-tail Order.—The cat-tail, *Typha*, may be taken as an illustration of this order (Fig. 280). These plants live in wet and marshy places and present several variations that adapt them to such conditions The main stem is a rhizome that branches through the mud and sends up each spring leafy stems. The rhizome grows on from year to year, and

owing to the decay of the older portions, the branches become independent plants. In this way a single plant may spread and thickly populate an extensive marsh. The aërial stems live but for a year and are rather weak, but owing to the sheathing bases of the leaves they become sufficiently rigid to support a heavy



FIG. 279. Sectional view of seed of *Veltheimia*. The embryo consisting of a large cotyledon, *c*, and laterally placed stem, *s*, below which is the root. *e*, endosperm; *mi*, micropyle; *f*, funiculus.

foliage and withstand the winds. Notice also the extreme lightness of these organs. This is due to the large air spaces which also permit a ready interchange of gases from the leaves to all parts of the plant, even throughout the submerged rhizomes. You would naturally expect to find this structure in all aquatics, since the roots at least are submerged and all living cells require an interchange of gases (see page 46). The leaves are long and narrow and point nearly straight up in the air. This prevents shading and permits the association of the plants in dense colonies. The leaves are covered with a waxy coating or bloom to prevent the adhesion of water and the plugging of the stomata. This device is often to be seen on the leaves of plants that are subject to heavy dews or rains. Moisture is frequently not evaporated from the leaves until near midday, and if the stomata

become filled with water, there can be no interchange of gases for photosynthesis during this time.

The flowers are of a very primitive type, consisting of naked sporophylls arranged in a compact inflorescence that assumes a spike-like structure at the tip of the stem. The sporophylls are protected until mature by modified sheathing leaves (Fig. 280, *A*). Each flower in the upper portion of the spike contains only two or three stamens supported upon a short stalk, which is associated with hair-like outgrowths (Fig. 280, *B*), while the lower



Fig. 280.



Fig. 281.

Fig. 280. Inflorescence and fruit of *Typha: A*, inflorescence—*b*, protecting bracts curving away from the sporophylls; *mi*, region of staminate flowers; *mg*, region of pistillate flowers. *B*, staminate flower consisting of two stamens sessile upon a short stem. *C*, pistillate flower of one carpel—*s*, stigma; *o*, ovary containing a single ovule. *D*, appearance of *A* in the fall—*mi*, region occupied by staminate flowers; *mg*, the pistils have increased greatly in size during the ripening of the seed.

Fig. 281. The mature fruit of *Typha: s*, remains of the stigma; *o*, ovary; *p*, elongated pedicel bearing numerous hairs. Compare Fig. 280, *C*.

flowers of the spike bear a single pistil each (Fig. 280, *C*), which consists of a large flat stigma, style and ovary containing a single ovule and supported upon a hairy stalk or pedicel. The microspores are carried to the stigmas of the pistils by the wind.

Such anemophilous flowers are characterized by several features well illustrated in the cat-tail. The microspores must be produced in large numbers since the chance of one reaching the stigma of another plant rapidly decreases as the distance traversed by the spores increases. This probably accounts for the growth of many anemophilous plants in dense colonies since the close proximity of the plants greatly increases the chance of the microspores reaching the stigmas. The Pinales, grasses, willows, oaks, etc., are other illustrations of a very large series of plants, some 10,000 in number, that have a similar habit. Anemophilous flowers are inconspicuous and simple in structure. You will notice that showy perianths, nectar and perfume glands are only developed in such flowers as utilize insects for the distribution of the microspores. The stigmas of anemophilous flowers are usually large and hairy or brush-like and conspicuously exposed so as to increase the chances of catching the microspores, thus reducing the dangers of this rather risky method of crossing. You will also observe that anemophilous flowers are usually characterized by having imperfect flowers, the stamens and pistils being developed on different parts of the plant, or on different plants. By this arrangement, the advantages of crossing are secured and it also happens that the microspores are either scattered before the stigmas of neighboring flowers are mature, or, more frequently after they have withered and are therefore no longer capable of catching and nourishing the spores. In this way it comes about that the microspores, even when close to the pistillate flowers as in *Typha,* are often only of service when carried to some earlier flowering plant whose stigmas are mature. In the cat-tail, the microspores are shed a day or so before the stigmas on the same spike are mature and so there must result the benefit that comes from crossing two more or less widely separated plants (page 128). The stamens soon perish after the discharge of their spores, but the pistils increase

greatly in size during the ripening of the seed and form conspicuous brown cylinders (Fig. 280, *D*). As winter approaches, the pedicel below the ovary and the hairs attached to it become greatly elongated and the sporophyll is thus transformed into a very stable parachute that is easily detached from the spike and capable of floating the seed in even the lightest winds (Fig. 281). If this fruit chances to fall in a marsh, the torpedo-like seed after a time falls from the ruptured ovary and sinks in the water. When the growth of the seed is renewed, the base of the cotyledon elongates, pushes off the lid at the end of the seed and curves down so as to bring the root of the embryo in contact with the mud. Hairs now develop from the lower part of the cotyledon, anchoring the young plant to the ground while the tip of the cotyledon remains in the seed until the food is absorbed, when it is withdrawn. In the meantime, roots have developed and penetrated the soil and the stem tip begins to elongate, lifting up the cotyledon and making possible the formation of other leaves. The curious screw-pines, *Pandanus,* a large group from the oriental tropics and frequently seen in conservatories, belong to this order, also the bur reeds (*Sparganium*), common about marshes and water ways, which are the highest members of the order.

131. **Graminales, the Grass and Sedge Order.**—This is one of the largest groups of the Spermatophyta, some 7,000 species being known, and in number of individuals it exceeds all others (Fig. 282). These plants are almost universally distributed over the earth and have become adapted to almost every condition of climate and soil. The ability of these plants, which are of aquatic origin, to establish themselves upon the drier and more diversified land surface doubtless promoted variation and accounts in part for the enormous display of forms. The Graminales and the palms (Principales) which belong to an order related to *Typha* are about the only groups of monocotyledons that are associated in sufficient numbers to form striking features of the vegetation of a country. This is particularly true of the Graminales in all lands where they constitute the principal vegetation of hills, plains, meadows and marshes. These plants are

also of the greatest economic importance, furnishing a variety of foods, as wheat, rye, barley, oats, rice, corn, hay, etc. The bamboo also belongs to the order. The stem in the majority of forms is a rhizome (as in *Typha*), which branches extensively through the soil and sends up numerous aërial branches (Fig. 282,



FIG. 282. Habit of growth of one of the grasses: *A*, aërial stem terminating in a branched inflorescence. *B*, underground stem or rhizome sending up a new shoot from one of the nodes. *C*, section through the node of a stem that has been placed horizontal, showing the sheathing leaf base, *l*, and the beginning of the upward curvature of the stem.

B). These interwoven rhizomes, with their numerous roots, form the firm swards of meadows and prairies. The aërial stems are models of mechanical construction. The ability of the long hollow stems of the grasses to support the heavy head of grain must appeal to every one. An examination of Fig. 282 will show you how the stem is reinforced by the long sheathing base of the leaves while tough, elastic strands of stereome fibers are so distributed as to give the best mechanical support. An un-

usual localization of the growing regions also characterizes the grasses which enables them to lift up again their stems if they by any means become prostrate. In case the stem becomes prostrate, the stimulus of gravity causes a renewal of growth in the cells at the base of the node on the side of the stem next to the ground, thus causing the stem to curve up (Fig. 282, *C*). You can easily demonstrate this peculiar localization of growth by cutting off a growing stem of grass and placing it horizontally with one end embedded in moist sand and noting the curvature of the node after a day. The leaves show wide variation in structure and are doubtless one of the factors that have enabled these plants to live under a variety of conditions. For example, where the plants are exposed to drought or drying winds the epidermal cells are very much thickened, stomata often sunken in narrow furrows and practically all species have the power of rolling up the leaves in dry seasons and thus reducing the leaf surface and lessening transpiration.

The flowers are exceptionally alike in structure and show about the same state of floral development. While resembling those of the *Typha* in some particulars, they present several features that indicate a decided advance over the previous group. Fig. 283 illustrates the common types of inflorescence found among the grasses. The flowers are arranged on elongated branches, but instead of a large bract ensheathing the inflorescence (see *Typha*), each flower is inclosed by one or more small bracts so arranged as to form a spike-like structure of overlapping bracts (Fig. 283, 2). The two kinds of sporophylls may be arranged in separate spikes or on different parts of the same spike, or, as in many grasses, the stamens and pistils are developed in the same flower. One or more of the lower bracts of a spike are usually without flowers and above them are one to several flowers enclosed by secondary bracts (Fig. 283, 4). Removing the secondary bracts, the flower proper is seen (Fig. 284, 3*A*, 3*B*). This consists in many of the grasses of three stamens and one pistil and usually two small scale-like organs, the lodicules, which assist by their expansion in forcing open the protecting bracts at the time of flowering. The stamen con-

sists of a long filament attached to the middle of the anther so that the latter organ can swing at the end of the filament, an arrangement known as the versatile anther. The stigma is often brightly colored and of a delicate feathery character, indi-

FIG. 283. Inflorescence of a grass: 1, tip of stem with flowers arranged in spike-like inflorescences. 2, a single spike enlarged at time of flowering. 4, diagram showing structure of spike. At base two sterile bracts, above three flowers, each enclosed by an outer firm bract and an inner more delicate bract. *l*, lodicules. At apex of spike a sterile flower. 5, another species of grass, showing the scattering of the microspores.

cating that the plants are anemophilous. The ovary contains a single ovule. The manner of flowering of the numerous genera of this order varies. In many cases the stigmas are first extruded from the bracts and can therefore only be crossed with the microspores from some earlier flowering plant. In other instances, stigmas and anthers are extended together and in some genera no injury results from the transfer of the microspores to the stigmas of the same flower. It is worth any one's

time to note the time of flowering, the region on the spike where it begins and the manner of opening of the bracts and extension of anther and stigma. Many open between 6 and 7 A. M. on pleasant days. Owing to the swelling of the lodicules, or, of



Fig. 284.

Fig. 285.

Fig. 284. Flower and fruit of grass: *3A*, a single flower with the two enveloping bracts opened, exposing the stamens and pistil with feathery stigmas. *3B*, flower with outer firm bract removed—*l*, lodicules; *st*, stigma. *6*, mature fruit or grain—*e*, region of embryo. *7*, section through base of grain, showing the root, stem, leaves, and scutellum, *sc*, or absorbing organ of the embryo; *en*, endosperm. *7A*, diagram of a few of the outer cells of the scutellum, *sc*.

Fig. 285. Inflorescence of one of the sedges, *Carex: p*, spike of pistillate flowers, each pistil is surrounded by a papery sac, through which the style and stigma protrude; *s*, spike bearing staminate flowers. Note the triangular stem, a characteristic of this large genus of over 1,000 species.

the base of the bracts, these latter organs are forced apart, permitting the extension of the stigma and anthers. The filament quickly elongates and soon curves so as to allow the anther to

swing back and forth in the wind. The two lobes of the anther now curve apart and open at their ends by a narrow slit, forming a cup in which the microspores collect (Fig. 283, 5). As the rocking of the anther in the wind sifts out the spores, more rattle down into the cup and so a very gradual scattering of the spores is effected. This entire process may be effected in a few minutes, and in the case of the wheat each flower is said to last for only 15 minutes. The Graminales includes two families: The Graminaceae, or grass family, and the Cyperaceae or sedge family. The grass family is by far the more important and is distinguished by the usually hollow aërial stems, two rows of leaves with sheathing bases split and the fruit is generally a grain (Fig. 284, 6). The sedges are largely paludose and of little value, the stems and leaves being often silicified. They usually have solid stems, three rows of leaves with sheathing bases entire, more simple flowers and the fruit is usually an akene (Fig. 285).

132. Arales, the Aroid Order.—These plants are largely tropical and are characterized by large and generally net-veined leaves and showy inflorescences and fruit (Fig. 286). They present an odd series of striking forms quite different from other monocotyledons. Many of them are familiar plants owing to their extensive cultivation in green-houses, as the calla lily (*Richardia*), *Caladium,* often with variegated leaves, and the gigantic leaves of *Dracontium* and *Colocasia,* known as the elephant ear, and the curiously perforated leaves of the *Monstera.* Many species of *Anthurium* are climbers and reach to the top of the highest trees, sending out with great regularity from the successive nodes, naked branches that may reach the ground and form roots. The order is represented in temperate regions by a few very familiar genera as jack-in-the-pulpit (*Arisaema*), skunk cabbage (*Spathyema*), golden club (*Orontium*), sweet flag (*Acorus*), water arum (*Calla*). The duckweeds that float upon every pond are extremely reduced allies of the order, representing the smallest seed plants known, *Wolffia* being an oval, rootless plant scarcely one millimeter in diameter.

The inflorescence is usually covered, as in *Typha,* with a bract

known as the spathe. This organ is variously colored and often the most attractive feature of the plant. The coloration and the odors and glands that are developed in some of the forms indicate that these plants are entomophilous. The flowers are arranged upon a fleshy axis, the spadix, and in the simplest forms



FIG. 286. Inflorescence of the Arales: *A*, habit of the Jack-in-the-pulpit. *B*, diagram of the inflorescence with spathe, *s*, opened on one side to show the spadix, *sp*, bearing staminate flowers at the base. *C*, a pistillate flower consisting of a single naked carpel. *D*, a staminate flower of four two-lobed anthers. *E*, section of the inflorescence of one of the arums—*p*, compartment containing pistillate flowers; *s*, staminate flowers.

are quite comparable to those of the *Typha*, consisting of one or a few stamens or a single pistil (Fig. 286, *B–D*). These imperfect flowers may be developed on different parts of the same spadix or on different spadices. In more advanced types, the

flowers are perfect, containing one or several pistils that may cohere and stamens which are surrounded by a very rudimentary perianth. The relation of these plants to insects is a very interesting one. The brightly colored spathe serves as an allurement and it also protects the microspores against wetting. The spores of comparatively few plants can endure water and it will be interesting to observe in the following lessons the devices that appear to guard against this danger. The spathe also affords shelter to the insect and the temperature is higher within the spathe since the food is being rapidly oxidized while the spores are forming. The temperature of the larger spadices may approach blood heat, being 10 to 25° C. higher than the outside air. Insects are not slow to avail themselves of these advantages and they also find an abundance of food in the microspores and glands that are often developed upon the spathe. Many aroids are characterized by the foul odors of decaying flesh and it is noteworthy that the coloration of the spathe is in singular harmony with these odors, resembling the hues of putrid meat. As a result of these variations, the lower orders of insects visit these flowers in great numbers and unwittingly serve in the transfer of the microspores; 250 carrion beetles and over 1,000 midges have been reported as taken from a single spathe. You can readily demonstrate that small insects are attracted to the spathe by examining the inflorescence of the skunk cabbage and jack-in-the-pulpit.

Common Examples of the Arales.—In the skunk cabbage (*Spathyema foetida*), which is of common occurrence along muddy streams, the inflorescence may appear as early as February, while the leaves are still rolled up into a compact bodkin to enable them to penetrate the soil. Later the leaves unroll and become very large, being good types of aroid leaves with broad blades, strongly net-veined and evidently adapted to abundant moisture and rapid transpiration. The spathe is shell-like, mottled with purple, green and yellow and encloses a fleshy oblong spadix that is entirely covered with flowers. You will find that small insects are attracted to these spathes in great numbers and spiders often take advantage of this fact and spin their webs

out of sight in the darker recesses of the spathe. In this plant, the flowers are decidedly in advance of those noted in *Typha,* being perfect, and in addition containing a perianth of four sepals which arch over the sporophylls. The crossing of the flowers is effected in a variety of ways. In some cases the inconspicuous stigmas first push out from between the sepals so that microspores must be brought from another earlier flowering plant. In other cases, the stamens at the top of the spadix are first extruded and shed their spores while at the bottom of the spadix the reverse condition obtains, the stigmas being in a receptive condition. This might result in a crossing of the upper flowers with the lower, and later, when the lower flowers put out their stamens and the upper their stigmas, insects could effect a reverse crossing. As the seeds mature the spadix and sepals become large and spongy, enclosing the ovaries which are finally set free by the decay of the spadix, as fleshy berries. This fruit is admirably adapted to the aquatic habit of these plants and readily floats in the water, owing to the spongy outer tissues, and so brings about the distribution of the seeds. The jack-in-the-pulpit has a surer method for effecting a crossing, since the sporophylls are usually borne on separate spadices. You will often find, however, spadices with a few stamens situated just above the pistillate flowers. The flowers are very simple, consisting of a single naked pistil or of four nearly sessile stamens (Fig. 286, *C–D*). The fruit is a shining bright red berry.

In some of the tropical genera the insects are held in the spathe by hairs that extend obliquely downwards, thus permitting the entrance but blocking the exit of the insects until crossing of the flowers has been effected. In one of the species of *Arum* there are two sets of hairs that divide the spathe into two compartments containing respectively staminate and pistillate flowers (Fig. 286, *E*). As soon as the spathe opens the insect ladened with microspores from another flower makes his way to the lower chamber, where the stigmas are in a receptive condition. Here he is held a prisoner and rubs the microspores upon the stigmas as he wanders about in the compartment. When the microspores in the upper chamber begin to discharge, the lower

set of hairs wither, giving him entrance to this chamber, where he feeds upon the microspores and becomes covered with them. Finally the upper set of hairs wither and he is free to leave the spathe and repeat his work in another inflorescence.

133. Liliales, the Lily Order.—The members of this order comprise nearly 5,000 species that are widely distributed and extensively cultivated for their showy flowers. We have now reached a point in the evolution of the flower where it has become perfect and the protective spathe of preceding orders is replaced by a well-developed perianth. The floral axis also becomes shortened and we pass from the spiral series of flowers with their variable number of organs to the cyclic flowers with a definite number of organs arranged in whorls. In the lily order there are five whorls of organs of three members each, the stamens being arranged in two whorls. Note also that this crowding usually results in the complete coherence of the carpels (Fig. 287, *C*). These plants are largely perennial, with underground stems in the form of bulbs or rhizomes. This feature adapts many of them to dry and steppe regions, from which source many of our cultivated lilies have been obtained. It will be seen that this habit of storing food in underground stems enables these plants to develop their leaves, flowers and fruit during the short rainy season, after which the entire aërial portion withers away and their life lies dormant in the buried stems. This habit is equally serviceable if these plants come into competition with larger forms, as in forests where plants with bulbs and rhizomes may complete their annual growth before the grosser vegetation that would crowd them out is fairly started (see page 39).

(*a*) *The Fawn Lily, Erythronium americanum.*—This species may be examined as typical of the order (Fig. 287). This plant has received the atrocious name of adder's-tongue, which is offensive and far-fetched, and also of dog-tooth violet, although it is not a violet at all. Burroughs has suggested the very appropriate name of trout lily, since the mottled leaves often form conspicuous beds on shady banks of streams; but to those who have experienced the spring time in the north country, the term fawn lily seems singularly appropriate. The leaves of the fawn

lily spring from deep-seated bulbs that are formed in a peculiar way. The seed germinates on the surface of the soil and forms a very small bulb and a single grass-like leaf. During each succeeding season a larger leaf and bulb are formed, and when of sufficient size, the bulb sends out one or more runners that penetrate the soil and develop new bulbs at their tips (Fig. 287, *B*).

FIG. 287. The fawn lily, *Erythronium americanum: A,* habit of the plant. *B,* the bulb, showing the origin of the stem and leaves shown in *A* —*r,* runners that penetrate the soil forming new bulbs at their tips. *C,* pistil of three carpels, at the right the fruit, a capsule, opening to scatter the seeds.

In this way the bulbs become deep-seated and rapidly increase in numbers, and after several years they attain sufficient size to develop two leaves and a flower. The mottled leaves have the same habit of rolling up in emerging from the ground, as noted in the skunk cabbage. It is to be observed that the position assumed by the mature leaf of many plants is often strikingly correlated with the extent of the root system. In the Liliales generally, which do not have extensive lateral roots, the hang of the leaves is such as to direct the water that falls upon them

towards the center of the plant, a feature doubtless of considerable advantage to plants living in semi-arid regions. The flower is of a decidedly higher type than any previously studied and it presents several features of special interest. The flower is laterally placed or slightly pendulous. The perianth is conspicuously developed, consisting of two whorls of three members each, though these organs are not as yet fully differentiated into calyx and corolla. The stamens are also arranged in two whorls of three members each and the three carpels form a single whorl which cohere into a compound pistil. The members of these whorls alternate with each other, so that the flower has three planes of symmetry. At maturity the walls of the ovary become papery and split down the side, thus freeing the seeds. This form of fruit is known as the capsule, and is of very common occurrence in the order (Fig. 287, *C*). The development of a conspicuous perianth is a noteworthy departure. It protects the microspores and therefore replaces the bracts and spathe of preceding orders, but owing to its peculiar coloration and form, it also serves as an attraction to special kinds of insects. In the preceding forms that attracted insects, the inducements were usually in the nature of shelter and food, which was largely the microspores that were offered freely to all. Such flowers are principally visited by a low order of stupid flies and beetles that are rather promiscuous feeders and are quite as likely to go from a flower of one species to that of a different kind, or to some other object and so defeat the principal object of the flower. With the development of the perianth, however, the flower is equipped with a device that primarily serves to exclude these less desirable visitors and to attract the more intelligent ones. In *Erythronium* the perianth is in a horizontal position, so that it offers a natural landing place for the insect, and the inducement is a sugary solution secreted by nectar glands concealed at the base of the perianth. It is an easy task for the more intelligent long-tongued insects, like the bees, wasps and butterflies, to reach the nectar in this type of flower. Such insects come to know by experience how to gather the food from a particular form of flower and consequently they will often confine their attention to

a single species during their entire flight; consequently the flower by excluding the less intelligent and slothful insects is more certain of being properly crossed. The development of nectar and also of odor glands are among the important variations that appear in the evolution of the flower. It is chiefly by means of the perfumes derived from the organs that the insect is directed to the flowers. The coloration is also of service, when the insect is near to the flower, thus supplementing the perfume glands by directing him to the proper entrance. Some of the larger and more brightly colored members of this and other orders apparently depend entirely upon the attraction of their colors, while other forms rely upon perfumes, the flowers being small and inconspicuous and often hidden. So we have reached a point in the evolution of the flower where it presents a number of variations that are adapted and of benefit to special kinds of insects. Insects have likewise varied and some have become of special use to flowers owing to their peculiar form. In this way certain insects and flowers have become dependent upon each other, and as a result of their mutually beneficial variations, the flowering plant and insects have greatly increased in numbers and exceeded all other groups in the plant and animal kingdoms.

(*b*) *Other Forms of the Lily Order.*—This order comprises an extensive series of plants that present nearly all the variations that occur in the flower. At the bottom of the series is the Rush family with grass-like stems and leaves (Fig. 288, *A, B*). The flower has the same structure as noted in *Erythronium,* but the perianth is composed of small scale-like leaves destitute of nectar glands, and as you would expect, is anemophilous. The Bunch flower family is a somewhat higher type. The flowers are entomophilous, though the perianth is small and the carpels are often but partially fused (Fig. 288, *C, D*). The flowers gain in conspicuousness, however, by being associated in compact inflorescences, but do not develop as a rule the bright colors characteristic of large flowers. The family is represented by *Tofieldia,* swamp pink (*Helonias*), blazing star (*Chamaelirium*), *Zygadenus,* hellebore (*Veratrum*), bellwort (*Uvularia*), autumn crocus, etc. The Lily of the Valley family is characterized by rather

larger flowers than those of the previous family, and they are less clustered and often provided with tubular perianths owing to the cohesion of the perianth segments. The fruit is a berry, unlike the majority of the families of this order. Here belong the asparagus, the false and true Solomon's seal, Indian cucumber, *Trillium,* and the closely related *Smilax,* etc. The Lily family includes nearly one half of the members of the order and

FIG. 288. Lower forms of the Liliales: *A,* a common rush, *Juncus,* showing grass-like appearance of the stem and inflorescence. *B,* a flower enlarged, showing the lily type. *C,* white hellebore, *Veratrum. D,* flower enlarged, showing the partial coherence of the carpels of this primitive type of the Liliales.

occupies a medium position in the evolution of the group. The flowers are large and of the *Erythronium* type. Here belong some of the most showy of our native and cultivated plants; the day lily, *Erythronium,* tulips, hyacinths, lilies, *Fritillaria,* onion, aloes and Spanish bayonets (*Yucca*) of arid regions. Among the higher genera of the lily family the organs of the perianth often cohere, forming a tubular structure. In the Amaryllis family we find the same type of flower and fruit as in the lily family, but the basal growth of the receptacle has enveloped the ovary so that the flower has become epigynous (Fig. 289, *A*).



FIG. 289. Advanced forms of the Liliales: *A, Narcissus* with inferior ovary, *o.* The six sepals cohere at their base, forming a tube and they also develop an outgrowth at the mouth of this tube, which surrounds the anthers and stigma like a cup. *B,* flower of *Iris. C,* section of same, showing inferior ovary, *o;* stigma, *s;* and anther, *a.*

This large family furnishes a number of showy flowers, as the jonquil and daffodil (*Narcissus*), amaryllis, snowflake, *Crinum,* star grass, century plant, etc. The flowers of some genera become irregular through the unequal development of certain leaves of the perianth. The Iris family marks the culmination of the variations noted in the order. The type of flower and fruit are like

those of the previous family, but the crowding on the receptacle has resulted in the obliteration of one whorl of stamens. The structure of the flowers often shows a series of variations that adapt them to insect visitors and crossing, as is well illustrated in the iris (Fig. 289, *B, C*). The styles have a very unusual form, resembling a leafy organ with the stigma on the upper side of a small projecting shelf. Beneath each of these curving styles is a stamen. The insect naturally alights upon the broad leaf of the outer whorl of the perianth and the peculiar coloration of this leaf probably directs him to the nectar secreted at the base of the perianth. In reaching this food he crowds down into the tube and rubs off the microspores upon his back. In leaving the flower he cannot hit the stigma, but in visiting another flower it can readily be seen that he will deposit some of the spores upon the stigma situated upon the top of the shelf, thus effecting a crossing. This order includes a large number of showy cultivated forms, as the iris, fleur-de-lis, crocus, gladiolus, friesia, blue-eyed grass, etc.

134. **The Scitaminales.**—This order is confined largely to the tropics and includes such familiar plants as the banana, the traveler's tree, ginger plant, canna, etc. It is mentioned here as furnishing an interesting illustration of the variations that often appear in epigynous flowers as a result of the crowding of the organs upon the receptacle. Tendencies toward irregularity appeared in the epigynous families of the Liliales, but in this order these variations become very marked and make less abrupt the transition to the next order. The flowers of the banana are suggestive of the amaryllis family, though somewhat irregular. All of the perianth leaves are united but one and only five perfect stamens are developed. These flowers are united into groups in the axils of bracts that form large buds at the ends of the stem. As the bud elongates, the basal or lower bract is first curved back from the bud, thus exposing the flower cluster, and this expansion goes on until all the flowers are exposed. The lowest flowers in the inflorescence are imperfect, containing only pistils, the central are perfect and the upper ones bear only stamens. Only the ovaries of the lower flowers of the inflorescence develop,

each "hand" of bananas representing the matured ovaries of the flowers in the axil of a bract. Singularly, seeds do not develop in the edible banana, although you can see minute dark-colored ovules forming three radiating lines in a cross-section of the fruit. The ability of these plants to propagate themselves by means of buds developed on the underground rhizomes may have resulted in the loss of the seed habit. A great many plants are so successful in propagating themselves by buds, bulbs, runners, etc., that they have ceased to produce seed. In the higher members of the order, as in the ginger and canna families, the flowers become very irregular through the unequal development of the leaves of the perianth or in some forms on account of the abor-



FIG. 290. Flower of *Canna*, the showy petal-like organs being modified stamens (staminodia) while the perianth proper, *p*, is reduced to green bract-like organs: *l*, labellum; *an*, anther on petal-like organ; *s*, stigma; *o*, ovary. *B*, section of flower, showing ovary, *o*, the modified stamen, *an*, and stigma, *s*, the other organs being removed.

tion or transformation of all the stamens save one into petal-like organs termed staminodia. Usually one of the leaves of the perianth, or one or more of the staminodia, are highly modified and known as the labellum. This organ is so placed as to afford a natural landing place for the insects visiting the flower and also to necessitate crossing (Fig. 290). Observe a bee visiting the

flowers of the canna and determine the significance of the position and movement of the labellum and its relation to the stamen and stigma.

135. Orchidales, the Orchid Order.—The orchids are the highest group of the monocotyledons and their flowers are a source of wonder and admiration, owing to the singular beauty and delicacy of their mechanical construction. Variation in this order has occurred on a gigantic scale, resulting in a larger number of species (over 7,000) than is found in any of the preceding orders. Nevertheless these elaborate variations have not been very successful in enabling them to compete with other plants, and as a result the orchids are rather rare and not at all comparable in number of individuals with the lilies and grasses. Though more widely distributed than any of the other monocoty-

FIG. 291. An epiphytic orchid growing upon the branch of a tree. The coarse roots, *r*, are surrounded by a mantle of cells which take up the moisture from the atmosphere. *b*, storage organs formed from the base of the leaves, enabling the plant to produce flowers and fruit. The smaller stalks, *a*, are the shrivelled remains of these organs after flowering and fruiting.

ledons, their variations have adapted them as a rule to peculiar conditions. They are especially abundant in the mountainous districts of the tropics, where they more commonly appear as epiphytes upon the trunks of trees and in the crevices of rocks. Such conditions are met by the development of a thick mantle of cells, the velamen, about the aërial roots which absorbs the moisture from the air and doubtless the enlargement of the leaf base into a bulbous storage organ enables these plants to anticipate in this way the heavy demands that will be made upon them in the flowering season (Fig. 291). The terrestrial forms are largely parasitic or saprophytic and associated with mycorrhiza, and this has resulted in some of the forms in the suppression of various organs of the plant as the primary root or even the entire root system, as is illustrated in the coral root orchid, where the leaves have also become reduced to mere scales and the chloro-



FIG. 292. A simple type of the Orchidales: *A*, the moccasin flower, *Cypripedium*—*l*, labellum; *p*, the two unmodified petals; *s*, sepals, two being united below the labellum; *b*, bract, partially concealing the inferior ovary. *B*, section of the flower—*l*, labellum; *s*, stigma; *an*, anther; *st*, shield-like sterile stamen covering the two anthers and stigma; *o*, ovary; *b*, bract.

phyll has disappeared. The organs of the flowers are subject to such remarkable variations that the various parts may appear at first somewhat difficult to recognize. It is evident that the same line of variation noted in the higher families of the Liliales has

been continued, for the perianth is arranged upon a compound ovary, but its parts are often sharply differentiated into calyx and corolla. The sepals and petals comprising these two whorls differ greatly in form, but especially to be noted is the oddly constructed petal known as the labellum (Figs. 292, 293, *l*). This organ is the most striking feature of the orchid and it assumes an almost endless variety of forms and colorations, being a good illustration of Wallace's law that the most highly modified part shows the greatest variation in coloration. In *Cypripedium* (Fig. 292, *A*) the labellum assumes the form of a moccasin, in other genera it resembles a vase, boat, tongue, body of insect, etc. It is entire or variously lobed, slit, fringed and often prolonged into a tube for the concealment of nectar. The stamens and stigmas are reduced in number and greatly modified, the former organs usually being reduced to one and so fused with the style that the anther is sessile upon it (Fig. 292, *B*). One or two of the stigmas are generally modified into a mucilage-secreting organ, called the rostellum (Fig. 293, *A*). The microspores are usually united into a waxy mass, pollinium (pl. pollinia) and attached to a sticky part of the rostellum (Fig. 293, *B, C*). The remarkable feature about these gaudy flowers is the relation that the labellum, anther, rostellum and stigma sustain to each other. The position of these organs is such that an insect visiting the flower touches with some part of his body the sticky part of the rostellum and the pollinia are thus made fast to him and carried to the stigma of another flower. These devices are so elaborate in many orchids that the microspores can only reach the stigma through the agency of an insect. In the lower types of orchids, as the moccasin flower (Fig. 292, *B*), a somewhat different arrangement is found. The bee enters the opening in the upper part of the labellum and feeds upon the glands distributed along the bottom. In leaving the flower he forces his way through the small opening on either side of the style and so he first comes in contact with the stigma and later with the anthers, which are two in number and located back of the stigma. In this flower the anthers are surrounded by a sticky mass and the microspores are thus fastened to the insect's body to be carried to another

flower. In the higher types of orchids, the insect probing for the nectar touches the sticky discs of the rostellum, to which the pollinia are attached, and in this way they are fastened to his body and drawn out of the open anthers when he leaves the flower (Fig. 293, *C*). In some cases the pollinia quickly curve after being withdrawn from the anther, with the result that this change of position brings them into line with the stigma of the next flower visited. In other genera certain cells are irritable



Fig. 293. Higher type of the Orchidales: *A*, flower of *Orchis*—*l*, labellum; *p*, the two unmodified petals; *s*, sepals; *r*, rostellum to which the two pollen masses, pollinia, in the two-lobed anther, *an*, are attached by sticky discs; *st*, stigma. *B*, enlarged lateral view of the anther which has opened, exposing the pollinia. *C*, one of the pollinia withdrawn from the anther, showing adhesive disc at base which is attached to the rostellum. At right a pollinium enlarged, showing the small masses of microspores, massula, that may be separately detached from the pollinium.—After Warming.

and in a high state of tension. A touch causes an explosion that results in hurling out the pollinia which always land on the end to which the sticky disc is attached and so become fastened to the insect. (See Darwin's Fertilization of Orchids for a discussion of the multiplicity of these arrangements.)

The seeds are among the most rudimentary of the Spermatophyta, consisting of an undifferentiated, few-celled embryo with-

out endosperm and surrounded by a delicate bladder-like integument. They are exceedingly small and produced in enormous numbers; in fact, the seeds of the rattlesnake orchid, *Peramium*, weigh but two millionths of a grain each and float in the air like dust particles. The fringed orchis (*Blephariglottis*), *Arethusa*, ladies' tresses (*Gyrostachys*), rattlesnake plantain (*Peramium*), grass pink (*Limodorum*), rose pogonia, showy orchis (*Galeorchis*), moccasin flower (*Cypripedium*) are among the common and more showy of our native orchids.

Class B. Dicotyledones

136. General Characters.—The structure of these plants is more complex than that of the monocotyledons and their variations have been more extensive and successful, over 100,000 species being known. For this reason they are adapted to a greater range of conditions and have become the dominant plants, forming the conspicuous and characteristic features of the vegetation of the earth.

The dicotyledons may be short-lived annual plants, or perennials, and they include a great variety of climbers, epiphytes, parasites, and saprophytes and comparatively few aquatics. The most extreme form of xerophytes are also found in this group. The leaves are highly differentiated, usually consisting of a blade and petiole which is often associated with small leaf-like organs, the stipules (Fig. 294). The blade varies in form but is often characterized by teeth and various forms of lobing, owing to the fact that the veins usually differ from those of the monocotyledons in that they repeatedly branch, becoming smaller and smaller and thus forming a network or reticulated venation with free ends on the margins and other parts of the leaf. The stems are markedly different from the previous class, owing to the arrangement of the vascular bundles in a circle about the pith and the formation of a cambium zone which brings about an increase in the diameter of the stem (Fig. 295). This arrangement gives the plant a great advantage, permitting the extensive system of branching that characterizes the group and the consequent increase in the display of foliage.

The flower is subject to the same modifications as noted in the monocotyledons. Among the lower orders, the flowers are quite as simple as those of the primitive monocotyledons and the development of imperfect and anemophilous flowers is of common occurrence. The spiral arrangement of the numerous organs

Fig. 294.



Fig. 295.

Fig. 294. Leaf of white birch, the blade, *b*, traversed by a network of veins that end in free branches (the margin irregularly toothed or dentate) and supported upon a petiole, *p*.

Fig. 295. Diagram of a cross-section of a stem of black oak four years old: *p*, pith; 1, 2, 3, 4, annual rings of xylem; *c*, cambium zone; *ph*, phloem; *c*, cortex; *ck*, cork; *m*, medullary rays.

of the flower will also be noted. The majority of the orders of the dicotyledons, however, are characterized by perfect, cyclic flowers and the various sets of organs usually consist of four or five members each. The perianth, when present, is generally differentiated into a green calyx and a variously colored corolla. The stamens are more frequently arranged in one or two whorls, equalling or twice the number of the sepals, and the pistils usually form a single whorl, equalling or less than the number of sepals. The crowding of the various organs, and the lateral growth of the receptacle results in the reduction in the number of organs and in their cohesion with one another so that in the higher types

the calyx and corolla become more or less tubular and the carpels unite, forming a compound ovary. The cohesion of the organs is often associated with other variations of the flower. Among the more common of these changes is the adhesion of the stamens to the corolla, which is usually associated with the basal growth



FIG. 296. Structure of dicotyledonous seeds: *A*, nearly mature seed of *Lepidium*. The embryo consists of the hypocotyl, *hy*, ending below in the root, *r*, and the root cap and above in the epicotyl or plumule, *pl*. Two cotyledons, *c*, arise laterally from the stem; *f*, funiculus; *mi*, micropyle; *in*, integuments; *en*, remains of endosperm. *B*, section of seed of water lily —*e*, embryo with two cotyledons attached laterally to the minute stem of the embryo and surrounded by a layer of endosperm cells; *mg*, sporangial cells or perisperm; *i*, integument.

of the receptacle, forming the perigynous type of flower, and the adhesion of the receptacle to the ovary so that the perianth and stamens apparently arise from the top of the ovary, epigynous type of flower (page 353). The irregularity that characterizes the highest types of flowers in many of the orders is the most helpful of all these variations, as is indicated by the abundance and more common occurrence of such forms.

The most characteristic feature of the dicotyledons is the

embryo which usually consists of a root, stem, and two laterally attached cotyledons. The region of the stem above the cotyledons is known as the epicotyl and frequently appears as a minute bud, the so-called plumule. The region of the stem below the cotyledons, the hypocotyl, terminates in the root (Fig. 296). The elongation of the basal portion of the hypocotyl frees the root from the seed and the growth of the upper region of the hypocotyl pushes up into the air the cotyledons and growing point or epicotyl. The formation of the stem is sometimes due to the elongation of the epicotyl alone, the cotyledons frequently containing the storage foods for the nourishment of the young plant and remaining buried in the seed.

In endospermous seeds it should be noted that the cotyledons are not developed as digestive and absorbing organs as observed in the monocotyledons (page 122). Unlike the latter group, the primary root often persists, forming the main or tap root of the plant. The Dicotyledons include two rather distinct series: *a.* The Choripetalae, distinguished by their free petals or lack of perianth. *b.* The Sympetalae, with united petals.

Series a. Choripetalae

137. General Characters.—This group comprises about 60,000 species and includes the majority of the trees and shrubs found in the temperate regions. The flowers as a rule are of a simple type but exceedingly variable in structure, so that it is not possible to separate them into so sharply characterized orders as in case of the monocotyledons, or in the following group of Sympetalae. Furthermore, these orders doubtless represent many parallel lines of development that are imperfectly understood. For this reason, the order of the presentation in the following pages does not attempt to represent the real relationship of the groups.

138. Salicales, the Willow and Poplar Order.—The willows are almost universally distributed along water ways, apparently demanding for their existence only light and water. The poplars are less restricted and some can endure moderately arid condi-

tions. Few plants have greater vitality. A bit of a twig and often a portion of a root is capable of developing buds and so starting the shoot. Trees are often pointed out that have originated through the careless sticking of a twig in the soil and several of the willows are naturally propagated by their twigs which are easily broken off by the winds. Their unusual ability



FIG. 297. Flowers and seed of the willow (*Salix*): *A*, winter appearance of a flowering twig, each boat-shaped scale concealing an ament. *B*, pussy willow stage of flowering, the aments emerging from the scales and exposing the hairy bracts that conceal the flowers. *C*, ament of pistillate flowers in full bloom. *D*, pistillate flower, consisting of a compound pistil of two carpels—*b*, bract; *n*, nectar gland. *E*, ament of staminate flowers. *F*, staminate flower of two stamens. *G*, ament of mature pistils which are opening to discharge the seeds. *H–I*, successive stages in the opening of the pistil. *J*, a seed with circle of hairs at base forming a parachute for dissemination.

to form numerous buds is well shown in the pollarded willows where the branches have been cut back and a large number of shoots develop about the wound. Perhaps the stimulus of the wound also awakens some of the dormant buds (page 100). The flowers, as in *Typha*, are arranged on an elongated axis forming

a compact inflorescence known as an ament or catkin (Fig. 297, *C, E*). The distinguishing feature of this kind of flower cluster is seen in the scale or bract that protects each flower (Fig. 297, *D, F*). The aments and also the leaves are concealed in buds that are peculiar in that they are protected by a boat-shaped scale (Fig. 297, *A*). These plants flower very early in the spring and as the ament emerges from the bud the overlapping bracts with their hairy coats form the "pussy willow" stage of the inflorescence (Fig. 297, *B*). Soon the bracts spread apart, exposing the flowers which are nearly as primitive as the simplest of the monocotyledons, being without perianth, imperfect and usually the two kinds of sporophylls are arranged on different plants. The pistil is compound and composed of two carpels, each of which contains numerous ovules, so that the pistillate flowers are not as primitive as those of the cat-tails, where the pistil is simple (Fig. 297, *D*). The staminate flowers show the same primitive characters (Fig. 297, *F*), consisting of one or more naked stamens. The absence of showy perianth, the extravagant production of microspores and the formation of the flowers before the leaves become large and so interfere with the distribution of the microspores are all characteristics of anemophilous flowers. It is noteworthy, however, that nectar glands are developed in the flowers of the willow (Fig. 297, *D, n*) and that the microspores are sticky, indicating that these flowers are entomophilous. Perhaps we have here an illustration of one of the earliest variations of the flower that served as an allurement to insects. Certainly, the nectar glands and the conspicuous display of microsporophylls are a very efficient attraction, as is attested by the variety of insects that swarm about the aments.

The most efficient factor in the distribution of the willows and poplars is found in the seed. The megasporophylls mature in the early summer, when the two carpels spread apart (Fig. 297, *G–I*), permitting the discharge of the seeds, which are provided with a circle of hair at the base (Fig. 297, *J*). This parachute is not so nicely constructed as in *Typha,* but it is so efficient that myriad numbers of minute seeds are carried a considerable distance and cover everything in the neighborhood of the trees

with their lint-like masses. The name cottonwood is popularly applied to several of the poplars because of the cotton-like clusters of seeds that emerge from their aments.

139. Fagales, the Beech Order.—This order includes many of the most important hard-wood trees of the temperate regions, comprising the family of the birches, with such representatives as the American hornbeam (*Carpinus*), hop hornbeam (*Ostrya*), hazel (*Corylus*), birch (*Betula*), alder (*Alnus*) and the beech family, which includes the chestnut (*Castanea*), beech (*Fagus*) and oak (*Quercus*). The inflorescence is more commonly an ament as in the preceding order (Figs. 298, *A*; 299, *A*), although



Fig. 298. Flowers and fruits of the birch family, order Fagales: *A*, inflorescence of hornbeam (*Carpinus*)—*s*, staminate ament; *p*, pistillate ament. *B*, staminate clusters from ament of alder (*Alnus*), each flower consisting of four stamens attached to a four-parted perianth, *pr*. *C*, upper view of the same cluster, showing the numerous bracts associated with the flowers. *D*, fruit of *Carpinus* attached to the greatly enlarged three-lobed bract. *E*, flower cluster of hazel (*Corylus*)—*b*, bract encircling the ovary and within a more delicate bract, *pr*, adnate to the ovary; *s*, stigma. *F*, fruit of *Corylus*, the bract, *b*, in *E* has grown out into a tubular beaked structure that completely envelops the nut.

in the hazel and in the beech family only one or a few pistils are developed in the bud-like clusters of overlapping bracts (Fig. 299, *mg*). The flowers are imperfect and the two kinds of sporophylls are usually developed upon the same plant. Several bracts are usually associated with sporophylls (Figs. 298, *B–F*; 299, *B–E*) so that the flowers are of a higher type than the willows. The innermost of these bracts is often of a delicate structure and has been referred to as a primitive form of the calyx (Fig. 298, *B, pr*) and when present in the pistillate flowers it adheres to the ovary (Figs. 298, *E, pr*; 300, *B, pr*). The pistils are provided with long delicate stigmas and are compound, containing several ovules but only one usually matures. The wall of the ovary develops into a tough coat about the seed, forming a fruit known as the nut. Certain bracts of the flower increase greatly in size as the ovary matures and form a conspicuous part of the fruit. Thus in the hornbeam (Fig. 298, *D*), one of the bracts develops into a large, three-lobed green leaf, in the hop hornbeam the bract forms a papery sac about the nut, in hazel a leafy husk (Fig. 298, *F*), in the birch and alders a woody peg-like structure. In the chestnut and beech the pistils are completely enveloped by prickly bracts or outgrowths that form the bur (Fig. 300, *A–C*), while in the oak these structures only cover the lower portion of the ovary, forming the cup (Fig. 299, *C–E*). The flowers of this order are typical of that great group of plants comprising about one tenth of all the flowery plants, which are called anemophilous. Note the small and simple flowers, absence of showy perianth, nectar, and odor glands, dry and light microspores lavishly produced to ensure crossing and the delicate bushy stigmas for catching the spores. The stigmas appear a day or so before the adjoining microspores are being shed so that a crossing from an earlier flowering plant is necessitated. These plants form the larger part of our deciduous forests and their association in colonies is doubtless connected with their anemophilous habit, as is also the appearance of the flowers before the leaves are fully developed. The positions assumed by the staminate aments is of service in protecting the microspores against wetting and also to assist in their distribution. During

the winter, these aments are quite erect but as the flowering stage approaches they become pendulous, the larger bracts protecting the sporophylls, like the shingles on a roof (Fig. 298, *A*). These bracts, being hygroscopic, remain closed during wet weather but on dry days they curve back, each bract forming a shelf

Fig. 299.

Fig. 300.

Fig. 299. The beech family, order Fagales: inflorescence of oak (*Quercus*)—*s*, staminate ament; *p*, pistillate inflorescence. *B*, staminate flower surrounded by a perianth of slightly united bracts. *C*, pistillate flowers with numerous bracts surrounding base of ovary. *D*, section of flower, the pistil being composed of three carpels and the inner bracts adnate to the ovary. *E*, fruit of oak, the cup consisting of the modified outer bracts shown in *C* and *D* and the nut has developed from the ovary and one of its ovules.

Fig. 300. Flower and fruit of the beech (*Fagus*), order Fagales: *A*, pistillate inflorescence, the three-lobed stigmas projecting beyond the bracts. *B*, section of the inflorescence—*pr*, inner bracts or perianth surrounded by an outer spiny set. *C*, the fruit, the outer bracts of *B* have become hard and spiny and are splitting into four valves, exposing the three-angled nuts.

which serves to catch the spores when no winds are stirring and thus prevent their falling to the ground.

140. Other Tree Orders Suggestive of the Fagales.—Closely allied to the Fagales is the order of the walnuts, Juglandales, including the black walnut and butternut (*Juglans*) and the hickories (*Hicoria,* Fig. 301). These trees are of very common

FIG. 301. Flower and fruit of the Juglandales: *A,* inflorescence of butternut—*s,* staminate ament; *p,* pistillate inflorescence. The butternut and black walnut (*Juglans*) may be recognized by the chambered pith, as shown at bottom of twig. *B,* section of a pistillate flower—*pr,* perianth adnate to the two carpels composing the pistil; *s,* stigma. *C,* fruit of hickory, the outer part of ovary wall splitting into four valves and exposing the hard inner part, the shell of the nut.

occurrence in the northern United States and characterized by their aromatic oils and large compound leaves. The flowers are very similar in structure and arrangement to those of the beech order, but at maturity the outer part of the wall of the ovary becomes transformed into a fleshy rind and the inner portion forms a hard shell. In the walnut this pulpy, aromatic rind

about the nut is finally destroyed by decay, while in the hickories the rind becomes leathery and splits into valves, freeing the nut (Fig. 301, *C*).

The order of the nettles, Urticales, also has many points in common with the beech order and contains several of our common trees as the elm (*Ulmus,* Fig. 302), hackberry (*Celtis*), mulberry (*Morus*), osage orange (*Toxylon*), numerous tropical forms, as the India rubber trees, banyan tree, etc., as well as a variety of valuable herbaceous plants, as the hop and hemp. These plants very generally contain a milky juice, latex, which in the case of *Ficus* and *Castilloa,* two of the India rubber trees, furnishes the raw material of India rubber and in the cow tree of South America yields a saccharine, nutritious milky juice. Tough stereome fibers are characteristic features in many of these plants and furnish the hemp derived from the *Cannabis,* and in Japan paper is manufactured from the fibers of the paper mulberry. The leaves are often provided with rough hairs which sometimes contain irritating acids, as in the nettles (page 35). The flowers are of a somewhat higher type than in the preceding orders, and associated in a variety of dense inflorescences. They are rarely perfect, though rudimentary (sterile) stamens may be developed with fertile pistils or *vice versa,* thus giving the appearance of perfect flowers (Fig. 302, *B–D*). The calyx does not adhere to the ovary as in the Fagales, though the sepals often cohere (Fig. 302, *D*). The ovary usually matures as a nut, but in many cases a kind of drupe (page 365) is formed, owing to the perianth becoming fleshy and enveloping the nut. The fruit of the mulberry is an aggregation of drupes. In the fig the stem envelops the minute curious flowers and becomes fleshy, forming the edible portion of the fruit.

141. Orders Showing an Advance over Preceding Forms.— The buckwheat order, Polygonales, containing such familiar weeds as the sorrel, dock (*Rumex*), and knotweed or smartweed (*Polygonum*), has perfect regular flowers with a distinct calyx. These characters also appear in the allied goosefoot order, Chenopodiales, with its great array of common weeds, as the goosefoot (*Chenopodium*), orache (*Atriplex*), tumbleweed and pigweed

(*Amaranthus*). The flowers are arranged in loose clusters and are for the most part small and anemophilous (Fig. 303). However, it is noteworthy that some of the genera are entomophilous, the calyx becoming larger and brightly colored and nectar glands

Fig. 302.

Fig. 303.

Fig. 302. Inflorescence and fruit of elm, order Urticales: *A*, twig of elm bearing principally staminate flowers. *B*, a staminate flower enlarged, showing lobed perianth enclosing numerous stamens. *C*, pistillate flower, two-lobed stigma projecting from perianth. *D*, section of pistillate flower, showing aborted (sterile) stamens and pistil (fertile) of two carpels, as indicated by the two stigmas. *E*, the fruit and perianth still attached. A wing has grown out from the sides of the ovary.

Fig. 303. A simple type of the Chenopodiales: *A*, shoot of Mexican tea (*Chenopodium*), showing character of inflorescence, *in*. *B*, flower enlarged, showing the perianth, stamens and pistil of three cohering carpels.

are associated with it. In the higher members of the order, as in the purslane and pink families (which includes the spring beauty, *Claytonia*), the perianth becomes differentiated into a calyx and showy corolla, and the flowers have become in a marked degree adapted to insect visitors (Fig. 304). These orders form a natural transition from the primitive flowers of the willows and beeches to the large flowers of the next order with their showy perianths, though the structure of the flower does not indicate a relationship between them.

142. Ranales, the Buttercup or Crowfoot Order.—This large

and interesting order includes a great variety of our common plants, herbs and trees, as the white and yellow water lilies (*Nymphaea* and *Castalia*), buttercups (*Ranunculus*), marsh



FIG. 304. Advanced type of the Chenopodiales: *A*, shoot of *Melandrynum* bearing flower with perianth differntiated into calyx, *ca*, and corolla, *c*. *B*, section of flower, showing the relation of calyx to corolla and the concealment of the nectar glands at the base of the ovary. Access to the corolla tube is guarded by an outgrowth on the petals, as shown in *C*, which, assisted by the styles or stamens, so effectually closes the mouth of the tube that only insects with long tongues can reach the honey.

marigold (*Caltha*), windflower (*Anemone*), *Hepatica*, rue (*Thalictrum*), columbine (*Aquilegia*), larkspur (*Delphinium*), monkshood (*Aconitum*), may apple (*Podophyllum*), magnolia, tulip tree (*Liriodendron*), *Sassafras*, spice bush (*Benzoin*), etc. In this order we have again reached the point, just as in the monocotyledons (see Liliales), where the flowers are more usually solitary and conspicuous, owing to the development of large, showy perianths that are more frequently differentiated into calyx and corolla. The flowers, however, are very simple, as is indicated by the regular and hypogynous arrangement of the parts (Fig. 305). The receptacle is frequently elongated

and consequently the organs of the flower, especially the sporophylls, are numerous and indefinite in number and spirally arranged. Were it not for the perianth, such a type of flower would be quite as primitive as any of the preceding orders.

(*a*) *The Buttercup, Ranunculus.*—The flower of the buttercup illustrates the more characteristic features of the Ranales (Fig. 305, *A, B*). The calyx and corolla form two whorls of five members each, but the sporophylls are indefinite in number and spirally arranged. It is interesting to note the relationship of the various parts of this simple flower and their behavior during

FIG. 305. A common type of the Ranales, *Ranunculus repens: A,* habit of the plant. *B,* early stage of flowering, the stamens clustered about the stigmas. *C,* late stage of flowering, nearly all the stamens bent over towards the petals, having discharged their spores. *D,* petal with nectar gland at base. *E,* fruit consisting of numerous spirally-arranged akenes.

the period of blooming. When the flower first opens, the anthers are still closed and clustered around the receptive stigmas, affording a natural landing place for the insects (Fig. 305, *B*). Thus crossing may only be effected with the spores carrried from older flowers. At a later period the outermost filaments elongate, curving over towards the petals, so that as their anthers discharge there is little chance of autogamy (Fig. 305, *C*). This position also brings the anther into the pathway leading to the nectar gland at the base of the petal (Fig. 305, *D*), so that the insect

in probing for the nectar is sure to receive some of the sticky microspores on his body. Each day, for about a week, successive series of stamens behave in this manner, and it is not until the innermost anthers are discharging on the last day of blooming of the flower that there is an opportunity for autogamy. In many members of this and other orders you will find that the stigmas and anthers of the innermost stamens actually meet, owing to the peculiar curvature of the styles and filaments, thus ensuring autogamy in case crossing has failed. It should be borne in mind that the devices for effecting autogamy, as a last resort, are quite as general and often as elaborate as the provisions for bringing about crossing. The microspores are protected against dews and rains by the folding of the perianth and also by the downward curvature of the pedicel at night, while on each succeeding bright day the perianth opens again and the flower becomes nearly or quite erect and turned towards the light, so that it is in a position from which the insect will naturally come. The opening of the flower is due to a growth in the morning of the basal, inner portion of the petals which bends them away from the center of the flower, thus opening the flower; while a corresponding growth towards evening of the basal outer part of the petal bends them towards the center of the flower. This growth slowly increases the size of the flowers, as may readily be observed by comparing the freshly opened flowers of the anemone, buttercup, etc., with those several days old. All these features are of common occurrence in a great variety of orders. The fruit of the buttercup is an akene, each carpel containing but a single ovule (Figs. 305, *E*; 261 *B*).

(*b*) *Some Variations of the Order.*—Many variations of this simple structure appear in other members of the order. The carpels may be reduced to one, as in sassafras and other groups, and in the green hellebore (Fig. 306, *A*) they are partially fused and even united with the receptacle in some water lilies, etc. (flowers perigynous and epigynous). The stamens are sometimes few in number and they are frequently modified into nectar glands and assume a variety of odd shapes (Fig. 306, *A*). The showy nectar-bearing petals, honey leaves, of many of the genera

are regarded by some as modified stamens, as in the monkshood, columbine, larkspur, etc. (Fig. 306, *B*).

The parts of the perianth, which are more frequently in three's or indefinite in number than in five's are also subject to consider-

FIG. 306. Modifications appearing in the Ranales: *A*, flower of *Helleborus—n*, nectar glands due to modifications of stamens. *B*, columbine—*n*, honey leaves due to modifications of the petals or possibly of the stamens.

able variation. Very frequently the corolla is not developed or partially suppressed, and the sepals become colored and petaloid, as in the marsh marigold, hepatica, certain windflowers, rue, spice bush, larkspur, etc. In a few of the genera the perianth becomes greatly modified and even irregular, as in the columbine (Fig. 306, *B*, *n*), where five of the petals, often regarded as modified stamens, are transformed into tubular honey leaves or into spurs and hoods, as in the larkspur and monkshood (Fig. 307, *B*, *D*). These flowers illustrate very well the progressive coloration that is often associated with the variation of the perianth. The simpler forms are usually yellow or white, while flowers with more highly modified parts are pink and pale blue (columbine), blue in higher types (larkspur) and ultra-marine blue in the most irregular and highly modified type (monkshood). Any of these flowers may take on by reversion the lower grades of color, so that a blue-colored form may vary and its offspring may produce white or yellow flowers, as in the colum-

bine. The remarkable brilliant yellow of the flowers, as seen in the buttercup and marsh marigold, etc., has been regarded as the primitive coloration of the perianth, and it is to be noted that the flowers showing reversion to this color assume a duller



FIG. 307. Highest types of the Ranales: *A*, inflorescence of monkshood. *B*, section of a flower—*s*, helmet-like sepal enclosing nectar organ, *n*. *C*, flower of larkspur. *D*, section of flower—*s*, spurred sepal enclosing two-spurred nectar organs, *n*. *E*, fruit of larkspur consisting of 3 pistils maturing as follicles.

yellow. This order furnishes a great variety of showy ornamental plants, but none with fragrant flowers. Many contain acrid juices and poisonous alkaloids, some of which are of medicinal value, as aconite, hydrastin, helleborine, etc. Others are of value for the volatile oils, as sassafras, cinnamon, camphor, nutmeg, etc. The simple structure of the more characteristic flowers indicates that this order is a very ancient one and the pronounced tendency to vary would suggest that possibly many of the higher orders of the Choripetalae have been derived from this group. The origin of the monocotyledons from this order has also been suggested, owing to the structural features of the embryo and vascular bundles of certain groups.

143. Papaverales, the Poppy Order.—This group includes two very well known families: (1) The poppies, with such familiar plants as the poppy (*Papaver*), sea poppy (*Glaucium*), celandine (*Chelidonium*), soldier's cap (*Bicuculla*), fumitory (*Adlumia*), corydalis (*Capnoides*) (Fig. 308). (2) The mustards, including the peppergrass (*Lepidium*), hedge mustard (*Sisymbrium*), white and black mustard (*Sinapis* and *Brassica*), yellow rocket (*Barbarea*), cress (*Roripa*), toothwort (*Dentaria*), shepherd's purse (*Bursa*), whitlow grass (*Draba*), rock cress (*Arabis*) (Fig. 309). The families of the caper and mignonette are also included in the order. These plants are closely related to the Ranales, as is indicated by their simple and showy flowers, the various organs usually being in multiples of two and quite distinct (flowers hypogynous) save in the case of the carpels. These organs are united as in the water lilies and more commonly reduced to two, thus forming a sharp contrast with the preceding order.

(*a*) *The Bloodroot, Sanguinaria.*—This plant is typical of the simpler members of the poppy family (Fig. 308). These plants form colonies in rather open, rich woods owing to their fleshy rhizomes, and like many plants with storage organs, flower very early in the spring. The flowers endure for about two days, but the large-lobed leaves are conspicuous features of the forest carpet for several weeks while they are manufacturing the food for the next season. The large leaves are tightly rolled about the solitary flower and enveloped in a papery sheath as they emerge from the soil. A juice, latex, is found in the majority of the plants of this order, and in this example it is of a blood-red color, thus giving the latin and popular name to the plant, *Sanguinaria,* or bloodroot. The flowers are of the simple open type noted in the buttercup, having two sepals which fall with the opening of the flower, petals 8–12, stamens indefinite and a compound pistil of two carpels, as is indicated by the two stigmas and two rows of ovules (Fig. 308, *C*). The stigmas are receptive as soon as the flower opens, but the anthers remain closed, and so the spores must be carried from an older flower. The petals close during the late afternoon and when opened on

the following day the anthers shed their spores and complete the flowering.

In higher types of the poppies we find the flowers becoming irregular and the various parts reduced in numbers. Thus, in the soldier's cap (Fig. 308, *D*), there are two scale-like sepals, four petals in two pairs, the two outer heart-shaped and spurred at the base, while the two inner are narrow and winged on the back, enclosing the sporophylls. The six stamens are somewhat united and arranged into two groups opposite the spurred petals. The carpel has the same structure as in the bloodroot. It is evident that the nectar concealed in this closed type of flower can only be secured by some long-tongued insect. Find out how



FIG. 308. Examples of the poppy family, order Papaverales: *A*, habit of the bloodroot at time of flowering—*r*, rhizome; *l*, leaf folded about flower stalk. *B*, leaf unfolded. *C*, section of flower, showing its hypogynous structure with numerous stamens and pistil of two carpels. *D*, flower of fumitory (*Adlumia*)—*s*, sepal; *p*, inflated petals; *w*, winged petals which conceal the sporophylls.

the bee procures the honey, the purpose of the wings on the narrow petals and the position of the nectar glands. Is the stigma mature before the anthers open? The structure of this higher type of the poppies makes the transition to the mustard family a simple one.

(*b*) *The Mustard Family, Cruciferae.*—Here we find the sepals and petals four in number, and the latter arranged crosswise (Fig. 309, *A, B*), thus explaining the family name, Cruciferae. The petals usually have rather long claws (Fig. 309, *D*)



FIG. 309. Example of the mustard family, order Papaverales: *A*, branch of *Brassica*, one of the turnips. *B*, a flower enlarged. *C*, flower in section. *D*, a petal, showing narrow claw and broad blade. *E*, pistil. *F*, fruit, the valves of the two carpels opened to show the seeds.

that are so associated with the sepals as to form a narrow tubular perianth that effectually conceals the nectar glands situated at the base of the ovary. There are usually six stamens, but the

two outer ones are shorter than the others (Fig. 309, *C*). The ovary is divided into two compartments by a membranous partition. The fruit is generally a capsule that usually opens by two valves, as seen in Fig. 309, *F*. This large family of over 1,800 species is almost universally distributed over the earth and of very common occurrence. They are among the most familiar weeds of waste places, fence rows and barren fields, and this unfavorable situation has perhaps resulted in the reduction in the size of the flowers which are usually small and of a pale-yellow or white color. They have the advantage of being grouped into conspicuous inflorescences, a tendency that will be noted again and again in higher orders. These flowers are of a more specialized type than in the open blossoms of the buttercups and poppies, and present a variety of interesting devices that can only be hinted at in this lesson. More commonly, perhaps, the stigmas are receptive as soon as the flower opens and the anthers are still closed, thus ensuring crossing. The short stamens are more frequently utilized only for crossing since they are below the stigma and so placed as to be in line with the nectar glands that are often located in sac-like enlargements of the perianth. The four long stamens are variously related to the stigmas. Sometimes they are below the stigma and when open are of service only for crossing, but later they elongate and bear some of the spores to the stigma. The reverse position of the sporophylls also occurs, but the outward opening of the anthers keeps the microspores from dropping by chance on the stigma until the stigma is finally lifted up to the anther by the elongation of the style, where it is sure to be dusted with spores, as the anthers open wider and wider. Certain species are characterized by an extraordinary twisting of the filaments, so that for a time the anther is turned away and bent over towards the perianth so as to be in line with the nectar glands, and later, at the close of the flowering period, a reversal of this bending brings the anther in contact with the stigma. These illustrations are sufficient to show that the flowers have many arrangements and exact movements that result in crossing, or, if this fails, in autogamy. It should be stated that several of these plants get along very well without crossing.

Many products of great commercial value are derived from various members of this order. Opium, from which morphine and other alkaloids are obtained, is the dried latex obtained from the incisions of the unripe capsules of the Turkish poppy. The majority of these plants contain acrid or peppery juices that render certain parts of the plant or their seeds of value as spices, oils, foods, etc. Many of them are biennial, forming a close rosette of leaves the first year and flowers and fruit the second. By cultivation an abnormal development of one part or another of the plant has been induced. This feature is well illustrated in the cabbage, where the elongated stem of the wild plant has become shortened by cultivation and covered with fleshy leaves, forming a large bud or head. Brussels sprouts are a modification of the cabbage in which the stem becomes more elongated and covered with numerous small heads. The great variety of kales are really headless cabbages in which the leaves remain free from one another, assuming a variety of forms. The cauliflower is a variety of the cabbage in which the inflorescence has been transformed into a fleshy mass of tissue, and in the kohlrabi the stem becomes swollen and herbaceous. The turnip is a related species of the cabbage genus in which the underground portion of the plant is modified. Radish, cress, horse-radish, caper, spices and oils of mustard, etc., are other products of the order, as well as many cultivated flowers, as the wallflower, stock, mignonette, etc.

144. Rosales, the Rose Order.—This enormous order, comprising over 14,000 species, is better known than any other, not only because of its great array of common field plants, but especially because of the large variety of our cultivated fruits and flowers that belong to it. The cultivated currant, blackberry, strawberry, apple, pear, cherry, peach, plum, pea, bean, hydrangea, syringa, rose, spiraea, wistaria, laburnum, as well as many native trees and shrubs, as the tulip tree, sycamore, witch-hazel, locusts, etc., belong to the rose order.

These plants may be looked upon as the most typical of the Choripetalae, just as the Liliales were the most representative of the monocotyledons. The parts of the flower are more com-

monly in five's and cyclic (Fig. 310, *A*) though the spiral arrangement still persists in nearly every family of the order. The most distinguishing feature of the group is seen in the adhesion of the receptacle to the calyx, thus lifting up the corolla and stamens about the ovary (perigynous flowers) and this growth also frequently results in an adhesion to the ovary (epigynous flowers). The simple forms of flowers that appear in many of the families are very suggestive of the Ranales, the sporophylls being free and often spirally arranged. Thus, in the live-for-

Fig. 310. Simple forms of the Rosales: *A*, flower of *Sedum*, showing the radial symmetry of the flower and five organs in each whorl. *B*, grass of Parnassus (*Parnassia*), a member of a closely-allied family. *C*, flower on first day of bloom, stamens converging over the pistil and encircled by row of modified stamens. *D*, section of flower, showing slight growth of base of receptacle and consequent adhesion to perianth. This flower is protandrous and the stamens in shedding the spores straighten up one by one on succeeding days and curve back towards the petals. The section shows three positions assumed by the stamens. You may calculate the significance of these features.

ever (*Sedum*), we have a flower that has always been cited as the typical flower of the angiosperms. It is very regular with five whorls of alternating parts of five members each, *i. e.*: 5 sepals, 5 petals, 5 to 10 stamens (one or two whorls) and usually 5 pistils (Fig. 310, *A*). This type closely resembles the arrangement of parts in the buttercups, save for the slight adhesion of the calyx and receptacle (Fig. 310, *D*). These fleshy plants as illustrated in the houseleek (Fig. 69), hen and chickens, etc., are very common forms of xerophytes. Owing to their ability to retain moisture (page 40) they can exist in the crevices of rocks and upon dry soil, while the formation of roots from a bit of stem or even a leaf, as well as their habit of producing buds on short stems that afterward become detached and grow into new plants, brings about a sure distribution and explains their popular name, live-forever.

(*a*) *Some Variations of the Rose Order.*—The active growth of the base of the receptacle and the consequent crowding result



FIG. 311. Higher forms of the Rosales: *A*, flower of the saxifrage. *B*, flower in section, showing the partial adhesion of receptacle to ovary. *C*, inflorescence of currant (*Ribes*). *D*, section of flower, showing receptacle forming the cavity for the ovules which is roofed over by the carpels, epigynous flower.

in an interesting series of departures from the simple type noted above that lead in very regular gradations to the highest forms of the order. In the higher family of the saxifrages, for example, which includes the saxifrage, false miterwort (*Tiarella*), heuchera, bishop's cap (*Mitella*), golden saxifrage (*Chrysosplenium*), the receptacle adheres to the ovary (Fig. 311, *A, B*) so that the flower is, in a measure, epigynous and the carpels are reduced

Fig. 312.



Fig. 313.

Fig. 312. A member of the rose family with the simple structure of the saxifrages: *A*, flower of the strawberry (*Fragaria*). *B*, section of flower, showing slight adhesion of receptacle to calyx. *C*, the fruit, akenes spirally arranged on the enlarged and fleshy receptacle.

Fig. 313. Higher forms of the rose family: *A*, flower of the rose. *B*, section of flower, showing the cup-like receptacle surrounding the pistils. *C*, flower of *Agrimonia*. *D*, section of flower, showing the bristle-covered receptacle completely surrounding the pistils.

to two and partly fused. Passing to the families of the hydrangeas, syringas (*Philadelphus*), currants (*Ribes*), etc., the receptacle and carpels are quite fused (Fig. 311, *C, D*) and the flower is strictly epigynous. In the rose family, which is very closely connected with the preceding group, this story of change in the evolution of the flower is repeated. In the simpler forms the receptacle adheres to the calyx, forming a cup-like

structure that may be rather broad and shallow as in the strawberry, cinquefoil and blackberry (Fig. 312), but in higher types the receptacle more or less completely surrounds the pistils as in the spiraea, avens, rose, agrimony (Fig. 313). In the apple family, including the apple, peach, quince, shadbush, thornapple, hawthorn, the ovules are completely enveloped by the receptacle (Fig. 314, *A–C*) as in the currants, while in the plum family, with its plum, prune, cherry, peach, almond, apricot members, the pistils are reduced to one and do not adhere to the cup-like receptacle (Fig. 314, *D–F*).

The senna family has essentially the same type of flower as the plum save that the pistil usually contains many seeds (nor-



FIG. 314. Flowers and fruits of the apple and plum families: *A*, inflorescence of the apple (*Malus*). *B*, section of flower, showing adhesion of receptacle to the ovary, epigynous flower. *C*, sections of the fruit—*c*, carpels of the pistil; *r*, fleshy receptacle. *D*, flower of cherry (*Prunus*). *E*, section of flower—*r*, cup-like receptacle which falls off as fruit matures. *F*, fruit in section, showing the outer part of the wall of the ovary as a fleshy rind and the inner part forming the stone or pit which enclosed a single seed.

mally but one develops in the plum) and splits at maturity into two valves, a form of fruit called a pod. For example, in the

honey locust (*Gleditsia*) and the Kentucky coffee bean (*Gymnocladus*), the receptacle forms a shallow cup which bears the regular sepals, petals and usually ten stamens about a single pistil (Fig. 315, *A*). The flowers are really monoecious, but in other respects are very suggestive of the plum flower. In the coffee bean tree, however, the petals are not quite equal and this irregularity becomes more noticeable in the sensitive pea (*Cassia*) (Fig. 315, *B–D*). In the redbud, or Judas tree (*Cercis*), the petals are very irregular, two of them being united into a boat-like structure and known as the keel which encloses the ten distinct stamens and single pistil, while two laterally placed petals, the wings, inclose the fifth petal known as the standard (Fig. 315, *E, F*). These three examples from the senna family make a



Fig. 315. Development of the irregular type of flower in the rose order: *A*, regular flower of the honey locust (*Gleditsia*). At left staminate flower. At right pistillate with single pistil, *c*, which develops into long flat pod, *s*, rudimentary stamens. *B*, flower of *Cassia*, showing slightly irregular corolla. *C*, section of flower—*s*, stamen; *c*, pistil. *D*, the fruit or pod. *E*, irregular flower of redbud (*Cercis*)—*c*, calyx; *k*, keel enclosing stamens and pistil; *w*, wings which arch over the standard, *s*. *F*, corolla removed, showing the ten stamens surrounding the simple pistil.

very easy transition from the regular flower of the plum to the highly modified flowers of the pea family.

(*b*) *The Pea Family, Papilionaceae.*—The pea family (Fig. 316) is the highest of the rose order and the largest family, with one exception, of all the angiosperms, comprising over 11,000 species. Here we find the same type of flower as in the redbud. The standard in this family, however, incloses the wings and the stamens, usually ten in number, may be distinct as noted above or united by their filaments into a sheath about the solitary pistil, more frequently a single stamen remains free, an arrangement called diadelphous (two brotherhoods). This type of flower is called papilionaceous from its fancied resemblance to a certain genus of butterfly, *Papilio.* The more important characteristics of this form of flower are very well illustrated



FIG. 316. Structure of the sweet pea (*Lathyrus*): *A*, flower of the pea —*c*, calyx; *s*, standard enclosing the two wings, *w*; *k*, keel. *B*, section of flower, showing the sporophylls concealed in the keel. *C*, one of the wings. *D*, perianth removed, showing relation of sporophylls.

in the pea (Fig. 316). The standard is the conspicuous and most highly-colored organ of the flower, overlapping the two wings which in turn practically cover the keel. By carefully removing the keel and wings, it will be seen that these organs are attached to the calyx and receptacle by rather narrow claws (Fig. 316, *C*) and that they are also locked together by a little process on each wing that fits into a groove on the keel. Nine of the filaments, the tenth being free, form a sheath about the ovary which terminates in a stylar brush of upwardly pointing hairs and in a stigma. The anthers and style are confined in

the tip of the keel (Fig. 316, *B, D*). The significance of these features will be discovered if you watch a bee probing into the sheath of filaments after the nectar secreted at the base of the ovary. The standard serves to attract and also direct his approach to the flower so that he alights at the proper place, *i. e.*, on the keel and wings. The weight of his body causes a slight depression of the keel and the stylar brush sweeps some of the microspores upon his body. It should be added that the flower is ready for the insect as soon as opened since the spores are discharged before the flower blooms. Only a portion of the spores are swept out by the insect and the dusting may be repeated to several visitors. The stigma does not mature until a later period and it is evident that there is a good chance of autogamy as soon as the stigma is mature. Some genera are always autogamous, but it has been observed in many cases that the spores grow more vigorously upon the stigma of another flower than upon the stigma of their own flower, so that if the stigma receives spores from its own and other flowers, the latter spores would develop better and effect fertilization. This is called the prepotency of foreign spores. In some species, autogamy is impossible owing to the failure of the spores to germinate except when transported to other flowers. There was a good illustration of this fact in Australia when the English clover was introduced and flourished exceedingly well, but produced no seed until bees from England were introduced for crossing. Plants transported to foreign countries are often sterile owing to the absence of the insect to which their variations are adapted. There are many variations of the mechanism shown in the pea. In some genera the style and stigma act like a piston, pushing out some of the spores from the apex of the keel with each visit of the insect and in some of these forms the wings are unlocked by the insect when the stamens that have been retained in the keel under considerable pressure are liberated by the dropping of the keel and spring up, scattering the spores with an explosive effect.

In many of these plants the pod splits at maturity into two valves, each of which twists with a sudden snap in opposite

directions, hurling the seeds to a considerable distance (Fig. 317, *A*). This reaction is due to the tension set up by the difference in the drying of the three strata of cells which compose the walls of the pod. The Papilionaceae contain very many nutritious food plants as the pea, bean and lentils, and also forage plants as the clover, alfalfa and vetches. In temperate regions the majority of the forms are herbaceous, only a few trees and shrubs as the locust, (*Robinia*), broom (*Cytissus*), dye weed (*Genista*), false indigo (*Amorpha*) belong to the family, but the tropics are richly represented by a great variety of woody forms that are valuable for lumber, resins and dyes, as the red sandalwood, licorice root, gum tragacanth, balsam of tolu, indigo, etc. Related to the pea family are the curious sensitive plant (*Mimosa*) and the acacias. One of the species of the *Acacia* develops enormous spines that are tunnelled by ants that protect the trees against leaf-destroying insects. Gum arabic is obtained from African and Australian species of *Acacia.*

One of the most interesting features of the Rosales is the variety of changes to which the receptacle and ovaries are subject in the ripening of the fruit. In the majority of cases, the pistil ripens as an akene or follicle (splitting along one side to free the seeds) or as a pod, without any considerable modification in the form of the ovary. In many instances, however, this growth is attended with pronounced alterations of the parts as in the currants and gooseberries where the receptacle adheres to or forms the ovary and the entire structures become succulent, forming a berry. The same organs in the witch-hazel develop into a horn-like capsule, the walls of which, after splitting open, contract and pinch out the hard, smooth seed with great force. In the raspberries, the ovaries are transformed into drupes which may be lifted off from the convex receptacle like a thimble, while in the blackberries, the drupes remain attached to the receptacle which becomes the most edible portion of the fruit. The attractive portion of the strawberry is the enlarged succulent receptacle while the objectionable hard particles on its surface are the akenes. The receptacle surrounds the pistils in the agrimony and rose, but in the former genus it is hard and covered with

hooked bristles for distribution of the nut-like fruit (Fig. 313), while in the rose it is fleshy and often brightly colored, serving in some cases as food for birds and thus effecting seed distribution. The most pronounced changes are seen in the apple and plum families. In the former group, the carpels become tough, forming the "core" (Fig. 314, *C*) and the edible part is the greatly enlarged receptacle. The plum, cherry, peach, etc., represent the ovary alone, the cup-like portion of the receptacle is cast off as the fruit matures and the ovary becomes a drupe (Fig. 314, *F*). In the pea family, the ovary usually becomes a pod with elastic valves but it is also variously modified. In the peanut, after the withering of the flower, the pistil is thrust into the ground where it develops as a pod that does not open at all. In other genera, the pod is nut-like as in the clovers, spirally coiled in alfalfa, or separable into nut-like joints that are provided with hooks as in the tick trefoil (*Meibomia*) (Fig. 317, *B*). Doubtless these methods of distributing their well-



Fig. 317. Fruits of the pea family: *A*, fruit or pod of the ground nut, showing the manner of seed dissemination by the snapping back of the valves with a twisting motion. *B*, fruit or lomentum of tick-trefoil (*Meibomia*). The lomentum breaks into as many nut-like parts as there are seeds in the fruit. *C*, hooked bristles on the surface of lomentum.

developed seeds, their symbiotic relations with the bacteria (page 57) and especially the elaborate mechanism of the irregular flowers have been the causes that have led to the abundance and

wide distribution of these plants. It will be noticed in the following orders as in the preceding Orchidales, that the irregular and more highly constructed flowers are generally represented by a great number of genera and, barring some weakness, as for example, the poorly developed embryos of the orchids and their peculiar habitats, by a great number of individuals in each species.

145. Sapindales, the Soapberry Order.—This group includes principally shrubs and trees, as the box, sumac (*Rhus*), smoke tree (*Cotinus*), holly (*Ilex*), burning bush (*Euonymus*), climbing bittersweet (*Celastrus*), maple (*Acer*), horse-chestnut and buckeye (*Aesculus*), etc. The flowers are more commonly small, of a yellow-green color and variously grouped into inflorescences (Fig. 318, *A*). We noticed in the rose order as the dominant characteristic the tendency towards the development of the basal region of the receptacle which was associated with the checking of its apical growth. In the Sapindales a similar shortening of the axis without its basal growth leads to quite a different series of variations. As a result of the crowding of the parts of the flower upon the shortened receptacle, the distinguishing features of the order are seen in the reduction of the number of the organs of the flower, a tendency towards the cyclic arrangement of parts and in the adhesion of the carpels. For example, the sepals and petals are four to five in number, usually distinct and the corolla may be entirely suppressed. There are two whorls of stamens, five to eight in number, rarely ten, while the pistils are reduced in number, ranging from two to more commonly three or rarely five. Thus we see that the spiral type of flower so often noticed in the rose and preceding orders has become reduced as a rule to the cyclic type and that five numerous sets of organs are not of common occurrence. This reduction of the flower and the suppression of parts is well illustrated in the maples (Fig. 318). The sepals cohere, forming a five-lobed calyx. The petals are often suppressed and a nectar disc is developed outside the stamens, a distinguishing feature of the order (Fig. 318, *C*). The two whorls of stamens are suppressed to a varying degree so that from four to eight

commonly appear and the pistils are normally reduced to two. More frequently the flowers are imperfect, either the pistils or stamens of each flower being aborted (Fig. 318, *B*). These two kinds of imperfect flowers are arranged on the same or different trees and they are adapted to small short-tongued lapping in-



FIG. 318. A common form of the Sapindales: *A*, inflorescence of sugar maple (*Acer saccharum*)—*s*, staminate flowers; *p*, pistillate flowers. *B*, staminate (right-hand) and pistillate flower enlarged. *C*, section of a pistillate flower, showing the two sterile stamens and nectar disc at base of filaments. *D*, section of ovary, showing early development of the wings of the fruit and the pendulous ovules with micropyle pointing down.

sects which visit these open types of flowers. The intelligent long-tongued bees and butterflies generally avoid such flowers, having learned by experience with colors and odors that a surer supply of food is to be found in those flowers that conceal their nectar and so exclude the promiscuous crowd of insects that swarm about the simpler types. It should be stated that the development of imperfect or incomplete flowers appearing in many of the orders is not to be looked upon as a primitive condition, though common in the lower monocotyledons and dicotyledons, since, throughout the Angiospermae, forms will constantly appear in which one or another set of organs fails to develop. The formation of anemophilous flowers, however, as in the box maple, is a return to a primitive condition.

The stimulation of fertilization results in a green wing-like outgrowth on each of the ovaries that assist at first in the manufacture of food for the embryo and later becomes a dry, membranous organ for seed distribution. This fruit, known as a schizocarp or samar, is at first partly loosened from its support and remains attached only by a small stalk (Fig. 319, *B*), which



FIG. 319. Fruit of the maple: *A*, mature fruit of red maple (*Acer rubrum*). *B*, schizocarp of Norway maple. The fruit split in half is still attached to the receptacle by delicate stalks.

requires a rather strong wind to snap it. Thus the fruit is freed under conditions that will result in the widest dissemination of the seed. The development of the chlorophyll-bearing tissue in immature fruits to assist in the work of food production is an economical arrangement of tissues often to be seen. It is noteworthy that these green fruits are often protected by bitter, acrid juices and poisonous properties that finally give place to attractive flavors, odors and colors, variations that are of considerable assistance to seed protection and distribution.

Many of the members of this order contain acid or poisonous juices, as in the scarlet fruit of the sumac, which were a source of acetic acid to the early settlers of this country, or poisonous oils, as in the poison ivy or poison oak (*Rhus radicans*) and the poison sumac (*R. Vernix*). The former species is a climbing

vine or sometimes a shrubby plant with leaves divided into three leaflets and with nut-like fruits (Fig. 320, *B*), while the poison sumac is an erect coarse shrub ten to fifteen feet high, with large pinnate leaves with reddish petioles and fruit clusters, as in the poison ivy (Fig. 320, *A*). As in the preceding orders, the lower

FIG. 320. Two poisonous species of the Sapindales: *A, Rhus vernix,* poison sumac. *B, Rhus radicans,* poison ivy.

members of this group are characterized by a regular alternation of the members of each whorl, but in the higher types the sets of organs vary in number and the flowers become irregular, as in the horse-chestnut and balsam weed or touch-me-not.

146. Orders Suggestive of the Sapindales.—The geranium order, Geraniales, shows essentially the same type and range of variation in the flower as the Sapindales, but there is this rather singular difference, namely, that the ovules, usually pendulous, are always turned away from the axis of the ovary with micropyle directed upwards, while in the Sapindales the opposite arrangement is to be found, micropyle pointing down (Figs. 318, *D*; 321, *D*). This order ranges from the regular flowers of the oxalis, flax and geranium families to irregular forms like the nasturtium, milkwort (*Polygala*), etc. Many of these plants are known by their peculiar juices, oils, gums, as in the spurges (*Euphorbia*), castor bean (*Ricinus*), citron, lemon, orange, etc. The fruit of the lemon and orange is a berry in which the outer

part of the ovary becomes leathery. The juicy pulp, which envelops the seeds, is formed from hair-like structures that arise on the inner side of the carpellary walls and by degrees entirely fill them. The navel orange is a chance variation in which a



Fig. 321. Flowers of the Geraniales and Rhamnales: *A,* inflorescence of *Geranium. B,* nearly mature fruit consisting of five united carpels. *C,* discharge of the seed. The carpels snap apart, owing to the tension set up by the drying out of the tissues. *D,* section of pistil, showing the ovules with micropyle directed upwards. *E,* flower of the grape (*Vitis*), a common form of the Rhamnales—*ca,* calyx, reduced to a rim; *c,* corolla, which opens at base and falls off as a cap; *n,* nectar glands within stamens. Compare Sapindales. *F,* corolla free from the receptacle. *G,* flower freed from corolla, ovary in section, showing micropyle pointing down as in Sapindales.

secondary receptacle, bearing compartments like the normal fruit, is introduced. The original stock from which the navel oranges grown in our country have been derived was obtained from Brazil.

The buckthorn order, Rhamnales, is very closely allied to the Sapindales and includes such familiar plants as the buckthorn

(*Rhamnus*), Jersey tea (*Ceanothus*), used in revolutionary times as a substitute for tea, the grape (*Vitis*), Japanese ivies, Virginia creeper (*Parthenocissus*), etc. The minute green or white flower shows a further reduction in parts, the petals not only being frequently suppressed, but the stamens are reduced to one whorl and placed opposite the sepals, whereas they alternate with the sepals in the Sapindales (Fig. 321, *E–G*), and the nectary is within the filaments.

147. Myrtales, the Myrtle Order.—This order marks a decided advance in the evolution of the flower over preceding groups, as indicated by the adhesion of the calyx and receptacle and usually of the ovary, the flowers being perigynous and usually epigynous. Another important character is that the flowers are always cyclic, the organs being regularly arranged in five whorls of four or less commonly of five members each, though the number of stamens may be greatly increased by the splitting of the original number of the set. The petals are frequently suppressed and the calyx is often highly colored, a variation to be seen in many groups when the corolla is wanting. The carpels, as a rule, contain numerous ovules (Figs. 322, 323). The Myrtales is an important tropical order and represented with us by many familiar plants, such as the meadow beauty (*Rhexia*), willow-herb (*Epilobium*), the large evening primrose family, with one of which, *Oenothera* (Fig. 323, *A, B*), deVries worked out his theory of mutation, and the common ditch and pond aquatics —the water milfoil (*Myriophyllum*) and mermaid weed (*Proserpinaca*).

The purple loosestrife (*Lythrum*) illustrates the characters of the lower members of the order before epigyny has become established (Fig. 322). The wand-like branches of this introduced plant with their terminal spikes of purple flowers are becoming rather common around the borders of marshes and water ways. The cup formed by the calyx and receptacle bears on its rim alternately with calyx lobes the rather twisted petals and at its base eight to twelve stamens. The pistil is composed of two carpels. This flower has become historic because of the attention that has been given to its devices for crossing. Fig. 322

shows that the sporophylls are of three lengths, short, medium and long, an arrangement called heterostyly. It can readily be seen that the three different lengths of the styles correspond exactly with the position of the anthers, consequently whatever



FIG. 322. Lower form of the Myrtales, flower perigynous: *A*, flower of *Lythrum*. *B*, flower in section—*c*, lobe of the tubular calyx, from the margin of which arise the petals, *p*. This flower has the long form of style and the short and medium form of stamens. *C*, flower with medium style and with short and long stamens. *D*, flower with short style and with medium and long stamens.

part of the proboscis or body of the bee or butterfly in visiting these flowers comes in contact with one of the sets of open anthers, exactly the same region of their bodies will touch the stigmas as soon as they visit another flower with a corresponding length of style. The spores from the three lengths of stamens differ in size and color, and it has been demonstrated that better results follow when crossing is effected between sporophylls of the same length, *i. e.*, long with long, short with short, etc. Heterostyly arose in many of the orders, as among some of the knotweeds, buttercups and rose families and in the gentians, primroses, forget-me-not, etc.

A higher and more characteristic type of the order is seen in the *Oenothera* and in the great willow-herb (*Chamaenerion*, Fig.

323). This latter plant flourishes in rather dry soils, forming large colonies by its underground stems and gaining in conspicuousness through its terminal racemes of large deep purple flowers. The parts of the flower are in fours. The receptacle adheres to the ovary and bears the linear segments of the calyx alternating with the round spreading petals. At the time of the opening of the flower the eight anthers are shedding their spores and in line with the nectaries, while the four-lobed stigma is closed and bent backward (Fig. 323, *C, s*). A day later the

Fig. 323. Higher forms of the Myrtales, flowers epigynous: *A,* flower of *Oenothera—o,* ovary. *B,* enlarged sectional view of flower, showing the receptacle enveloping the ovules. The sepals cohere, forming a tube, *t,* that bears at its summit the petals, *p,* and stamens; *c,* lobes of the calyx. *C,* inflorescence of the great willow herb (*Chamaenerion*)—*s,* closed stigma in young flowers; *o,* opened stigmas in older flowers; *a,* stigma touching anthers in withered flowers. *D,* capsule opening and discharging the seeds. *E,* a seed enlarged.—*C* after Kerner.

stigmas assume the position shown in the lower flowers, the lobes curving backward so as to lie in the pathway leading to the nectaries (Fig. 323, *C, o*). It is evident that these positions must necessitate a crossing if the flowers receive the proper vis-

itors. If, for any reason, crossing should fail autogamy results, owing to the continued curvature of the stigmas, which are finally brought into contact with the anthers, in which work the positions of the flowers and stamens coöperate, as shown in the lowest flower (Fig. 323, *C, a*). The microspores are fastened together in threads by a sticky substance, viscin, so that they adhere to the insect's body and are drawn out from the anthers in net-like skeins. This may be tested by applying the moistened finger to the opened anthers; this is also true of the related plants, as the fuchsia, evening primrose, fireweed (*Epilobium*), enchanter's nightshade (*Circaea*).

The numerous seeds developed in the capsule are provided at the end with tufts of long white hair (Fig. 323, *D, E*), which readily transport them and explains the quick appearance of these plants in forest lands that have been devastated by fire or cutting. In addition to the native plants mentioned above, many well-known tropical plants are represented in this order, as the mangrove of our southern swamps and the large myrtle family, often characterized by leathery leaves and aromatic oils. This family furnishes the clove, which is a flower bud, the pomegranate, guava, bay rum, the Brazil nut and the eucalyptus trees, some of which (Australian) attain the greatest height of any tree, over four hundred feet, with a diameter of twenty-four feet.

148. Umbellales, the Carrot Order.—This order marks the consummation in the reduction and adhesion of parts that we have seen steadily progressing through the various orders of the Choripetalae. Notice that the strictly epigynous, cyclic flowers (Fig. 324, *C, D*) are now reduced to four whorls, the five lobes of the calyx alternating with the five petals and these in turn with the four to five stamens. The reduction also appears in the pistil which usually consists of two carpels that contain but a single ovule each. The order contains two rather small families, the ginseng family (Araliaceae) represented by the sarsaparilla or spikenard (*Aralia*) and the ginseng (*Panax*) and the dogwood family (Cornaceae), including the pepperidge or sour gum trees (*Nyssa*) and dogwood or cornel (*Cornus*); and the large carrot family (Umbelliferae) of 2,100 species. The carrot family

contains many familiar native and cultivated plants which may be recognized by the hollow internodes of the stems, leaves variously lobed and attached by conspicuous sheathing petioles, peculiar odors derived from oils and resins, small flowers that are usually white or yellow and grouped into flat-topped inflorescences (umbels), which are usually surrounded by bracts, called the involucre (Fig. 324, *A, B*). These features are well seen in the wild carrot

FIG. 324. A common form of the Umbellales: *A*, stems of wild carrot (*Daucus*) with flowers arranged in compound umbels. *B*, inflorescence with all the umbels removed but one—*in*, bracts of the involucre. *C*, forms of flowers, the one on the right being a flower from the margin of the umbel. The corolla is enlarged and irregular, thus adding to the conspicuousness of the inflorescence. *D*, corolla partially removed to show the epigynous character of the flower and the cushion-like nectary at the base of the styles; *o*, ovary. *E*, fruit splitting into two nutlets. *F*, portion of hollow stem with sheathing leaf base.

which has become a troublesome weed in pasture lands and meadows. The calyx lobes are very small, a feature likely to be seen in any epigynous flower. The incurved petals alternate with the

five stamens and in the center of the flower, the two styles broaden out into a conspicuous nectary above the ovaries. The fruit is a schizocarp, splitting into two nut-like parts after the manner of the maple (Fig. 324, *E*). The flowers on the outside of the umbel become somewhat irregular, owing to the unequal enlargement of the outermost petals. These small flowers that individually cannot be seen at a distance of a few feet become the most conspicuous feature of our fields owing to their association in large numbers into compact, lace-like umbels. The aggregation of flowers has been noticed in various groups, as in the mustards, various families of the Rosales, in the Sapindales and buckthorns, of which latter simple regular type the carrot flower would appear to be a natural sequence (compare Figs. 321, *G*; 324, *D*). But in no group has so successful and varied an arrangement of the flowers been achieved. The advantages of this umbellate arrangement are apparent. The crowding of the flowers is attended with the reduction in their size and in saving of material while they have gained in the number of flowers and in conspicuousness. This arrangement also increases the chances of crossing since the insect in a single visit may cross a score of flowers in crawling over the umbel. Autogamy is prevented at first by the difference in the maturation of the anthers and stigmas, frequently one or more days intervening between these two conditions and in many genera the flowers are imperfect, the stamens and pistils being arranged on different parts of the umbel and so further assist in crossing. But even if the great variety of small lapping insects that swarm over these flowers were absent the construction of the umbel is such that the flowers are usually able to effect a crossing unassisted. The variety of devices for the accomplishment of this work are without parallel in any other group. The opening of the flowers of an umbel proceeds either from the circumference towards the center or from the center outwards. In some genera, as *Eryngium*, the flowers on the margin open first and the stigmas are ready for crossing while the anthers are closed and bent down upon the petals. On the following day, the inner adjoining set of flowers are in the same condition while the anthers of the

marginal flowers have been lifted up by their filaments so that they reach out and come in contact with the stigmas of the adjacent inner flowers, the curvature of the styles often assisting in bringing the two organs together. In this way, ample opportunity for crossing with the older flowers of another umbel is first given and later a crossing between the adjacent flowers of the same umbel is almost sure to result. In other cases, as the snakeroot (*Sanicula*) a few of the simple flowers are perfect while the remaining flowers of the umbel bear only stamens. The perfect flowers are the first to open, being in the same condition as the marginal flowers of the preceding example. Later the filaments straighten out, curve away from the stigmas and the anthers discharge their spores and finally drop off. Now the adjoining flowers open and extend their filaments so that the anthers are pushed over to the stigmas of the perfect flowers. In the chervil, a similar crossing is effected after the perfect flowers have shed their anthers by the numerous imperfect flowers growing above them and extending their anthers so that the microspores will drop straight down upon the stigmas (Fig. 325).

FIG. 325. Flowers of chervil: *A*, perfect flowers only in bloom. The anthers are discharging their spores before the stigmas are receptive. *B*, later stage—the staminate flowers are now in bloom and shedding their spores upon the mature stigmas of the perfect flowers.—After Kerner.

In the water parsnip (*Sium*), and beaked parsley (*Anthriscus*), etc., there are two kinds of umbels, one containing principally perfect flowers and the other only staminate flowers. The perfect flowers blossom first and shed their spores and anthers before the stigmas are receptive. After all the anthers have been shed, the stigmas mature and continue in this condition for two

days, during which time the umbels of imperfect flowers have grown above them and assume such a position that when the anthers finally open there results a rain of microspores upon the stigmas below.

The members of the carrot family are characterized by the presence of essential oils and resins in the roots, stems, leaves or especially in the fruits which usually contain oil cavities between the ribs of the ovary. These substances are the sources of the peculiar odors that make easy the recognition of these plants. These oils give the commercial value to several fruits, as in the caraway, parsley, fennel, anise, coriander. Several species contain very poisonous alkaloids, as the water hemlock (*Cicuta*), among the most poisonous of our native plants, and the hemlock (*Conium*) which perhaps furnished the potion drunk by Socrates. Several valuable food plants, as the carrot, parsnip, celery, parsley, are cultivated species of the family.

Series b. Sympetalae.

149. General Characteristics.—This group contains the most common and abundant of our flowering plants, approximating 42,000 species, and ranks as the most specialized of the angiosperms. The variations of these plants have been very successful and those are especially noteworthy that adapt them to the most intelligent and active of the insects. As a result of these advantages, they exceed all other groups in the number of individuals, though they comprise fewer orders as a consequence of the striking uniformity in the floral structure.

While a few of the simpler members of this group have distinct petals, the crowding of the organs on the receptacle has resulted, in nearly all the forms, in a cohesion of the petals into a sympetalous corolla, in the cyclic arrangement of the parts, and in the reduction in the number of stamens, so that each whorl does not exceed the number of petals, and more frequently one whorl of stamens is partly or entirely suppressed. The pistils are usually less numerous than the petals. The flowers range from regular hypogynous forms to epigynous and irregular

types through the same series of variations, as noted in the Choripetalae. The Sympetalae are the most recently evolved of the Angiospermae, as is apparent from their uniformity of structure and they appear to be adapted to temperate and northern conditions where they have preëmpted the open country and flourished exceedingly, forming the most characteristic features of the herbaceous flora. Aquatics are of rare occurrence and but few tree forms appear; notably the ash, persimmon, catalpa, paulownia, etc. Heath-like shrubs, however, are widely distributed in northern regions. The invasion of the tropics with its favorable conditions has led to an enormous increase in some of the groups and the development also of a great variety of woody plants, as trees and climbers.

150. Ericales, the Heath Order.—This order is the simplest of the Sympetalae and includes a great variety of plants that are largely northern in their distribution. Many are cultivated, as the azaleas, rhododendrons and laurels (*Kalmia*), and others are

FIG. 326. A simple form of the Ericales: *A*, the shin leaf (*Pyrola*) with flower just opened. *B*, flower with part of perianth removed to show the retention of the anthers by the petals. *C*, later stage of flowering. The anthers have been released by the spreading of the petals and the flower stalk has inclined so that the spores sprinkle down on the stigma. *D*, flower in section, showing the relation of parts in autogamy.—After Kerner.

familiar plants of bogs and woods as the sweet-pepper bush (*Clethra*), wintergreen (*Chimaphila*), Labrador tea (*Ledum*), Rhodora. *Leucothoë,* rosemary (*Andromeda*), leather leaf (*Chamaedaphne*), arbutus (*Epigaea*), checkerberry (*Gaultheria*), bearberry (*Arctostaphylos*), huckleberry (*Gaylussacia*), blueberry (*Vaccinium*), heather (*Calluna*), cranberry (*Oxycoccus*). The flowers of the heaths are very characteristic and sharply distinguished from other orders. More frequently they consist of five regular whorls of five members each, anthers usually opening by pores and often provided with two horn-like appendages, and the pistil is composed of five carpels, the ovary maturing as a capsule or berry (Fig. 326). The lack of adhesion between the whorls is an important feature of the order as contrasted with other sympetalae, each set being as a rule separately attached to the receptacle. While the sympetalous corolla is a step in advance of previous types, it is evident that these regular hypogynous flowers are in other respects of a rather simple character. This feature is further illustrated in some of the genera like *Clethra, Monotropa* and *Pyrola* where the free or slightly fused petals form a very natural transition from the Choripetalae (Figs. 327, *A*; 326).

The simpler forms of flowers are illustrated in *Pyrola* with its regular hypogynous and polypetalous flowers, though some species show a slight cohesion of the petals. The stigmas are mature with the opening of the flower while the anthers are bent back out of the way, their filaments being held under considerable tension by the petals (Fig. 326, *A, B*). A bee laden with spores would effect a crossing as soon as he alighted upon the flower, but in probing after the nectar he presses back the petals, thus releasing the anthers which snap down, spilling the spores upon him. Doubtless autogamy results in many of these forms if crossing fails owing to the expansion of the petals which would release the stamens, and the drooping of the flower (Fig. 326, *C, D*).

Other genera show varying degrees in the cohesion of the petals but in the majority of cases, tubular corollas appear (Fig. 327, *C*) as in the *Leucothoë,* bearberry, *Andromeda,* blue berries,

etc. The majority of such forms have horned anthers which are shaken by the insect and so assist in sifting out the spores. In the blueberry family, we note that an adhesion of the receptacle and ovary has begun (Fig. 327, *D*) while in *Rhodora,* rhododendron and azaleas another line of variation appears in the irregular corolla (Fig. 327, *F*). In the two latter genera, the styles and stamens extend far out, furnishing a natural landing place for hovering animals as moths and humming birds.



Fig. 327. Common forms of Ericales: *A,* Indian pipe (*Monotropa*). Leaves reduced to scales and without chlorophyll, owing to saprophytic habit of plant. *B,* section of flowers, showing polypetalous corolla. *C,* inflorescence of the blue berry (*Vaccinium*). *D,* section of flower, showing sympetalous corolla and epigynous type of flower. *E,* a stamen enlarged, opening by pores at end of anther. *F,* azalea, showing irregular type of flower. *G,* spores attached by viscid threads.

In this position the spores cannot be sifted out and we find them fastened together in fours, just as they were formed in the mother cell, by sticky threads which cause them to adhere to

the insect's body in fringe-like masses (Fig. 327, *G*). One of the lower basidiomycetes (*Exobasidium*) infests the leaves and flowers of the azalea and other genera, causing large watery outgrowths that are often eaten under the impression that they are the fruit of the plant. The members of this order are adapted to a wide range of conditions though more characteristic of north temperate and arctic regions where they constitute the conspicuous features of the tundras, bogs, heaths and moors. These plants are often characterized by their thick leathery, evergreen leaves of pronounced xerophytic character and their common distribution in wet moors and bogs presents a most puzzling problem in the association of plants. Many of the heaths have been cultivated from very early times and they have lent themselves so readily to crossing that perhaps no order furnishes so many forms for decorative and landscape effects. Several yield valuable fruits, as the blueberries, huckleberries and cranberries

151. Orders of a Higher Type than the Ericales.—The primrose order (Primulales) forms a very natural transition from the Ericales. As the next step in advance, we note the slight adhesion of the stamens to the corolla (Fig. 328, *A*) and their frequent reduction in number. The occurrence of staminodia, page 388, in certain forms marks the first transition to the reduction of the number of stamens. The pistil is generally composed of five carpels forming a capsule without partitions (unilocular) and containing numerous centrally placed seeds. This order includes the loosestrife (*Lysimachia* and *Steironema*), star flower (*Trientalis*), shooting star (*Dodecatheon*) and cultivated forms of the cyclamen and primroses.

Proceeding to the gentian order (Gentianales), it will be noted that the flowers are variable and not as clearly characterized, the corolla being sometimes wanting or polypetalous. The reduction of the stamens to a single whorl and the pistils usually to two in number indicate points of advance that are to be associated with the twisting of the petals in the bud and the opposite arrangement of the leaves as distinguishing features of the order (Fig. 328, *B–D*). Note should be made of the fact

that the two pistils often do not cohere, doubtless a survival of a more primitive condition. The Gentianales include the ash (*Fraxinus*), lilac (*Syringa*), *Forsythia,* olive (*Olea*), valued for fruit and oil, fringe tree (*Chionanthus*), privet (*Ligustrum*), jasmine, *Strychnos,* yielding the alkaloid strychnine, oleander (*Nerium*),



Fig. 328. Primulales and Gentianales: *A,* flower of loosestrife (*Steironema*). Above flower in section, showing cohesion of single row of stamens to base of corolla and the numerous ovules on a free central placenta. These are the most important characteristics of the order. *B,* fringed gentian (*Gentiana*), showing leaf arrangement and twisting of petals in bud, *b,* characteristics of the order. *C,* section of flower. *D,* fruit, the two carpels separated.

the gentians, marsh pink (*Sabbatia*), the aquatic floating heart (*Limnanthemum*) and the highly specialized plants of the dogbane and milkweed families. The latter family is distinguished by the microspores being united into club-shaped pollinia that

are attached, one from each of the two adjoining anthers, to a peculiar hook, so that the leg of the insect becomes caught in the hook and withdraws the pollinia in pairs.

152. Polemoniales, or Phlox Order.—This group is the richest in the number of species, over 14,600, of any order of angiosperms. The advance over previous orders is seen in the more complete adhesion of corolla and stamens, the filaments appearing to arise at a higher point on the corolla (Fig. 330, *D*) and

Fig. 329.



Fig. 330.

Fig. 329. Flower of the morning-glory (*Ipomoea*), showing the tubular corolla characteristic of the Polemoniales.

Fig. 330. Boraginaceae: *C*, inflorescence of comfrey (*Symphytum*). Note the coiled inflorescence, *a*, a feature of this family. *D*, section of flower, showing the deeply four-lobed ovary and the stamens cohering high on the corolla and alternating with small tongue-like scales.

the stamens are reduced to a single whorl, and frequently less than five in number. The pistils cohere completely and are usually composed of but two carpels. This reduction is associated with a pronounced irregularity of the corolla in the higher families and a high degree of specialization in the construction of the flower which accounts for the occurrence of the large number of individuals. The order is noted for its great number

of showy flowers and the large tubular corollas which attain their highest perfection in several of the families.

(*a*) *The More Important Families of the Phlox Order.*—The flowers of the lower families are regular, as in the morning-glory, sweet potato, dodder (*Cuscuta*), a yellow thread-like parasite that twines about various plants, and the phlox (Fig. 329). The flowers of the rough-leaved borages (Boraginaceae) are also regular but they are distinguished by having the ovary of the two carpels deeply lobed so that the fruit appears as four nutlets and also by their coiled inflorescences as shown in Fig. 330. This family includes the heliotrope, hound's-tongue (*Cynoglossum*), forget-me-not (*Myosotis*), comfrey (*Symphytum*), and the blueweed (*Echium*) in which the corolla becomes irregular.

The mint family, Labiatae, is world-wide in its distribution and the largest of the order with 3,000 species. These plants are sharply characterized by square stems, opposite leaves, aromatic oils, nut-like fruits as in the borages, but the corolla is very irregular and usually two-lipped. The stamens, usually two short and two long ones, together with the two-lobed style are more frequently concealed under the upper lip of the corolla (Fig. 331). The flowers are generally so placed that the lower lip serves as a landing place for insects and in entering the flower, the top of their bodies rubs against the stigmas or anthers. Crossing is secured by the difference in the time of the maturation of the anthers and stigmas, assisted by a variety of movements, such as the alternate curving down of the anthers and stigmas which brings first one set of organs and then another into the pathway of the insect. One form of this arrangement is seen in the sage, where there are but two strangely modified stamens, the two anther lobes being connected by a long-curved rod which is fastened near one end to a short filament so that the anther becomes a lever with the filament as a fulcrum (Fig. 332). Only the anther lobes concealed under the upper lip are fertile. As the bee enters the flower, he pushes up the short arm of the lever, thus causing the long arm to swing down, pressing the fertile anther upon his back and dusting him with



FIG. 331. FIG. 332.

FIG. 331. A common species of the mint family: *A,* inflorescence of the skull cap (*Scutellaria*). Note the square stem, opposite leaves. Why are all the flowers facing one way? *B,* flower enlarged, showing the two-lobed under lip and the three-lobed upper lip which conceals the sporophylls. *C,* section of the flower. Ovary four-lobed, stamens cohering with the corolla and anthers concealed with the stigma beneath upper lip. Purpose of the crest, *c,* on the calyx?

FIG. 332. Flower of the sage (*Salvia*): *A,* flower after the anthers have shed their spores. The two-lobed stigma is bending down into the position occupied by them. *B,* sectional view of the flower, showing four-lobed ovary with nectar glands at the base, stigma not receptive and bent back. As the insect enters the flower he pushes against the sterile lobe of the anther, *t,* and thus causes the fertile lobe, *a,* to swing down upon his back. *f,* filament of anther.

spores. Later, the style which is at first immature and above this apparatus, grows out and curves down and at maturity comes to lie in such a position as to rub upon the back of a visiting insect at the same point as the anthers. The peculiar forms of the corolla, the relation of the anthers and stigmas and the frequent occurrence of hairs and bristles that direct the insect with the greatest precision to particular parts of the flower with a view to crossing, mark this family of the mints as the most specialized of the order. A fine series of devices are seen in such common mints as the blue curls (*Trichostema*), skullcap (*Scutellaria*), hyssop (*Agastache*), ground ivy (*Glecoma*), catmint (*Nepeta*), selfheal (*Brunella*), motherwort (*Leonurus*), dead nettle (*Lamium*), hedge nettle (*Stachys*), sage (*Salvia*), bergamot (*Monarda*), pennyroyal (*Hedeoma*), mountain mint (*Koellia*), bugle weed (*Lycopus*), etc. Many of these plants are of commercial importance. *Mentha* yields valuable oils as spearmint, peppermint, and menthol and from other genera are derived oils used in perfumery, medicine and as condiments, as rosemary, lavender, origanum, thyme. Several are cultivated for their flowers and foliage, as the coleus, monarda, stachys, etc.

The large family of figworts, Scrophulariaceae, with its 2,400 species, closely resembles the mints but is distinguished by the ovary being undivided and containing many seeds on a central axis (Fig. 333, *C*). While the simpler forms have almost regular corollas and five stamens, the higher forms are bilabiate with two to four stamens as in the mints. This family contains some of the most showy of our cultivated plants, as the foxglove (*Dasystoma* and *Digitalis*), *Gerardia,* snapdragon (*Antirrhinum*), rattlebox (*Rhinanthus*), toadflax (*Linaria*), *Paulownia,* beardtongue (*Pentstemon*), monkey flower (*Mimulus*); and the allied *Catalpa* and trumpet creeper; also many other attractive plants that are not cultivated as the mullen (*Verbascum*), figwort (*Scrophularia*), turtle head (*Chelone*), hedge hyssop (*Gratiola*), speedwell (*Veronica*), painted cup (*Castileja*), lousewort (*Pedicularis*), cow-wheat (*Melampyrum*). These plants are among the most characteristic features of our flora, the family being largely confined in its distribution to the north temperate

regions. The flower of the toadflax (*Linaria*) illustrates a common type of the figwort family (Fig. 333). Such flowers are said to be personate, that is bilabiate but with the under lip arched so as to meet the upper lip and entirely closing the mouth of the corolla. This arrangement very effectively protects the microspores and conceals the nectar and it requires a rather muscular insect alighting upon the knobbed lower lip to force his way into the flower. It is worth any one's time to sit by



FIG. 333. Examples of the figwort family: *A,* inflorescence of the toad flax (*Linaria*). *B,* flower viewed from beneath, showing the under lip arching up against the upper, two-lobed lip—*s,* nectar spur of the corolla. *C,* section of the flower viewed from the side, showing the undivided ovary with central ovules. Note the stigma and anthers concealed at the lips of the corolla. *D,* sectional view of flower of *Rhinanthus* with the four stamens arranged in pairs. *E,* appearance of the stamens as viewed from the mouth of the corolla. The filaments have been pressed apart, thus separating the opened anthers.

this plant and examine the mechanism of the flower while the bee is at work. The sporophylls present the same variety of arrangements for crossing as noted in the mints. The nectar glands are situated at the base of the carpels but the nectar does not remain upon the glands as in the majority of flowers, but

runs down through a narrow duct between the filaments and collects in the spur-like prolongation of the corolla (Fig. 333, *s*).

The yellow rattlebox (*Rhinanthus,* Fig. 333, *D, E*) shows another form of the flower, common in this family. The general arrangement of the floral organs is the same as in *Linaria,* but the anthers are in pairs so that when they open, owing to the rigidity of the filaments or the pressure of the upper lip, they are kept tightly pressed together like a pair of sugar tongs, thus preventing the shedding of the spores. Owing to the form of the corolla and often because of numerous hairs and bristles, the insect is directed in such a definite way to the flower that he comes in contact with appendages of the anthers, or causes a slight deflection of the rigid filaments or of the lip of the corolla, any of which movements are sufficient to separate the anthers and bring down upon him a shower of spores. These flowers are largely protogynous so that at first only a crossing is possible. In many species, autogamy results from the downward bending of the corolla and style, accompanied by a loosening of the anthers so that during the last days of flowering the spores may fall directly upon the stigma.

Many of these flowers are characterized by mottlings and blotches. It appears to be a rather general law that the most highly modified parts are variegated in this manner—compare the mints and orchids. Less specialized flowers present an association of various colors or veinings, a feature that is illustrated in many families as violets, peas, geraniums, etc., whereas, the simplest types of flowers are usually of a uniform color. The figworts have been called a suspicious group, and if not actually poisonous, none at least serve as food. Several are medicinal, as *Digitalis, Veronica, Gratiola,* etc. Many are parasitic, as the foxglove, *Gerardia,* eyebright (*Euphrasia*) and the related and reduced brownish rapes (*Thalesia* and *Orobanche*), beech drops (*Leptamnium*) and squawroot (*Conopholis*).

The curious family of the bladderworts is closely related to the figworts. The bladderwort (*Utricularia*) is a very common aquatic in still waters and ponds. Certain leaves become modified into elaborate sacs with trapdoors for enticing and capturing

small insects which die and decay in these prisons and are ultimately absorbed and so contribute to the nourishment of the plant. The butterwort (*Pinguicula*) is another member of this family found in the northern countries and also in our southern states. It has more the appearance of a violet with a rosette of leaves which are provided with glands that secrete a viscid substance for the capture of insects and also digestive fluids. The contact of the insect acts as a stimulus, causing the leaves to roll up and so bringing to bear upon him a larger surface of digestive glands. One of the species of *Pinguicula* is used in northern countries to curdle milk in place of "rennet" which contains the similar digestive fluids of calves' stomachs.

The potato family (Solanaceae) shows many of the irregularities of the figworts, but the majority of its 1,700 species have regular flowers as illustrated in Fig. 334. While the family

FIG. 334. Flower of the potato (*Solanum tuberosum*), showing the stamens inserted on the tube of the corolla and encircling the style.

is largely tropical, many of the species are cultivated, as the potato, eggplant, tomato, cayenne pepper (*Capsicum*). Poisonous, acrid and narcotic properties are characteristic features of the family. Belladonna and atropine from *Atropa,* stramonium from *Datura* and nicotine from *Nicotiana tabacum* are characteristic drugs. Ground cherry (*Physalis*), nightshade (*Solanum*), *Petunia,* etc., are cultivated forms.

153. A Transitional Order.—It would appear that all the changes possible in the hypogynous type of flowers had been wrought in the members of the Polemoniales and that the next step in advance must be to the epigynous flower. This is seen to be the case with the madder order, Rubiales. Here the

flowers are epigynous and the parts are generally in fives though the carpels vary greatly in number. The simplicity in the structure of the flower and often also the form of the inflorescence, as in the elderberry (*Sambucus*), bedstraw (*Galium*) and arrow-

FIG. 335. Simpler forms of the Rubiales: *A*, inflorescence of the arrow-wood (*Viburnum*). *B*, simple, epigynous flower of the elderberry (*Sambucus*)—*o*, ovary. *C*, the twin flower (*Linnaea*).

wood (*Viburnum*) is strikingly suggestive of the Umbellales (Figs. 335, *A*, *B*; 324). It appears that each step in advance is attained in a very simple way and that only gradually are alterations made that lead up to the specialized types. Look back over the preceding groups and orders and note that many of the irregular types of flowers are preceded by a long series of regular forms. So in the madder order, this new departure of the flower is attended with such a simplicity of structure as to render difficult the separation of some of the genera from the open flowers of the Umbellales which have also arrived at the same stage of development, page 432. This would not necessarily imply a relationship, since it repeatedly happens both among plants and animals that identically similar structures arise in groups in no way related. From these simple flowers that may

be no more sympetalous than certain genera of the carrot order, we pass to more pronounced tubular forms in the bluets (*Houstonia*), buttonbush (*Cephalanthus*), twin flower (*Linnaea,* Fig. 335, *C*), snowberry (*Symphoricarpus*) and finally to irregular and even labiate types as in the honeysuckle (*Lonicera,* Fig. 268), valerian, teasel (*Dipsacus*) and scabious (Fig. 336). It is noteworthy that in the three latter genera the flowers become massed in a dense inflorescence, known as a head, which are subtended by modified leaves, the involucre (Fig. 336) and the pistils are reduced to a single fertile carpel with two-lobed style. These



Fig. 336. Advanced forms of the Rubiales: *A,* inflorescence of *Scabiosa,* at the right showing the involucre, *in. B,* a single flower enlarged, showing the somewhat irregular corolla. At the right the sectional view of the flower shows the calyx, *ca,* terminating in bristle-like teeth; *br,* bract-like cup surrounding the calyx. *C,* irregular flower of valerian—*n,* nectar sac; *ca,* rudimentary calyx, which matures in the fruit as a mass of delicate feathery organs, known as the pappus.

variations will become very prominent in the next order. The Rubiales are an important tropical group and furnish the coffee (*Coffea*) and the cinchonas which yield such drugs as quinine and calisaya.

154. Campanulales, the Bellflower Order.—This order marks the culmination of the tendencies that we have seen steadily progressing through the monocotyledons and the dicotyledons. The variations have been of so peculiar and successful a nature that no group of plants are so widely distributed and in the more specialized families of so common occurrence. Over 14,500 species are known. The parts of the epigynous flowers are arranged in four whorls of usually five members each. The anthers are aggregated and usually cohere, forming a sheath



FIG. 337. Lower forms of the Campanulales: *A*, inflorescence of the bellflower (*Campanula*). *B*, section of a young flower. The hairy style is pushing up between the encircling anthers and sweeping the spores out of them. Note the closed stigmatic lobes. *C*, older flower. The anthers are withering and the stigmas are curving back towards the spore-covered style. *D*, flower of *Lobelia*. The tubular corolla is opened at one side, permitting the style and encircling stamens, *g*, to protrude. *E*, section of flower, showing the anthers, *a*, united about the style and stigma. *F*, relation of stigma to anthers. At right the section shows the anthers cohering about the bushy style, which acts later as a brush sweeping out the spores. At the left the style has grown beyond the anthers and the stigmatic lobes are spreading apart.

about the style which is frequently covered with hairs and acts like a piston rod in the cylinder of anthers. The flowers are protandrous with few exceptions, the anthers opening on their inner sides and discharging the spores upon the style which later sweeps them out as it elongates. The pistils are generally reduced to a single one-ovuled carpel that ripens as an akene. Leaving out of consideration the gourds which include the melons, pumpkin, cucumber and gourd, as a family of uncertain alliance, we find a very natural sequence leading to the higher families. For example the lower members of the bell-flower family have nearly or quite regular flowers, parts in five, anthers grouped about the style but not united, fruit a capsule with many seeds as in the harebell (*Campanula*) and *Specularia* (Fig. 337, *A, C*). In the *Lobelia* the corolla usually becomes irregular and slit down one side, thus approaching the form assumed in the highest groups of the order. The anthers are united about the style which is two lobed and provided with a whorl of hairs to sweep the spores from the cylinder of anthers. The fruit is a capsule of two carpels (Fig. 337, *D–E*). This brings us to that great group of familiar plants that were formerly known as the Compositae, but that are now separated into the chicory family (Cichoriaceae) of 1,400 species, the ragweed family (Ambrosiaceae) of 55 species, and the thistle family (Cardu-aceae) of over 11,000 species. The most conspicuous feature of these three families is the aggregation of numerous small flowers in heads subtended by one or more rows of bracts that form a calyx-like involucre (Fig. 338, *in*). This type of inflorescence might readily be mistaken for a single flower as the buttercup, rose, etc. This tendency to group the flowers in heads and compact clusters has been attained in several orders, notably the mustards, peas, Umbelliferae, mints, scrophularias, and especially in the teasel family, page 450. But in no group has the aggregation been so successful and coupled with such efficient types of flowers. Leaving out of consideration the degenerate ragweeds the individual flowers of a head are very uniform in structure. They are epigynous, parts usually in fives, calyx wanting or more often appearing as tufts of hairs, plumose

or barbed bristles, and known as the pappus (Figs. 339, 341). The corolla is tubular or partially split open, forming a strap-shaped blade known as the ligulate corolla (Fig. 339, *F*). The anthers are united about the style which is often hairy and pushes

FIG. 338. Habit of the dandelion (*Taraxacum*), an example of the Cichoraceae: *a*, appearance of the inflorescence or head; *b*, appearance of the head during the ripening of the seed; *in*, involucre; *c*, appearance of the fruit.

out the microspores like a piston (Fig. 341, *B–D*). The style is two lobed, the stigma usually appearing as a line on the inner surface of the lobes and the single, one-ovuled carpel matures as an akene. Scale-like bracts, the chaff, are often associated with the flowers (Fig. 341, *B*). The nectar formed from the glands at the base of the style is concealed at the bottom of the corolla tube.

(*a*) *The Chicory Family.*—This family is distinguished by all the flowers of the head being ligulate (Fig. 339) and mostly yellow in color and by the possession of milky, bitter or acrid juices.

Here belong the goat's beard (*Adopogon*), hawkbit (*Leontodon*), sow thistle (*Sonchus*), hawkweed (*Hieracium*), rattlesnake root (*Nabalus*), as well as several introduced and native plants that are cultivated, as the dandelion (*Taraxacum*), species of chicory (*Cichorium*), salsify (*Tragopogon*), lettuce (*Lactuca*). The character of the head, flower and fruit of this family are well shown in the dandelion (Fig. 339). All the flowers are ligulate,



FIG. 339. Flowers and fruit of *Taraxacum: A,* sectional view of inflorescence—*in,* involucre. The flowers in the center of the head not as yet in bloom. *B,* an unopened flower. The thread-like calyx (pappus) and corolla arising from the ovary, *o. C,* corolla opening. *D,* later stage, the style has elongated, sweeping the spores from the sheath of anthers, *an,* and the two stigmatic lobes are beginning to open. *E,* flower in full bloom, the stigma lobes recurving. *F,* mature fruit. The pappus is lifted up on a long, slender outgrowth of the ovary, *o.*

the five lobes frequently to be seen on the margin of the corolla indicating the number of petals, and the calyx assumes the form of minute hairs (Fig. 339, *B–E*). The five filaments adhere to the tube of the corolla while the anthers form a cylinder about the style and discharge their spores before the flower opens. The style begins to elongate as soon as the flower opens and pushes

out the spores which may be seen as little piles of dust at the top of the anthers in freshly opened flowers. The filaments of this genus are sensitive to touch and curve outward, pulling down the sheath of anthers, thus assisting in sweeping out the spores and exposing the style when an insect irritates them in his quest for nectar. This device appears in other genera and is sometimes the only means of exposing the microspores and styles. Autogamy is at first impossible owing to the fact that the two lobes of the style are closely pressed together effectually excluding the microspores from its inner, stigmatic, surface. Later the style grows considerably above the anthers and the lobes separate, exposing the stigmas for crossing (Fig. 339, *E*). The association of this type of flower is more effective than any as yet noticed, and it would appear impossible for any of the numerous insects which frequent these flowers to miss crossing a score of them at a single visit. Crossing between the flowers of a head is effected in some genera by the movements of the flowers. The inner leaves of the involucre fold over the flowers at night and in rainy weather, protecting them like a perianth. This results in crowding of the flowers together, and the stigmas are often brought in contact with the spores that have been dusted upon the various parts of adjoining flowers. The outward curving of the stylar lobes may also bring the stigmas in contact with the spore-covered parts of adjoining flowers with a like result. Autogamy is brought about in the dandelion, as in nearly all other members of the order, by the curvature of the lobes of the style which continue to bend back until the stigmatic surface is brought in contact with the spore-covered style (Fig. 341, *D*). The blooming of the flowers progresses from the margin to the center of the head so that during several days new sets of flowers are exposed for crossing. When the period of bloom is passed the involucre closes over the head and remains in this condition until the fruit is mature (Fig. 338, *b*). During this period, the stalk bearing the head elongates considerably, so as to lift the fruit above the surrounding vegetation. The corolla withers away and the hair-like calyx grows out into a white, delicate pappus, which is lifted up on a beak-like outgrowth of the ovary (Fig.

339, *F*). The fruit is now mature and in a position for distribution. The involucre opens, the hairs of the pappus expand, loosening the akenes so that the least touch or breath of air floats off the fruit as a most perfect type of parachute (Fig. 338, *c*).

(*b*) *The Thistle Family, Carduaceae.*—This is the largest

FIG. 340. Inflorescence of the bur-marigold (*Bidens*), a common ray form of flower of the thistle family: *in*, involucre.

family of the angiosperms. The flowers have the same arrangement and structure as in the Cichoriaceae, save that the corollas are either all tubular, as in the thistle and ironweed (Fig. 342), or the marginal flowers of the head may be ligulate, thus increasing the conspicuousness of the inflorescence, as in the asters, daisy, etc. (Figs. 340, 341). The marginal ligulate flowers are termed ray flowers to distinguish them from the inner tubular or disc flowers. The ray flowers may be sterile, though more frequently imperfect and provided with pistils. The calyx may be wanting, but more often assumes the form of silky or plumose pappus or of membranous scales for wind transportation or of barbed bristles of various kinds for distribution by animals. In the case of the burdock (*Arctium*) the involucre is covered with hooks. The Carduaceae includes a great array of plants, many of which are among our troublesome weeds: Ironweed (*Vernonia*), thoroughwort (*Eupatorium*), blazing star (*Lacinaria*),

golden-rod (*Solidago*), aster, fleabane (*Erigeron*), cat's-paw (*Antennaria*), everlasting (*Gnaphalium*), rosinweed and compass plant (*Silphium*), Spanish needles (*Bidens*), daisy (*Chrysanthemum*), groundsel (*Senecio*), thistle (*Carduus*). Culti-

FIG. 341. Flowers and fruit of *Bidens: A,* sectional view of the inflorescence—*r,* ray flowers; *d,* disc flowers; *in,* bracts of the involucre. *B,* disc flower before opening, the shaded region showing the position of the anthers in the corolla—*c,* calyx or pappus in the form of downwardly barbed bristles; *b,* bract or chaff associated with the disc flowers. *C,* early stage in the opening of the flower. The stamens have been lifted beyond the mouth of the opened corolla by the growth of their filaments and the style is elongating, pushing out the spores, which appear in little piles at the top of the anthers. *D,* last stage in the bloom of the flowers. The style has grown beyond the anthers and the stigmatic lobes have reflexed, touching the spore-covered style. *E,* a sterile ray flower, much less enlarged. *F,* the fruit, showing the barbed pappus for dissemination.

vated for their showy flowers: Tickseed (*Coreopsis*), cone flower (*Rudbeckia*), sunflower (*Helianthus*), cornflower (*Centaurea*), cinerarias, etc. Medicinal: Wormwood (*Artemisia*), one species of which yields absinth, milfoil (*Achillea*), colt's foot (*Tussi-*



FIG. 342. A common tubular form of flower of the thistle family: *A*, inflorescence of the ironweed (*Vernonia*). *B*, sectional view of the inflorescence, only the outer flowers in bloom. *C*, enlarged view of one of the flowers—*c*, calyx or pappus. *D*, fruit.

lago), tansy (*Tanacetum*), chamomile (*Anthemis*), arnica, burdock (*Arctium*).

The ragweed family, Ambrosiaceae, is a small group, principally North American in its distribution, that have become separated from the Carduaceae through degeneration. The heads contain a few greatly reduced anemophilous flowers that are always imperfect and generally lacking in calyx or corolla, or both, and characterized by free anthers. The more familiar are: The marsh elder (*Iva*), ragweed (*Ambrosia*), and clot-bur and cockle bur (*Xanthium*).

These three alliances are apparently the most recently evolved of the angiosperms, and owing to their numerous variations that have been so successful in meeting the present conditions upon the earth, they have become the dominant plants the world over. Their success can be traced to a variety of causes, as the occurrence in many genera of perennial underground stems, which make them to a degree independent of climate conditions and the depredations of grazing animals. A great many are avoided

because of their protective armor of prickles, bitter, acrid juices or oils and resins. Especially to be noted is the seed-like fruit with its numerous devices for distribution. All these features, among others, explains their extensive distribution in open countries. Without doubt the greatest advantage appears in the structure of the flower and the inflorescence. The highest type of a flower, as in the case of a machine, is the one that is most efficient. The flower is not so elaborate as the orchid or as those of some other groups, but it is superior because it accomplishes its work with greater certainty and more economically. Reduction has been made in every part without loss of conspicuousness and variations have also appeared that practically eliminate the uncertainties of crossing and autogamy, and consequently the formation and distribution of the seed is ensured. In a word, every evolutionary tendency that you have noticed in the development of the flower finds its expression in this group of plants.

INDEX

Absorption, 12, 20, 45
Accompanying cells, **65, 66,** 67
Achlya, **198**
Actinomorphic, 353
Adiantum, **298**
Aecial stage, **222**
Aecidiospores, **222**
Agaricaceae, **230,** 233
Agaricales, 227, **228**
Agrimony, **417**
Akene, **347,** 365, **406**
Albugo, **200,** 201
Algae, 154
  blue green, 145
  brown, 181
  conjugating green, 165
  filamentous green, 170
  green, 155
  red, 189
  tubular green, 174
Alga-like fungi, 197
Allen, 194
Alternation of generation, 123, 181 290
Amanita, **230**
Ambrosiaceae, 458
Ament, **397,** 398
Anemophilous, 355, 371
Angiospermae, 345
Ankistrodesmus, **162**
Annular rings, 77, **78**
Annuals, 60
Annulus, **230,** 272, **278, 295**
Anther, **102, 103**
Antheridial cell, 107
Antheridium, antheridia, 176
Antherozoids, 176, 248
Anthoceros, **264, 266**
Anthocerotales, 266
Antipodal cells, 105, **106**
Antitoxin, 145
Apophysis, 277, **278**
Apple, **418**
Arales, 377
Archangiopteris, **286**
Archegonium, 248, **249**
Arcyria, **135**
Aroid order, 377
Arrow-wood, **449**
Ascocarp, 206, **211**
Ascomycetes, 205
Ascospores, 206
Ascus, asci, 206
Asexual generation, 122, 173, 180, 193, **254,** 290
Aspergillales, 209
Aspergillus, **210, 211**
Autogamy, 350

Bacillus, **140**
Bacteria, **139, 140**
  of decay, 142
  of disease, 143
  nitrifying, 57, 142
Bark, 72
Basidial stage, 224
Basidiomycetes, 221
Basidiospore, **224**
Basidium, basidia, 221, 223, 231
Bean seed, **116**
Beard, 39
Beech, **401**
  order, 399
Bellflower, **451**
  order, 451
Berry, 365
Bidens, **456, 457**
Biennials, 59
Black knot, **215**
Blackman, 223

Bladderwort, 447
Blade, 7, 8, 20
Bloodroot, 410, 411
Blueberry, 439
Body cell, 326, 337
Bogs, 270
Boletaceae, 233
Boletus, 233
Botrychium, 285, 288
Brassica, 412
Bryales, 273
Bryophyta, 242
Buckthorn order, 428
Buds, 97, 98
Bulbs, 96
Bur-marigold, 456, 457
Buttercup, 406
  order, 404
Butternut, 402
Butterwort, 448

Cactus, 40
Calcium cyanamide, 58
Callithamnion, 190
Callus, 85
Calyptra, 271, 276, 278
Calyx, 102
Cambium, 64, 67, 68, 75
Campanula, 451
Campanulales, 451
Canal cells, 248
Canna, 388
Carbondioxide, 12, 20
Capillitium, 135, 137, 138, 234
Capsule, 249, 271, 277, 278, 279, 382, 383
Carbohydrates, 12
Carbon, 15
Carduaceae, 456
Carex, 376
Carpel, 102, 366
Carpinus, 399
Carrot, order, 432
  wild, 433
Caruncle, 119
Cassia, 419
Castor bean, 119, 63, 65, 75, 76
Catkin, 397, 398
Cat-tail, 348
  order, 368
Cedar apple, 225
Cell, division of, 48
  nature of, 1
  sap, 1, 3
Cellulose, 11
Ceramium, 190
Cercis, 419
Chaetophorales, 170
Chamaecyparis, 343
Chamaenerion, 431
Chenopodiales, 403
Chenopodium, 404
Chervil, 435
Chicory family, 453
Chlamydomonas, 159
Chlorenchyma, 11
Chlorophyceae, 155
Chlorophyll, 13
Chloroplasts, 2, 9, 12, 13, 17
Choripetalae, 396
Chromatin, 48
Chromoplasts, 2
Chromosome, 49
  reduction of, 123, 193, 253
Cilium, 136
Cladonia, 237
Classification of plants, 131
Clavaria, 232
Clavariaceae, 233
Claviceps, 217
Climbing plants, 89
Closterium, 168
Club moss fern, 309
Cluster-cup, 222
Cohesion, 352
Coenocytes, 174
Coleochaete, 179
Collecting cells, 9, 20
Collenchyma, 63, 70
Colors of flowers, 391, 408, 447
Columbine, 408
Columella, 203, 265
Comfrey, 442
Commensalism, 59

Compass plant, **29**, 30
Compositae, 452
Conducting system, 68
Cone, 306
Conidium, 205
Conn, 141
Cordyceps, **218**
Cork, 71, **72**
Cork cambium, 71, **72**
Corm, **96**, 97
Corn, root of, 50
seedling of, 120, **121**
Corolla, **102**
Cortex, **50**, 52, 62,
Corylus, **399**
Coscinodiscus, **150**
Cosmarium, **168**
Cotyledon, **110**, 112, **117**, 118, **364**
Cribraria, **135**
Crossing, 107, 128, 349
Crowfoot order, 404
Crucibulum, **236**
Cruciferae, **412**
Currant, **354**
Cuticle, 10, 23, **32**
Cutin, 10, 32
Cyanophyceae, 145, **147**
Cycadales, 321
Cycads, 321
Cycas, **323**, **328**
Cymbella, **151**
Cypripedium, **390**
Cystocarp, **192**, 193
Cytoplasm, 2

Daldinia, **216**
Dandelion, **453**, **454**
Darwin, 55, 118
Dasya, **150**
Daucus, **433**
Deciduous, 37
Delesseria, 150
Desert plants, 40
Desmidium, **168**
Desmids, **168**, **169**
De Vries, 128, 429
Diadelphous, 420
Diastase, 19, 208
Diatomaceae, 148
Diatoms, **149**
Dicotyledons, 393
Dioecious, 257
Dodder, 56, 443
Drupe, 365
Dryopteris, **294**
Duckweed, 337
Ducts, 64

Earth star, 235
Ectocarpus, 181, **182**
Elaters, 259, **260**, **307**
Elderberry, **449**
Elfvingia, **233**
Elm, **404**
Embryo, 112, 362
cell, **110**, 111, 362
sac, 111, 361, **362**
Empusa, **204**, 205
Endodermis, **52**, 63
Endosperm, **110**, 111, 341, 361, **364**
nucleus, 105, **109**, 357
Energy of plant, 13
Entomophilous, 355, 398
Enzyme, 17, 122
Epicotyl, 112, **364**, 396
Epidermis, 8, **9**, 32
Epigynous, 353, **354**
Epiphragm, 277, **278**
Epiphyte, 56
Equisetales, 303
Equisetum, **304**, **307**
Ergot, 218
Ericales, 437
Erysiphe, **213**, **214**
Erythronium, 381, **382**
Etiolation, 14
Eumycetes, 195
Euphyceae, 154
Evolution of Algae, 155, 170
of Angiospermae, 345
of Bryophyta, 281
of Eumycetes (true fungi), 199
of flower, 348
of Gymnospermae, 321

Evolution of Pteridophyta, 284
of sex, 159, 183
of sexuality, 158, 160, 172, 183
Excretion, 20, 22, 45
Exoascales, 209
Exoascus, 209

Fagales, **399**
Fagus, **401**
Fairy rings, 229
Family, 132
Farmer, 189
Fecundation, 108
Fermentation, 143, 207
Ferns, **282**
adder tongue, 284
club moss, 309
common, 291
horsetail, 303
Fertilization, 108, 361
significance of, 124
Ficus, 403
Filament, 102
Filicales, 291
Flower, 394
coloration of, 408
development of, 348
perfect and imperfect, 349
structure of, 101
Follicle, 365
Food of plants, 11, 44
distribution of, 16
storage of, 18
Foot, 258, **260**, 261
Formaldehyde, 13
Formative regions, 50, 82
Fruit, 364
Fucus, **185, 186, 187, 188**
Fungus, 195
alga-like, 197
basidia-forming, 221
bird's nest, 235, **236**
black, **215**
blue-green, 209
brown, 209
coral, **232**, 233
Fungus, cup, **219**
fly, **204**, 205
gill, **230**, 233
pore, **233**
prickly, **232**, 233
sac, 205
Funaria, **266, 272, 275, 279**
Funiculus, **104, 362**

Gager, 128
Gametangium, 172, **186**
Gamete, 104, 152, 158, 160, 168, 172, 191, 247
female, 105, **106**
male, 107, **108**
Gametophyte generation, 122, 173, 193, 244, 254, 287, 290, 299, 313, 318, 335
reduction of, 289
female, 105, **106**
male, **108, 326, 337, 359**
Gametospore, 104, 109, 180, 244
germination of, 110, 193, 249, 258, 288, 301, 309, 313, 319, 327, 338, 361
Gasteromycetes, 234
Gemmae, 243, **255, 256**
Generation, alternation of, 122, 173, 180, 193, 254, 290
Gentian, **441**
order, 440
Gentianales, 440
Genus, 131
Geoglossum, **220**
Geraniales, 427
Geranium, **428**
order, 427
Gills, 229, **230, 231**
Ginkgo, 330
Gleditsia, **419**
Gleocapsa, **147**
Grafting, **86**
Grain, **121**, 365
Graminales, 372
Grape, **428**
Grass, **373, 375, 376**
order, 372

Growth, conditions of, 115
relation of chemical changes to, 4
Growing point, **51, 82,** 243, 284, 314
Guard cells, **9, 10**
Gymnospermae, 321
Gymnosporangium, **225**

Hairs, plant, **10,** 34, **35**
root, 42, **43,** 44
Harper, 219
Haustoria, 212
Hazen, 157
Heath order, 437
Heartwood, 81
Helleborus, **408**
Helvellales, **220**
Hepaticae, 244
Hepatics, leafy, 261
thallose, 244, **245**
Heterocyst, **147**
Heterogamous, 160
Heterospory, 316, 320
Heterostyly, 430
Hilum, **116**
Honey locust, **419**
Honeysuckle, **354**
Hop, twining, 91
Hormogonium, 146, **147**
Horsetail fern, 303
Hybrid, 125
Hydnaceae, **232,** 233
Hydnum, **232**
Hydrodictyon, **164**
Hydrophytes, 35
Hymenium, 220, **231**
Hypha, 196
Hypocotyl, 112, **364,** 396
Hypocreales, 216, **217**
Hypogynous, 353
Hypoxylon, **215**
Hysteriographium, **216**

Indian-pipe, **439**
Indusium, **294, 295,** 324
Inflorescence, 349
Insectivorous plants, 269, 447
Insects, crossing by, 106, 349, 355, 379, 380, 383, 391, 443
sight and sense of smell, 384
Integument, 103, 323, **364**
Intercellular spaces, **9,** 20
Involucre, 258, 264, **433**
Iris, **386**
Ironweed, **458**
Irritability, 29, 54, 88, 91
Isogamous, 159

Jacket-cells, 325, 339
Jack-in-the-pulpit, **378**
Jeffery, 287
Juglandales, 402
Juglans, **402**
Juncus, **385**
Jungermaniales, **261**
Juniperus, 344

Keel, 419, 420
Kelps, 185
Kerner, 60
Klebs, 165
Knees of cypress, 41
Knot, 85

Labellum, **388, 390,** 391
Labiatae, 443
Laminaria, **184**
Larkspur, 409
Lateral roots, 52
Lathyrus, **420,** 421
Leaf, arrangement of, 24–28
compound, **28**
fall of, **37, 38**
movement, 28–32
reduction of, 39
structure of, 7–12, 32–37
work of, 11–24
Leaf-scar, **98**
Legume, 365
Lenticel, **73, 74**
Leotia, **220**
Lepidium, **356, 362**
Lessonia, **184**
Leucoplasts, **2,** 18

Lichens, 237–241
Licmophora, 149
Light, 4, 13–14, 17, 28, 32, 36, 55, 88
Liliales, 381
Lily, fawn, 381, 382
Lily order, 381
Linaria, 446
Linin, 48
Linnaea, 449
Liverworts, 244
  horned, 264
Living substance, 1–6
Livingston, 163
Lobelia, 451
Lodicules, 374, 376
Lomentum, 423
Loosestrife, 441
Lycogala, 135
Lycoperdales, 234
Lycoperdon, 234
Lycopodiaceae, 311
Lycopodiales, 309
Lycopodium, 310, 311, 313
Lythrum, 429, 430

MacDougal, 128
Macrocystis, 184
Madder order, 448
Maple, 424, 426
Marchantia, 247, 255–260
Marchantiales, 244
Mechanical structure, 63, 69
Medullary ray, 77, 79
Megasporangia, 316, 324, 347
Megaspore, 103, 104, 308, 316
Megasporophyll, 103, 316, 322, 347
Melandryum, 350
Melosira, 149
Mendel, 127
Mesophyll, 9, 10
Mesophytes, 35
Micropyle, 103, 104, 323
Microsphaera, 214
Microsporangia, 316, 347
Microspore, 102, 308, 316, 317
  germination of, 107, 108, 307, 317, 326, 337
Micrasterias, 168
Mildews, downy, 199
  powdery, 212, 213
Minerals in soils, 44
Mint family, 443
Monocotyledons, 82, 367
Monoecious, 257
Monotropa, 439
Moore, 58
Morchella, 220
Morel, 220
Morning-glory, 442
Mosaics, 27
Moss, 267
  bog or peat, 268
  true, 273
Mould, black, 202, 203
  brown, 209
  green, 196, 206, 209
  water, 197
Movements, gases, 12, 20, 41
  leaf, 29
  root, 54
  stem, 87, 90
  water, 23, 52, 64, 67, 68
Mucorales, 202
Mushrooms, 227, 230
Musci, 267
Mustard family, 412
Mutants, 128
Mycelium, 196
Mycorrhiza, 59, 269, 287, 390
Myrtales, 429
Myrtle order, 429
Myxomycetes, 134, 135

Narcissus, 386
Navicula, 149
Nectar glands, 348, 355, 371, 398, 408
Nemalion, 191
Nereocystis, 185
Nidulariales, 235
Netted veined leaf, 293
Nettle, 35, 403
Nitrogen, 57, 142
Nostoc, 147

Nucellus, 103, **104**, 356
Nucleoli, **48**
Nucleus, **1**, **2**, 48, 124
Nut, 400
Nutations, 87, 92

Oak, 28, **401**
Odors, 379, 384, 400
Oedogonium, **177**, **178**
Oenothera, 27, **43**
Onoclea, **299**
Oögonium, 176
Oösphere, 176, 248
Operculum, **271**, 272
Ophioglossales, 284
Ophioglossum, **285**
Orange, 427
Orchid order, 389
Orchidales, **389**
Orchis, **392**
Order, 132
Oscillatoria, **147**
Osmosis, 45
Osmunda, **296**, **297**
Ovary, **102**, 103, 347
Ovule, **102**, 103, **104**, 347, 366
Oxygen, 3, 13, 15, 20

Palisade mesophyll, 10, 36
Pallavicinia, **261**
Palms, 272
Pandanales, 368
Pandorina, **159**, 160
Papaverales, 410
Papillae of stigma, **356**, 359
Papilionaceae, 420
Pappus, 453, **454**
Paraphyses, **188**, **275**, 276
Parasites, 56, 195
Parenchyma, 11, 64
Parmelia, **237**
Parnassia, **415**
Pasteur, 145
Pea, family, 420
  sweet, **420**
Peach curl, 209
Peat moss, 269
Pediastrum, **164**
Penicillium, **196**, **206**, 209, **210**
Perennials, 60
Perianth, 101, **260**, 264, 346, 350
Periderm, 234, **235**
Perigynous, **352**, 353
Perisperm, 365
Perisporiales, 212–214
Peristome, 277, **278**
Perithecia, 206, **211**
Peronospora, **199**, **200**
Peronosporales, 199
Petals, 101, **350**
Petiole, **7**, 8
Peziza, **218**
Pezizales, **219**
Phaeophyceae, 181
Phallales, 236
Phallus, **236**
Phloem, 52, 64, **65**, **66**, 67
Phosphorescence, 299
Phlox order, 442
Photosynthesis, 11
Phycomycetes, 197
Phyllactinia, **214**
Phylloglossum, **312**
Phytophthora, 199, **200**
Picea, **343**
Pierce, 170
Pileus, 229, **230**
Pinales, 330
  classification of, 343
Pine, 331, **333**, **334**, **336**
Pinguicula, 448
Pinnularia, **150**, **153**
Pistil, **102**, 366
Pith, 64
Placenta, 355
Plankton, 148
Planktoniella, **150**
Plant, nature of, 1
  cell, 1
Plasmodium, 136, **137**
Plasmopara, **200**, 201
Plastids, **2**, 17
Pleurococcus, **162**
Plowrightia, **215**

Plumule, 116, **364**, 396
Pod, **116**, 365, **423**
Polar nucleus, 105, **106**
Polemoniales, 442
Pollen grains, 102
Pollinium, 391, **392**, 441
Polygonales, 403
Polypodium, **298**
Polyporaceae, 233
Polysiphonia, **192**, 194
Polytrichum, **274**, **275**, **278**, **282**
Pond scum, 166
Poplar order, 396
Poppy order, 410
Pore, 11, 79, 331
Porella, **262**
Potato, **448**
  rot, 199
Primrose order, 440
Primulales, 440
Pro-embryo, **327**, 339, 362, 364
Prosenchyma, 11
Proteids, 16
Protandrous, 349
Protection of microspores, 379
Proteid grains, **18**, **19**
Prothallium, **283**, 299, **300**
Protogynous, 349
Protonema, **280**
Protoplasm, function of, 3
  nature of, 2, 3
Protoplasmic connection of cells, 3
Pteridophyta, 282
Pteridium, **298**
Ptomaines, 143
Puccinia, 221, **224**
Puff ball, 234, **235**
Pulvinus, 30
Pustules, 226, **227**
Pycnidium, pycnidia, **222**
Pyrenoids, **165**, 166
Pyrola, **351**, **437**
Pyronema, **219**

Quercus, **350**, 399, **401**

Ragweed family, 458
Ranales, 404
Ranunculus, **406**
Ray, flower, 456
  medullary, 79
Rattlebox, yellow, **446**, 447
Receptacle, 348–353
Red algae, 154, 189–194
Red snow, 157
Respiration, 20, 46
Reproduction, 101, 124
Rhamnales, 428
Resin, 332
Resurrection plant, 314
Rhinanthus, **446**, 447
Rhizoids, 242
Rhizomes, 94, **95**, **373**
Rhizopus, **202**, **203**
Rhodophyceae, 189, **190**
Rhus, **427**
Riccia, **245**
Ricciocarpus, **245**, **246**, **254**
Richards, 100
Ring, annual, 77
Rivularia, **147**
Root, 1, 4, 42, 265, 283
  anchoring and supporting, 60
  bacteria on, **57**
  contraction of, 60
  lateral, **53**
  modification of, 56
  sensitiveness of, 54
  structure of, **50**, **51**, **52**
  surface of, 47, 48
Roots as storage organs, 59
Root cap, **50**, **364**
  hairs, 42, **43**, 44
Rootstock, 94
Rosales, 414
Rosette, 27
Rose order, 414
Rostellum, 391, **392**
Rubiales, 448
Rusts, 221
  white, 199

Saccharomycetes, 206
Sac fungi, 205–221

Sage, **444**
Salicales, 396
Salix, **348**, 397
Salvia, **444**
Samara, 365, **426**
Sambucus, **449**
Sanguinaria, 410
Sap, 3, 52, 62, 68
Sapindales, 424
Saprolegnia, 197, **198**
Saprolegniales, 197
Saprophytes, 137
Sapwood, 81
Sargassum, **185**
Saxifrage, **354**
Scabiosa, **450**
Scenedesmus, **162**
Schizocarp, **426**
Schizophyta, 139
Scitaminales, 387
Sclerenchyma, 63
Sclerotium, 218
Scouring rush, 305
Scrophulariaceae, 445
Scutellaria, **444**
Scutellum, **121**, 122, **376**
Secondary growth, 71
Sedge order, 372
Sedum, **102**, **346**, **415**
Seed, 320, 328, **340**, **364**
  development of, **111**, 112
  growth of, 113, **117**
Seedling, castor bean, **119**
  corn, 121
Seed plant, 320
Selaginella, 314, 315, 317
Selaginellaceae, 314
Sensitive plant, 30, 31
Sepals, 350
Seta, 264, **278**
Sex, evolution of, 158, 165, 173, 177, 179, 183, 187
Shade plants, 32–36
Shull, 126
Sieve tubes, **65**, **66**
Siphonales, 174
Skull-cap, **444**
Skunk cabbage, 379
Sleep movements, 30
Slime mould, 134, **135**, **137**
Smuts, 226, **227**
Soapberry order, 424
Soils, nature of, 44, 46
Solanum, **448**
Soredium, **239**, 240
Sorus, 286, **294**
Spadix, **378**
Spathe, **378**
Spathyema, 379
Species, 131
Sperm, 176, 248
Spermatophyta, 320
Sphaerella, 155, **156**
Sphaeriales, **215**
Sphagnales, 268
Sphagnum, **269**, **271**
Spirillum, **140**
Spirogyra, 166, **167**
Spongy mesophyll, **9**, 11, 20
Sporangia, 135, **285**, 286, **294**
Spore, 104
  differentiation of, 308
  germination of, **251**, 307, 317
Sporophylls, 101, 294, 306, **346**
Sporophyte generation, 122, 173, 180, 193, 254
Sporophyte, development of, **265**, 281, 284, 289, 313, 327, 338
Stamen, **102**, 366
Staminodia, **388**
Starch, **2**, 12, 16, 18, 19
  test for, 14
Staurastrum, **168**
Steironema, **441**
Stem, dicotyledon, **75**, **78**
  function of, 62
  growth of, **75**, **76**, **82**
  healing of, 85
  modifications of, 89
  monocotyledon, **70**, 82
  sensitiveness of, 88
  structure of, 62, **63**, **70**, **78**
Stemonitis, **135**
Stereome, **32**, 63, **64**

Stereum, 232
Sticta, 237
Stigma, 102, 106, 356
Stink horns, 236
Stimuli, effects of, 4
  reaction to, 5
Stipe, 229, 230
Stipules, 393
Stolon, 94
Stoma, 8, 9, 10, 20, 24, 32, 33
Stone fruit, 365
Storage organs, 18, 59
Strawberry, 351, 417
Strobilus, 304, 306, 309, 341
Stroma, 215
Style, 103
Suber, 72
Sugar, 12, 16
Surface reduction, 40
Suspensor, 110, 111, 314, 362
Symbiosis, 238
Sympetalae, 436
Symphytum, 352
Synergid, 105, 106, 357, 362

Tabellaria, 149
Taraxacum, 453, 454
Thaxter, 138
Taxus, 343
Teleutospore, 223, 224
Telial stage, 223
Telophoraceae, 232
Tendrils, 91, 92
Tetraspores, 191
Thallophyta, classification of, 134
Thallus, 255, 256
Thistle family, 456
Thuja, 343
Tissue, strengthening, 69
Toadflax, 446
Toadstools, 227, 229
Toxins, 143
Tracheids, 9
Transpiration, 22
Transport of water, 52
Tree ferns, 293
Truffles, 211
Tube cell, 107, 108, 326, 338
Tubercles, 57
Tubers, 95
Turgor, 69
Twigs, 98
Twining stems, 90
Typha, 348, 368, 370

Ulothrix, 170, 171, 172
Umbel, 433, 434
Umbellales, 432
Uncinula, 213, 214
Uredinial stage, 223
Uredinales, 221
Uredospore, 223, 224
Urticales, 403
Ustilaginales, 226
Ustilago, 227
Utricularia, 447

Vaccinium, 439
Vacuoles, 1, 2, 51, 196
Valerian, 450
Variation, 89, 124–129, 157, 165, 170
Variegated plants, 14
Vascular bundles, 11, 12, 16, 64, 65, 66, 81, 293
Vaucheria, 174, 175, 176
Veins, 11, 111, 268, 283
Velamen, 390
Veltheimia, 369
Velum, 228, 229
Veratrum, 385
Vernonia, 458
Vessels, 64, 66
Viburnum, 449
Volva, 229, 230
Volvocales, 155
Volvox, 160, 161

Wall cells, 247, 287, 318, 325, 337, 361
  formation of, 48–50
Ward, 23
Water, capillary, 44
  hygroscopic, 44
  lily, 365, 405, 407

Weeds, 403
Willow, **348**
  order, 396
Wilting, cause of, 22
Winged fruits, 365
Witches' broom, 226
Wood, heart, 81
  oak, 80
  pine, 331
  sap, 81
  sections, 78, 79
  spring, 77
  summer, 77
Woodwardia, **295**
Work of plants, 15

Xanthidium, 168
Xerophytes, 35, 40
Xylaria, **216**
Xylem, 52, 64, **65**

Yamanouchi, 194
Yeast, 206, **207**
Yucca, 386

Zamia, **322, 323, 324, 326**
Zoöspore, 136
Zygnema, **165**, 167
Zygnematales, 165
Zygomorphic, **354**

105960
Author Curtis, Carlton Clarence
Title Nature and development of plants.
Bot
C
A

University of Toronto
Library

DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET

Acme Library Card Pocket
Under Pat. "Ref. Index File"
Made by LIBRARY BUREAU